

LATTICE EMBEDDINGS IN PERCOLATION

BY GEOFFREY R. GRIMMETT AND ALEXANDER E. HOLROYD

*Cambridge University and Microsoft Research,
and University of British Columbia*

Does there exist a Lipschitz injection of \mathbb{Z}^d into the open set of a site percolation process on \mathbb{Z}^D , if the percolation parameter p is sufficiently close to 1? We prove a negative answer when $d = D$ and also when $d \geq 2$ if the Lipschitz constant M is required to be 1. Earlier work of Dirr, Dondl, Grimmett, Holroyd and Scheutzwow yields a positive answer for $d < D$ and $M = 2$. As a result, the above question is answered for all d, D and M . Our proof in the case $d = D$ uses Tucker's lemma from topological combinatorics, together with the aforementioned result for $d < D$. One application is an affirmative answer to a question of Peled concerning embeddings of random patterns in two and more dimensions.

1. Introduction and results.

1.1. *Preliminaries.* Let \mathbb{Z}^d denote the d -dimensional integer lattice. Elements of \mathbb{Z}^d are called *sites*. Let $\|\cdot\|_r$ denote the ℓ^r -norm on \mathbb{Z}^d , and abbreviate $\|\cdot\|_1$ to $\|\cdot\|$. We say that a map $f: \mathbb{Z}^d \rightarrow \mathbb{Z}^D$ is M -Lipschitz, or M -Lip, if $\|f(x) - f(y)\| \leq M$ for all $x, y \in \mathbb{Z}^d$ with $\|x - y\| = 1$.

For $p \in [0, 1]$, consider the site percolation model on \mathbb{Z}^D . That is, declare each site to be *open* (or p -open) with probability p , and otherwise *closed*, with different sites receiving independent designations. Let $W_p(\mathbb{Z}^D)$ denote the random set of open sites, and write \mathbb{P}_p and \mathbb{E}_p for the associated probability measure and expectation operator.

We are interested primarily in the probability

$$(1) \quad L(d, D, M, p) := \mathbb{P}_p(\exists \text{ an } M\text{-Lip injection from } \mathbb{Z}^d \text{ to } W_p(\mathbb{Z}^D)).$$

Clearly L is increasing in D, M and p , and decreasing in d . Furthermore, L is $\{0, 1\}$ -valued, since \mathbb{P}_p is a product measure and the event in (1) is invariant under translations of \mathbb{Z}^D . We define the critical probability

$$p_c(d, D, M) := \inf\{p : L(d, D, M, p) = 1\}$$

and furthermore

$$M_c(d, D) := \min\{M \geq 1 : p_c(d, D, M) < 1\}$$

Received March 2010; revised September 2010.

AMS 2000 subject classification. 60K35.

Key words and phrases. Lipschitz embedding, lattice, random pattern, percolation, quasi-isometry.

Theorem 1(a) is an immediate consequence of a substantially stronger statement proved in [2], which we state next. For $x = (x_1, \dots, x_{d-1}) \in \mathbb{Z}^{d-1}$ and $z \in \mathbb{Z}$ we write $(x, z) := (x_1, \dots, x_{d-1}, z) \in \mathbb{Z}^d$. Write $\mathbb{Z}_+ := \mathbb{Z} \cap (0, \infty)$.

THEOREM 2 (Lipschitz percolation [2]). *Let $d \geq 2$, and suppose $p > 1 - (2d)^{-2}$. There exists \mathbb{P}_p -a.s. a (random) 1-Lip function $F : \mathbb{Z}^{d-1} \rightarrow \mathbb{Z}_+$ such that for every $x \in \mathbb{Z}^{d-1}$, the site $(x, F(x)) \in \mathbb{Z}^d$ is open.*

With F given as in Theorem 2, the map $x \mapsto (x, F(x))$ is evidently a 2-Lip injection, thus establishing Theorem 1(a). Other applications of Theorem 2 appear in [3, 6]. Further properties of F are explored in [5], where an improved bound on the value of p in Theorem 2 is given.

The proof of Theorem 1(b) is relatively straightforward and may be found in Section 2. (The proof involves showing that any 1-Lip injection from \mathbb{Z}^2 to the full lattice \mathbb{Z}^D must satisfy rather rigid conditions.) The principal contribution of the current paper is Theorem 1(c). Interestingly, our proof of this nonexistence result makes use of the above existence result, Theorem 2. Another essential ingredient of this proof is Tucker’s lemma from topological combinatorics (see [10, 12]).

It is immediate from the definition of $M_c(d, D)$ that, if $M_c(d, D) = \infty$, then $p_c(d, D, M) = 1$ for all $M \geq 1$. On the other hand, we have the following result when $M_c(d, D) < \infty$ [which occurs if and only if $d < D$, as noted in (2) above].

PROPOSITION 3. *Let d, D be positive integers such that $M_c(d, D) < \infty$. Then $p_c(d, D, M) \rightarrow 0$ as $M \rightarrow \infty$.*

1.3. Embeddings of patterns. The above results concerning maps from \mathbb{Z}^d to the open sites of \mathbb{Z}^D have implications in the more general setting of maps that preserve values indexed by \mathbb{Z}^d , as follows. Let $\Omega_d := \{0, 1\}^{\mathbb{Z}^d}$ be the space of percolation configurations, in which the value 1 (resp., 0) is identified with the state “open” (resp., “closed”). An *embedding* of a configuration $\eta \in \Omega_d$ into a configuration $\omega \in \Omega_D$ is an injection $f : \mathbb{Z}^d \rightarrow \mathbb{Z}^D$ such that $\eta(x) = \omega(f(x))$ for all $x \in \mathbb{Z}^d$. We call a configuration $\eta \in \Omega_d$ *partially periodic* if there exist $x \in \mathbb{Z}^d$ and $r \in \mathbb{Z}_+$ such that $\eta(x) = \eta(x + ry)$ for all $y \in \mathbb{Z}^d$.

PROPOSITION 4 (Embedding). *Let d, D be positive integers.*

(a) *Let $d \geq 2$, $p \in (0, 1)$ and $\eta \in \Omega_d$. For \mathbb{P}_p -a.e. $\omega \in \Omega_D$, there exists no 1-Lip embedding of η into ω .*

(b) *Let $d < D$. For every $p \in (0, 1)$, there exists $M \geq 1$ such that for \mathbb{P}_p -a.e. $\omega \in \Omega_D$, it is the case that for every $\eta \in \Omega_d$, there exists an M -Lip embedding of η into ω .*

(c) *Let $d = D$ and let $\eta \in \Omega_d$ be a partially periodic configuration. For every $p \in (0, 1)$, $M \geq 1$ and for \mathbb{P}_p -a.e. $\omega \in \Omega_D$, there exists no M -Lip embedding of η into ω .*

The current work was motivated in part by the problem of Lipschitz embeddings of random one-dimensional configurations (see [4, 7]). Proposition 4(a) extends Theorem 1(b) to more general configurations than the all-1 configuration. Part (b) answers affirmatively a question posed by Peled concerning the existence of M -Lip embeddings of d -dimensional random configurations into spaces of higher dimension; see [4], Section 5. Part (c) leaves unanswered the question of whether or not there exist $d \geq 1$, $p \in (0, 1)$, $\eta \in \Omega_d$ and $M < \infty$ such that with strictly positive probability (and therefore probability 1), there exists an M -Lip embedding from η into a random configuration $\omega \in \Omega_d$ having law \mathbb{P}_p .

1.4. *Quasi-isometry.* There is a close connection between the existence of embeddings and of quasi-isometries. A *quasi-isometry* between two metric spaces (X, δ) and (Y, ρ) is a map $f : X \rightarrow Y$ such that there exist constants $c_i \in (0, \infty)$ with:

- (a) $\forall x, x' \in X, c_1\delta(x, x') - c_2 \leq \rho(f(x), f(x')) \leq c_3\delta(x, x') + c_4,$
- (b) $\forall y \in Y, \exists x \in X$ such that $\rho(f(x), y) \leq c_5.$

We call such f a ***c-quasi-isometry*** when we wish to emphasize the role of the vector $\mathbf{c} = (c_1, \dots, c_5)$. It is not difficult to see that the existence of a quasi-isometry is a symmetric relation on metric spaces. Quasi-isometries of *random* metric spaces are discussed in [11]. A *subspace* of a metric space (X, δ) is a metric space (U, δ) with $U \subseteq X$.

PROPOSITION 5 (Quasi-isometry). *Let d, D be positive integers, and let E be the event that there exists a quasi-isometry between (\mathbb{Z}^d, ℓ^1) and some subspace of $(W_p(\mathbb{Z}^D), \ell^1)$.*

- (a) *If $d < D$ then for all $p \in (0, 1)$ we have $\mathbb{P}_p(E) = 1$.*
- (b) *If $d \geq D$ then for all $p \in (0, 1)$ we have $\mathbb{P}_p(E) = 0$.*

The proofs of Theorem 1(b), (c) appear, respectively, in Sections 2 and 3. The remaining propositions are proved in Section 4. Section 5 contains four open problems.

2. Nearest-neighbor maps. In this section we prove Theorem 1(b). A (self-avoiding) *path* in \mathbb{Z}^d is a finite or infinite sequence of distinct sites, each consecutive pair of which is at ℓ^1 -distance 1. Let $e_1, \dots, e_d \in \mathbb{Z}^d$ be the standard basis vectors of \mathbb{Z}^d , and let 0 denote the origin.

LEMMA 6. *Let $x_1, \dots, x_k \in \mathbb{Z}^D$ be distinct, and let $A = A(x_1, \dots, x_k)$ be the event that there exists a singly-infinite path $0 = y_0, y_1, \dots$ in \mathbb{Z}^D such that the sites $(x_i + y_j : i = 1, \dots, k, j = 0, 1, \dots)$ are distinct and open. If $p < (2D)^{-1/k}$, then $\mathbb{P}_p(A) = 0$.*

PROOF. Let A_n be the event that there exists a path $0 = y_0, y_1, \dots, y_n$ of length n in \mathbb{Z}^D such that the sites $(x_i + y_j : i = 1, \dots, k, j = 0, \dots, n)$ are distinct and open. Note that A is the decreasing limit of A_n as $n \rightarrow \infty$. Let N_n be the number of paths $0 = y_0, \dots, y_n$ with the properties required for A_n . Then

$$\mathbb{P}_p(A_n) \leq \mathbb{E}_p N_n \leq (2D)^n p^{nk} \xrightarrow{n \rightarrow \infty} 0 \quad \text{if } 2Dp^k < 1.$$

Here, $(2D)^n$ is an upper bound for the number of n -step self-avoiding paths (y_j) starting from 0, while for those paths for which the sites $x_i + y_j$ are distinct, p^{nk} is the probability they are all open. \square

PROOF OF THEOREM 1(b). We must prove that, for any fixed $p < 1$ and $D \geq 2$, a.s. there exists no 1-Lip injection from \mathbb{Z}^2 to $W_p(\mathbb{Z}^D)$.

First, suppose f is a 1-Lip injection from \mathbb{Z}^2 to the full lattice \mathbb{Z}^D , and consider the image of a unit square. Specifically, take $(i, j) \in \mathbb{Z}^2$ and let

$$\begin{aligned} r_1 &= f(i+1, j) - f(i, j), & r'_1 &= f(i+1, j+1) - f(i, j+1), \\ r_2 &= f(i, j+1) - f(i, j), & r'_2 &= f(i+1, j+1) - f(i+1, j). \end{aligned}$$

Note that the four vectors r_1, r_2, r'_1, r'_2 are elements of $\{\pm e_j : j = 1, \dots, D\}$ (by the 1-Lip property); they satisfy $r_1 + r'_2 = r'_1 + r_2$ (by definition); the pair r_1, r_2 are neither equal to nor negatives of each other and similarly for r'_1, r'_2 (a consequence of injectivity). It follows that $r_1 = r'_1$ and $r_2 = r'_2$. Since this holds for every unit square, for any distinct $i, i' \in \mathbb{Z}$, the images under f of the two paths $\{(i, j) : j \in \mathbb{Z}\}$ and $\{(i', j) : j \in \mathbb{Z}\}$ are two disjoint self-avoiding paths that are translates of each other. (Another consequence, which we shall not need, is that there exists $\Delta \subset \{1, \dots, D\}$ such that all horizontal edges have images in $\{\pm e_j : j \in \Delta\}$, and all vertical edges have images in the complement $\{\pm e_j : j \notin \Delta\}$.)

Let B be the event that there exist $x_1, x_2, \dots \in \mathbb{Z}^D$ and a self-avoiding path $0 = y_0, y_1, \dots$ in \mathbb{Z}^D such that the sites $(x_i + y_j : i \geq 1, j \geq 0)$ are distinct and open. The above argument implies that, if there exists a 1-Lip injection $f : \mathbb{Z}^2 \rightarrow W_p(\mathbb{Z}^D)$, then B occurs. We shall now show that $\mathbb{P}_p(B) = 0$ for all $p < 1$ and $D \geq 1$. Let k be large enough that $p < (2D)^{-1/k}$. We define B_k analogously to B , except in that we now require the existence of only k sites x_1, \dots, x_k . Lemma 6 implies that $\mathbb{P}_p(B_k) = 0$ because B_k is the countable union over all possible x_1, \dots, x_k of the events $A(x_1, \dots, x_k)$. Finally, we have $B \subseteq B_k$. \square

3. The case of equal dimensions. In this section we prove Theorem 1(c). We denote integer intervals by $\llbracket a, b \rrbracket := (a, b] \cap \mathbb{Z}$, et cetera. Fix any $d \geq 2$, $M \geq 1$ and $p \in (0, 1)$. We will prove that a.s. there exists no M -Lip injection from \mathbb{Z}^d to $W_p(\mathbb{Z}^d)$.

The idea behind the proof is as follows. Suppose that f is such an injection. By a *hole* we mean a cube of side length M in \mathbb{Z}^d all of whose sites are closed (actually, a slightly different definition will be convenient in the formal proof, but

this suffices for the current informal sketch). Holes are rare (if p is close to 1), but the typical spacing between them is a fixed function of d , M and p . We will consider the image under f of a cuboid $\llbracket 1, n \rrbracket^{d-1} \times \llbracket 1, m \rrbracket \subset \mathbb{Z}^d$, where $m \gg n \gg 1$. We will arrange that the images of the two opposite faces $\llbracket 1, n \rrbracket^{d-1} \times \{1\}$ and $\llbracket 1, n \rrbracket^{d-1} \times \{m\}$ are far apart, and separated by a $(d-1)$ -dimensional “surface of holes” (at the typical spacing). This implies that the image of the interior of the cuboid must pass through this surface, avoiding all the holes. To do so, the image must be in some sense be folded up so as to be locally $(d-1)$ -dimensional, and this will give a contradiction to the injectivity of f if n is chosen large enough compared with the spacing of the holes.

In the case $d = 2$, it is possible to formalize the above ideas using fairly direct ad hoc geometric arguments. It is plausible that a similar approach could be pushed through (with substantially more difficulty) for $d = 3$. For general d , a less direct (but more systematic) approach appears to be required. Specifically, the surface of holes will be constructed using Theorem 2, and, crucially, we will augment it with a coloring of the nearby open sites using exactly $d-1$ colors, in such a way that the colored sites separate the two sides of the surface from each other, but the sites of any given color fall into well-separated regions of bounded size. Via the map f , this coloring will induce a coloring of the cuboid that contradicts a certain topological fact.

The following notation will be used extensively. A *coloring* of a set of sites $U \subseteq \mathbb{Z}^d$ is a map χ from U to a finite set Q . A site $u \in U$ is said to have *color* $\chi(u) \in Q$. We introduce the graph $G(U, \ell^r, k)$ having vertex set U and an edge between $u, v \in U$ if and only if $0 < \|u - v\|_r \leq k$. An important special case is the *star-lattice* $G^* = G_d^* := G(\mathbb{Z}^d, \ell^\infty, 1)$. Given a graph G and a coloring χ of its vertex set, a *q -cluster* (of χ with respect to G) is the vertex set of a connected component in the subgraph of G induced by the set of vertices of color q . The *volume* of a cluster is defined to be the number of its sites.

We next state the two main ingredients of the proof: a topological result on coloring a cuboid and a result on existence of random colored surfaces in the percolation model.

PROPOSITION 7 (Color blocking). *Let d, n, m be positive integers, and consider a coloring*

$$\chi : \llbracket 1, n \rrbracket^{d-1} \times \llbracket 1, m \rrbracket \rightarrow \{-\infty, +\infty, 1, 2, \dots, d-1\}.$$

If χ satisfies:

- (a) *all sites in $\llbracket 1, n \rrbracket^{d-1} \times \{1\}$ have color $-\infty$;*
- (b) *all sites in $\llbracket 1, n \rrbracket^{d-1} \times \{m\}$ have color $+\infty$; and*
- (c) *no site of color $+\infty$ is adjacent in G^* to a site of color $-\infty$,*

then, for some $j \in \{1, 2, \dots, d-1\}$, χ has a j -cluster with respect to G^ of volume at least n .*

In fact, Proposition 7 remains valid if “volume” is replaced with “diameter,” as we shall see.

PROPOSITION 8 (Colored surfaces). *Fix $d \geq 2$, $J \geq 1$ and $p \in (0, 1)$. There exist constants $K, C < \infty$ (depending on d, J and p) such that \mathbb{P}_p -a.s. there is a (random) coloring*

$$\lambda: W_p(\mathbb{Z}^d) \rightarrow \{-\infty, +\infty, 1, 2, \dots, d-1\}$$

of the open sites of \mathbb{Z}^d with the following properties:

(a) No site of color $+\infty$ is adjacent to a site of color $-\infty$ in $G(W_p(\mathbb{Z}^d), \ell^\infty, J)$.

(b) For each $j \in \{1, 2, \dots, d-1\}$, every j -cluster with respect to $G(W_p(\mathbb{Z}^d), \ell^\infty, J)$ has volume at most K .

(c) There exists a (random) nonnegative real-valued function $g: \mathbb{Z}^{d-1} \rightarrow [0, \infty)$, with the Lipschitz property that $|g(u) - g(v)| \leq d^{-1} \|u - v\|_1$ for all $u, v \in \mathbb{Z}^{d-1}$, such that all open sites in

$$S^- := \{(u, z) : u \in \mathbb{Z}^{d-1}, z < g(u)\}$$

are colored $-\infty$, while all open sites in

$$S^+ := \{(u, z) : u \in \mathbb{Z}^{d-1}, z > g(u) + C\}$$

are colored $+\infty$.

In (c) above, note in particular that all open sites in the half-space $\mathbb{Z}^{d-1} \times (-\infty, 0)$ are colored $-\infty$.

The proof of Theorem 1(c) will proceed by playing Propositions 7 and 8 against one another to obtain a contradiction. The number of permitted colors is crucial—if one color more were added to $1, \dots, d-1$, then the conclusion of Proposition 7 would no longer hold, while with one color fewer, the conclusion of Proposition 8 would not hold. It should also be noted that the use of the star-lattice G^* is essential in Proposition 7—the statement does not hold for the nearest-neighbor lattice $G(\mathbb{Z}^d, \ell^1, 1)$. Another key point is that the presence of closed sites is essential for Proposition 8—the conclusion does not hold when $p = 1$ (for any K, C), since such a coloring would give a contradiction to Proposition 7.

The choice of the Lipschitz constant d^{-1} in Proposition 8(c) above is relatively unimportant—the result would hold for any positive constant, while any constant less than $(d-1)^{-1}$ would suffice for our application (see Lemma 11 below).

Our proof of Proposition 7 will use Tucker’s lemma, a beautiful result of topological combinatorics. The general version of [8, 12] applies to triangulations of a ball, and is a close relative of the Borsuk–Ulam theorem (see [10] for background). We need only a special case, for the cuboid, which is also proved in [1].

For $t \in \llbracket 1, \infty \rrbracket^d$, consider the integer cuboid $T = T(t) := \llbracket 0, t_1 \rrbracket \times \cdots \times \llbracket 0, t_d \rrbracket \subset \mathbb{Z}^d$ with opposite corners 0 and t , and define the boundary $\partial T := T \setminus [\llbracket 0, t_1 \rrbracket \times \cdots \times \llbracket 0, t_d \rrbracket]$. We say that boundary sites $x, y \in \partial T$ are *antipodal* if $x + y = t$.

LEMMA 9 (Tucker's lemma for the cuboid [1]). *Let $T \subset \mathbb{Z}^d$ be a cuboid as above, and suppose $\beta: T \rightarrow \{\pm 1, \dots, \pm d\}$ is a coloring such that for each antipodal pair $x, y \in \partial T$ we have $\beta(x) = -\beta(y)$. Then there exist $u, v \in T$ that are adjacent in G^* (and, in fact, that satisfy $u_i \leq v_i \leq u_i + 1$ for all i) such that $\beta(u) = -\beta(v)$.*

PROOF OF PROPOSITION 7. Throughout the proof, adjacency and clusters refer to G^* . The (ℓ^∞ -)diameter of a cluster is the maximum ℓ^∞ -distance between two of its sites. It suffices to show that for a coloring χ satisfying the given conditions, there is a j -cluster of diameter at least n for some $j \neq \pm\infty$. Suppose that this is false. We will construct a modified coloring that leads to a contradiction.

First define a coloring χ' of the larger cuboid $T := \llbracket 0, n + 1 \rrbracket^{d-1} \times \llbracket 0, m + 1 \rrbracket$ as follows. Let χ' agree with χ on $T \setminus \partial T$, except with color ∞ everywhere changed to d , and $-\infty$ changed to $-d$. Color ∂T as follows. For each $i = 1, \dots, d$, let χ' assign color $-i$ to the face $\{x \in T : x_i = 0\}$, and color $+i$ to the antipodal face (this rule creates conflicts at the intersections of faces; for definiteness assign such sites the candidate color of smallest absolute value). Thus χ' satisfies the condition of Lemma 9 on the boundary.

Now let β be the coloring of T obtained by modifying χ' as follows. For each $i = 1, \dots, d - 1$, recolor with color $-i$ all i -clusters that are adjacent to the face colored $-i$. Since there were no i -clusters of diameter as large as n in χ , this does not affect the colors on ∂T . Hence Lemma 9 applies, so there are adjacent sites $u, v \in T$ with $\beta(u) = -\beta(v)$, which contradicts the manner of construction of β . \square

The proof of Proposition 8 relies on Theorem 2 concerning Lipschitz surfaces in percolation, together with the following deterministic fact.

LEMMA 10 (Periodic coloring). *For any integers $d \geq 1$ and $R \geq 2d$, there exists a coloring $\alpha: \mathbb{Z}^d \rightarrow \{0, 1, \dots, d\}$ with the following properties:*

- (a) *The coloring is periodic with period R in each dimension; that is, $\alpha(x + Ry) = \alpha(x)$ for all $x, y \in \mathbb{Z}^d$.*
- (b) *For each $j \in \{0, 1, \dots, d\}$, every j -cluster with respect to G^* has volume at most R^d .*
- (c) *The 0-clusters with respect to G^* are precisely the cubes $Rx + \llbracket -(d - 1), (d - 1) \rrbracket^d$, for $x \in \mathbb{Z}^d$.*




FIG. 1. Part of the coloring α of Lemma 10, for $d = 1$ (top), $d = 2$ (middle), $d = 3$ (bottom; color 3 is shown as transparent, and only selected slabs are shown).

PROOF. The construction is illustrated in Figure 1. Define a *slice* to be any set of sites of the form $Y = Rx + (I_1 \times \cdots \times I_d)$, where $x \in \mathbb{Z}^d$ and each I_i is either $\{0\}$ or $[[1, R - 1]]$. If $[[1, R - 1]]$ appears k times in this product then we call Y a k -*slice*. The set of all slices forms a partition of \mathbb{Z}^d . Let $a_k := d - 1 - k$. For a k -slice Y , define the associated k -*slab* to be the set obtained from Y by replacing each occurrence of $\{0\}$ in the product $I_1 \times \cdots \times I_d$ with $[[-a_k, a_k]]$ (thus “thickening” the slice by distance a_k). We now define the coloring: for each site x , let $\alpha(x)$ be the smallest k for which x lies in some k -slab.

The required properties (a) and (c) are immediate [the cubes in (c) are precisely the 0-slabs]. For (b), note that any k -cluster is contained within a single k -slab; it is straightforward to check that, for any given k , any connection in G^* between two

different k -slabs is prevented by sites of smaller colors (here it is important that a_k is strictly decreasing in k). The volume of a k -slab is $(R - 1)^k (2a_k + 1)^{d-k} < R^d$. □

In the following, we sometimes refer to the d coordinate as vertical, with positive and negative senses being up and down, respectively, and the other coordinate directions as horizontal.

PROOF OF PROPOSITION 8. See Figure 2 for an illustration of the construction. Let L be a large constant, a multiple of J , to be determined later, and let α be the coloring from Lemma 10 with parameters $d - 1$ and $R := L/J$. Let α' be the coloring obtaining by dilating α by a factor J ; that is, for $u \in \mathbb{Z}^{d-1}$, let $\alpha'(u) = \alpha(\lfloor u/J \rfloor)$ (where $\lfloor v \rfloor$ denotes v with each co-ordinate rounded to the nearest integer, rounding up in the case of ties). Note from property (a) in Lemma 10 that α' has period L in each dimension, while by (b), for $j \in \{0, 1, \dots, d - 1\}$, each j -cluster of α' with respect to $G(\mathbb{Z}^{d-1}, \ell^\infty, J)$ has volume at most L^{d-1} . Write




FIG. 2. Part of the random coloring λ of Proposition 8, for $d = 2$ (top), and $d = 3$ (bottom, with colors $\pm\infty$ shown as transparent). Holes are shown as black.

$r := J(2d - 3)$. From Lemma 10(c), the 0-clusters of α' are $(d - 1)$ -dimensional cubes of side length r centred (approximately) at the elements of the lattice $L\mathbb{Z}^{d-1}$.

For $u \in \mathbb{Z}^{d-1}$ we will use $\alpha'(u)$ to determine the colors (other than $\pm\infty$) assigned by λ to sites in the vertical column $\{u\} \times \mathbb{Z}$. Colors $1, \dots, d - 1$ will be used unchanged, while color 0 will be treated in a different way.

We now introduce a renormalized percolation process, starting with certain sets to be used in its definition. For a site $x = (x_1, \dots, x_d) \in \mathbb{Z}^d$, write $\underline{x} := (x_1, \dots, x_{d-1}) \in \mathbb{Z}^{d-1}$ and $\bar{x} := x_d$, so that $x = (\underline{x}, \bar{x})$. Let $\underline{C}_{\underline{x}} \subset \mathbb{Z}^{d-1}$ be the 0-cluster of α' centred at $L\underline{x}$. Let $s := \lfloor L/d \rfloor$ (where $\lfloor \cdot \rfloor$ denotes the integer part). Let $\overline{C}_{\bar{x}}$ be the interval $\llbracket s\bar{x}, s(\bar{x} + 1) \rrbracket \subset \mathbb{Z}$. Define the *cell* corresponding to $x \in \mathbb{Z}^d$ to be the set of sites $C_x := \underline{C}_{\underline{x}} \times \overline{C}_{\bar{x}}$. Thus, each cell is a cuboid of height s , and side length r in each horizontal dimension. The centers of the cells are spaced at distance s vertically (so that they abut each other), and at distance L horizontally.

Define a *hole* to be any cube of the form $z + \llbracket 1, r \rrbracket^d$, where $z \in \mathbb{Z}^d$, all of whose sites are closed in the percolation configuration. We say that the cell C_x is *holey* if it contains some hole as a subset. Now we return to the issue of choosing L . Since a hole has volume r^d (a function of J and d), and a cell has height $s = \lfloor L/d \rfloor$, we may choose L a sufficiently large multiple of J (depending on J, d and p) so that the probability that a cell is holey exceeds $1 - (2d)^{-2}$. For later purposes, ensure also that L is large enough that $s > J$ and $\lfloor (L - r)/2 \rfloor > J$. By Theorem 2, there exists a.s. a 1-Lip function $F : \mathbb{Z}^{d-1} \rightarrow \mathbb{Z}_+$, such that all the cells $C_{(u, F(u))}$ for $u \in \mathbb{Z}^{d-1}$ are holey.

We specify next a set of sites surrounding each of the holey cells considered above, to be colored according to α' . For any $\underline{x} \in \mathbb{Z}^{d-1}$, let $\underline{B}_{\underline{x}}$ be the cube $\{v \in \mathbb{Z}^{d-1} : \lfloor v/L \rfloor = \underline{x}\}$ (so that these cubes partition \mathbb{Z}^{d-1}). Let $\overline{B}_{\bar{x}}$ be the interval $\llbracket s\bar{x}, s\bar{x} + L \rrbracket$. Define the *block* corresponding to $x \in \mathbb{Z}^d$ to be the set of sites $B_x := \underline{B}_{\underline{x}} \times \overline{B}_{\bar{x}}$. Thus B_x is a cube of side L which contains the cell C_x (at its bottom-center).

Now we define the coloring λ . For each $u \in \mathbb{Z}^{d-1}$, call the block $B_{(u, F(u))}$ *active*. To each open site $y \in B_{(u, F(u))}$, assign the color $\alpha'(y)$, provided this is one of the colors $1, 2, \dots, d - 1$. For the remaining sites y in the active block [those satisfying $\alpha'(y) = 0$], we proceed as follows. Since the cell is holey, choose one hole $H_u \subset C_{(u, F(u))}$. Since the sites in H_u are closed, they receive no colors. Assign color ∞ to all open sites in the block that lie above the hole H_u , and assign color $-\infty$ to those that lie below H_u . (We say that a site x lies *above* a set S if there exists $y \in S$ with $\underline{x} = \underline{y}$, and for all such y we have $\bar{x} > \bar{y}$; *below* is defined similarly with the inequality reversed.) We have assigned colors to all open sites lying in active blocks. Finally, assign color ∞ to all open sites that lie above some active block, and color $-\infty$ to all those that lie below some active block.

Now we must check that the coloring λ has all the claimed properties. For property (b), note first that if the function F were constant, then each j -cluster for $j = 1, \dots, d - 1$ would have volume at most $L^{d-1} \times L = L^d$, since the coloring

α' has merely been “thickened” vertically to thickness L . The effect of taking a nonconstant F is to displace the active blocks in the vertical direction, and this clearly cannot make these clusters any larger, so we can take $K = L^d$.

Property (c) follows easily from the Lipschitz property of F . The constant d^{-1} arises because for $u, v \in \mathbb{Z}^{d-1}$ with $\|u - v\|_1 = 1$, the centers of the corresponding blocks are at horizontal displacement L from each other, and vertical displacement at most $s \leq L/d$. Once the function g is determined for the centers of the blocks, it can be defined elsewhere by linear interpolation.

To check property (a), suppose on the contrary that there exist two sites x, y with respective colors $+\infty, -\infty$ within ℓ^∞ -distance J of each other. If there is a single active block such that both x and y lie above, below or within it, this contradicts the presence of a hole (which has side length $r > J$) in the corresponding cell. Also, if one of x, y lies within an active block, then the other cannot lie above, below or within a different active block, since $\lfloor (L - r)/2 \rfloor > J$. Therefore the only other case to consider is that x and y lie, respectively, above and below two different active blocks, say $B_{(u, F(u))}$ and $B_{(v, F(v))}$, for some $u, v \in \mathbb{Z}^{d-1}$. In this case we must have $\|u - v\|_\infty = 1$ and therefore $|F(u) - F(v)| \leq \|u - v\|_1 \leq d - 1$, so the height intervals $\overline{B}_{F(u)}$ and $\overline{B}_{F(v)}$ overlap by at least $L - (d - 1)s \geq s > J$, giving again a contradiction. \square

To complete the proof of Theorem 1(c) we will need the following simple geometric fact in order to find an appropriate separating surface. For a vector $x = (x_1, \dots, x_d) \in \mathbb{R}^d$, write \widehat{x}_r for the $(d - 1)$ -vector obtained by dropping the r -coordinate.

LEMMA 11. *Let $a_{\pm 1}, \dots, a_{\pm d}$ be positive constants and define for $i = 1, \dots, d$ the sets*

$$A_i := \{x \in \mathbb{R}^d : x_i \leq d^{-1} \|\widehat{x}_i\|_1 + a_i\};$$

$$A_{-i} := \{x \in \mathbb{R}^d : x_i \geq -d^{-1} \|\widehat{x}_i\|_1 - a_{-i}\}.$$

Then $\bigcap_{i=\pm 1, \dots, \pm d} A_i$ is bounded.

PROOF. We may assume without loss of generality that the a_i are all equal, to a say. For $x \in A_i \cap A_{-i}$ we have $|x_i| \leq d^{-1} \|\widehat{x}_i\|_1 + a$, hence for x in the given intersection, summing the last inequality over i gives

$$\|x\|_1 \leq \frac{d - 1}{d} \|x\|_1 + da,$$

hence $\|x\|_1 \leq d^2 a$. \square

PROOF OF THEOREM 1(c). Fix $d \geq 2$, $M \geq 1$ and $p \in (0, 1)$, and suppose that f is an M -Lip injection from \mathbb{Z}^d to $W_p(\mathbb{Z}^d)$. Let K be the constant from

Proposition 8 for the given values of p, d and with $J := dM$. Let $n := K + 1$. Let N be large enough so that the image $f(\llbracket 1, n \rrbracket^{d-1} \times \{1\})$ is a subset of $\llbracket -N, N \rrbracket^d$.

Now apply Proposition 8 (again with parameters p, d and $J = dM$), but to the translated lattice having its origin at $(N + 1)e_d$, to obtain (a.s.) a coloring of $W_p(\mathbb{Z}^d)$ in which all open sites in $\mathbb{Z}^{d-1} \times \llbracket -\infty, N \rrbracket$ have color $-\infty$. Call this coloring λ_d , and let S_d^+ be the set corresponding to S^+ in Proposition 8(c) (all of whose open sites are colored ∞). Similarly, for each of the two senses of the d coordinate directions, apply Proposition 8 to the lattice rotated and translated so that the part of the half-axis at distance greater than N from the origin is mapped to the positive d -axis. Thus we obtain $2d$ colorings λ_i of $W_p(\mathbb{Z}^d)$, with associated sets S_i^+ , for $i = \pm 1, \dots, \pm d$, such that all the colorings assign color $-\infty$ to $\llbracket -N, N \rrbracket^d$, and λ_i assigns color ∞ to sites sufficiently far along the i coordinate half-axis.

For each i as above, let S_i^{++} be the set of sites y such that every site within ℓ^1 -distance dMn of y lies in S_i^+ . We claim that

$$Z := \mathbb{Z}^d \setminus \bigcup_{i=\pm 1, \dots, \pm d} S_i^{++}$$

is a finite set. This follows from Lemma 11 because $\mathbb{Z}^d \setminus S_i^{++}$ lies in a set of the form A_i in the lemma. [Here it is important that the Lipschitz constant in Proposition 8(c) is d^{-1} .] Since f is injective, it follows that, for some $m > 1$, the site $f((1, \dots, 1, m))$ lies outside Z , and hence lies in S_I^+ for some I . Since $f(\llbracket 1, n \rrbracket^{d-1} \times \{m\})$ has ℓ^1 -diameter at most dMn , this implies that $f(\llbracket 1, n \rrbracket^{d-1} \times \{m\})$ is a subset of S_I^+ , and is therefore colored ∞ in λ_I .

Now define a coloring

$$\chi : \llbracket 1, n \rrbracket^{d-1} \times \llbracket 1, m \rrbracket \rightarrow \{\infty, -\infty, 1, 2, \dots, d - 1\}$$

via $\chi := \lambda_I \circ f$. By the construction, χ satisfies properties (a) and (b) of Proposition 7. Now, if x, y are adjacent sites in G^* then $\|x - y\|_1 \leq d$, and therefore the M -Lip property gives

$$\|f(x) - f(y)\|_\infty \leq \|f(x) - f(y)\|_1 \leq dM = J,$$

so $f(x), f(y)$ are adjacent in $G(W_p(\mathbb{Z}^d), \ell^\infty, J)$. Hence, property (i) in Proposition 8 implies that χ has no two adjacent sites in G^* with colors $+\infty$ and $-\infty$, which is property (c) of Proposition 7. Therefore by Proposition 7, for some $j \neq \pm\infty$, χ has a j -cluster of volume at least n with respect to G^* . Let A be such a cluster. Since f is injective, $f(A)$ also has volume at least n . But by the above observation on adjacency, $f(A)$ is a subset of some j -cluster of λ_I with respect to $G(W_p(\mathbb{Z}^d), \ell^\infty, J)$. This contradicts property (b) in Proposition 8 because $n > K$. □

4. Embedding and quasi-isometry. We will use the following simple renormalization construction. Fix an integer $r \geq 1$. For a site $x = (x_1, \dots, x_D) \in \mathbb{Z}^D$ define the corresponding *clump* (or *r-clump*) to be the set of r sites given by

$$K_x := \{(x_1, \dots, x_{D-1}, rx_D + i) : i \in \llbracket 0, r - 1 \rrbracket\}.$$

The clumps $(K_x : x \in \mathbb{Z}^D)$ form a partition of \mathbb{Z}^D , with the geometry of \mathbb{Z}^D stretched by a factor r in the D th coordinate. If $\|x - y\| = k$, then, for all $u \in K_x$ and $v \in K_y$, we have $\|u - v\| \leq (2r - 1)k$.

PROOF OF PROPOSITION 3. Let $d < D$, which is to say that $M_c(d, D) < \infty$. Any given r -clump contains one or more open sites with probability $1 - (1 - p)^r$. If this probability exceeds $p_c(d, D, M)$, there exists a.s. an M -Lip injection $f : \mathbb{Z}^d \rightarrow \mathbb{Z}^D$ such that, for each $y \in f(\mathbb{Z}^d)$, the clump K_y contains some open site. By choosing one representative open site in each such clump, we obtain a $(2r - 1)M$ -Lip injection from \mathbb{Z}^d to $W_p(\mathbb{Z}^D)$. Hence,

$$p_c(d, D, (2r - 1)M) \leq 1 - (1 - p_c(d, D, M))^{1/r}.$$

The claim follows by the monotonicity of p_c in M . \square

PROOF OF PROPOSITION 4. (a) We may assume with loss of generality that $d = 2 \leq D$. The proof follows that of Theorem 1(b) as presented in Section 2, without one difference. Let $\eta \in \Omega_2$. The event $A = A(x_1, \dots, x_k)$ of Lemma 6 is redefined as the event that there exists a singly-infinite path $0 = y_0, y_1, \dots$ in \mathbb{Z}^D such that the sites $(x_i + y_j : i = 1, \dots, k, j = 0, 1, \dots)$ are distinct, and $\omega(x_i + y_j) = \eta(i, j)$ for all such i, j . As in the proof of Lemma 6, $\mathbb{P}_p(A) = 0$ whenever $\max\{p, 1 - p\} < (2D)^{-1/k}$. The proof is now completed as for the earlier theorem.

(b) Let $d < D$ and write $m = M_c(d, D) < \infty$. Given $p \in (0, 1)$, choose r sufficiently large that any given r -clump contains both an open and a closed site with probability exceeding $p_c(d, D, m)$. There exists a.s. an m -Lip injection $f : \mathbb{Z}^d \rightarrow \mathbb{Z}^D$ such that, for each $y \in f(\mathbb{Z}^d)$, the r -clump K_y contains both an open and a closed site. Hence, for any configuration η , by choosing the open or the closed site as appropriate in each r -clump, we obtain a $(2r - 1)m$ -Lip embedding of η into ω .

(c) Let $d \geq 1$, $M \geq 1$, and let $\eta \in \Omega_d$ be partially periodic. Without loss of generality, we may assume, for some $r \in \mathbb{Z}_+$ and all $y \in \mathbb{Z}^d$, that $\eta(ry) = \eta(0) = 1$. Let $\omega \in \Omega_d$ and assume there exists an M -Lip embedding f from η into ω . Let $g : \mathbb{Z}^d \rightarrow \mathbb{Z}^d$ be given by $g(x) = f(rx)$. Then g is an rM -Lip injection from \mathbb{Z}^d into $W_p(\mathbb{Z}^d)$. By Theorem 1(c), such an injection exists only for ω lying in some \mathbb{P}_p -null set. \square

PROOF OF PROPOSITION 5. (a) We assume without loss of generality that $D = d + 1$. Given $p < 1$, take r sufficiently large that a given r -clump in \mathbb{Z}^D contains some open site with probability exceeding $1 - (2D)^{-2}$. By Theorem 2,

there exists a 1-Lip map $F: \mathbb{Z}^{D-1} \rightarrow \mathbb{Z}_+$ such that for all $u \in \mathbb{Z}^{D-1}$, the r -clump $K_{(u, F(u))}$ contains some open site. By choosing an arbitrary open site to represent each such clump, we obtain a quasi-isometry of the required form.

(b) Let $d \geq D$, and suppose that with positive probability there exists a quasi-isometry from (\mathbb{Z}^d, ℓ^1) to some subspace of $(W_p(\mathbb{Z}^D), \ell^1)$. We will prove that, for some $p' \in (0, 1)$ and $M \geq 1$, there exists an M -Lip injection $g: \mathbb{Z}^d \rightarrow W_{p'}(\mathbb{Z}^D)$, which will contradict (2).

Recall the parameters $\mathbf{c} = (c_1, c_2, \dots, c_5)$ in the definition of a \mathbf{c} -quasi-isometry, and let $Q_{\mathbf{c}}$ be the event that there exists a \mathbf{c} -quasi-isometry from (\mathbb{Z}^d, ℓ^1) to some subset of $(W_p(\mathbb{Z}^D), \ell^1)$. Since each $Q_{\mathbf{c}}$ is invariant under the action of translations of \mathbb{Z}^D , it has probability 0 or 1. Under the above assumption, the event $\bigcup_{\mathbf{c}} Q_{\mathbf{c}}$ has positive probability. By the obvious monotonicities in the parameters c_i , this union is equal to the union $\bigcup_{\mathbf{c} \in (\mathbb{Q} \cap (0, \infty))^5} Q_{\mathbf{c}}$ over rational parameters, and hence there exists a *deterministic* \mathbf{c} such that $Q_{\mathbf{c}}$ has probability 1. We choose \mathbf{c} accordingly, and let $\mathcal{F}_{\mathbf{c}}$ be the (random) set of quasi-isometries of the required type.

A quasi-isometry $f \in \mathcal{F}_{\mathbf{c}}$ is not necessarily an injection, but, by the properties of a \mathbf{c} -quasi-isometry, there exists $C = C(d, D, \mathbf{c})$ such that, for all $y \in W_p(\mathbb{Z}^D)$ we have $|f^{-1}(y)| \leq C$. Let $r = C$, and take $p' \in (0, 1)$ sufficiently large that, with probability at least p , every site in any given r -clump is p' -open. Let $f \in \mathcal{F}_{\mathbf{c}}$ be such that: for $y \in f(\mathbb{Z}^d)$, every site in K_y is p' -open. Since the pre-image under f of any $y \in \mathbb{Z}^D$ has cardinality C or less, we may construct an injection $g: \mathbb{Z}^d \rightarrow W_{p'}(\mathbb{Z}^D)$ such that, for $y \in \mathbb{Z}^D$, every $x \in f^{-1}(y)$ has $g(x) \in K_y$, and furthermore distinct elements $x \in f^{-1}(y)$ have distinct images $g(x)$. It is easily seen that g is an M -Lip injection for some $M = M(d, D, \mathbf{c})$. \square

5. Open questions.

5.1. Derive quantitative versions of Theorem 1. For example, fix d, M, p , and let $N = N(n)$ be the smallest integer such that there exists an M -Lipschitz injection from the cube $[1, n]^d \cap \mathbb{Z}^d$ to the open sites of $[1, N]^d \cap \mathbb{Z}^d$ with probability at least $\frac{1}{2}$. How does N behave as $n \rightarrow \infty$?

5.2. For which graphs G and which M is it the case that for p sufficiently close to 1 there exists an M -Lipschitz injection from $V(G)$ to the open sites of $V(G)$ (where M -Lipschitz refers to the graph metric of G)? Theorem 1(c) shows that for \mathbb{Z}^d , no $M < \infty$ suffices. On the other hand, for the 3-regular tree, $M = 2$ suffices, by the well-known fact that percolation on a 4-ary tree contains a binary tree for p sufficiently close to 1.

5.3. We may interpolate between 1-Lipschitz and 2-Lipschitz maps as follows. Let S be a subset of $\{-1, 0, 1\}^D$, and let G be the graph with vertex set \mathbb{Z}^D and an edge between u, v whenever $u - v$ or $v - u$ belongs to S . For which d, D and S does there exist an injection from \mathbb{Z}^d to the open sites of \mathbb{Z}^D that maps neighbors in \mathbb{Z}^d to neighbors in G ?

5.4. Does there exist a configuration $\eta \in \{0, 1\}^d$ such that, with positive probability there exists a Lipschitz embedding of η into the percolation configuration ω on \mathbb{Z}^d ? When $d = 1$, this is related to the main problem of [7].

Acknowledgments. We thank Olle Häggström and Peter Winkler for valuable discussions. G. R. Grimmett acknowledges support from Microsoft Research during his stay as a Visiting Researcher in the Theory Group in Redmond. This work was completed during his attendance at a programme at the Isaac Newton Institute, Cambridge.

REFERENCES

- [1] BAKER, J. K. (1970). A combinatorial proof of Tucker's lemma for the n -cube. *J. Combin. Theory* **8** 279–290. [MR0253909](#)
- [2] DIRR, N., DONDL, P. W., GRIMMETT, G. R., HOLROYD, A. E. and SCHEUTZOW, M. (2010). Lipschitz percolation. *Electron. Comm. Probab.* **15** 14–21. [MR2581044](#)
- [3] DIRR, N., DONDL, P. W. and SCHEUTZOW, M. (2009). Pinning of interfaces in random media. Unpublished manuscript. Available at [arXiv:0911.4254](#).
- [4] GRIMMETT, G. R. (2010). Three problems for the clairvoyant demon. In *Probability and Mathematical Genetics* (N. H. Bingham and C. M. Goldie, eds.) 379–395. Cambridge Univ. Press, Cambridge.
- [5] GRIMMETT, G. R. and HOLROYD, A. E. (2010). Geometry of Lipschitz percolation. *Ann. Inst. Henri Poincaré Probab. Stat.* To appear. Available at [arXiv:1007.3762](#).
- [6] GRIMMETT, G. R. and HOLROYD, A. E. (2010). Plaquettes, spheres, and entanglement. *Electron. J. Probab.* **15** 1415–1428.
- [7] GRIMMETT, G. R., LIGGETT, T. M. and RICHTHAMMER, T. (2010). Percolation of arbitrary words in one dimension. *Random Structures Algorithms* **37** 85–99. [MR2674622](#)
- [8] LEFSCHETZ, S. (1949). *Introduction to Topology*. Princeton Mathematical Series **11**. Princeton Univ. Press, Princeton, NJ. [MR0031708](#)
- [9] LYONS, R. (2000). Phase transitions on nonamenable graphs. *J. Math. Phys.* **41** 1099–1126. [MR1757952](#)
- [10] MATOUŠEK, J. (2003). *Using the Borsuk–Ulam Theorem*. Springer, Berlin. [MR1988723](#)
- [11] PELED, R. (2010). On rough isometries of Poisson processes on the line. *Ann. Appl. Probab.* **20** 462–494. [MR2650039](#)
- [12] TUCKER, A. W. (1946). Some topological properties of disk and sphere. In *Proc. First Canadian Math. Congress, Montreal, 1945* 285–309. Univ. of Toronto Press, Toronto. [MR0020254](#)

STATISTICAL LABORATORY
CENTRE FOR MATHEMATICAL SCIENCES
CAMBRIDGE UNIVERSITY
WILBERFORCE ROAD
CAMBRIDGE CB3 0WB
UNITED KINGDOM
E-MAIL: g.r.grimmett@statslab.cam.ac.uk
URL: <http://www.statslab.cam.ac.uk/~grg/>

MICROSOFT RESEARCH
1 MICROSOFT WAY
REDMOND, WASHINGTON 98052
USA
AND
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF BRITISH COLUMBIA
VANCOUVER, BC V6T 1Z2
CANADA
E-MAIL: holroyd@microsoft.com
URL: <http://research.microsoft.com/~holroyd>