

## INFLUENCE AND SHARP-THRESHOLD THEOREMS FOR MONOTONIC MEASURES

BY B. T. GRAHAM AND G. R. GRIMMETT

*Cambridge University*

The influence theorem for product measures on the discrete space  $\{0, 1\}^N$  may be extended to probability measures with the property of monotonicity (which is equivalent to ‘strong positive-association’). Corresponding results are valid for probability measures on the cube  $[0, 1]^N$  that are absolutely continuous with respect to Lebesgue measure. These results lead to a sharp-threshold theorem for measures of random-cluster type, and this may be applied to box-crossings in the two-dimensional random-cluster model.

**1. Introduction.** Influence and sharp-threshold theorems have proved useful in the study of problems in discrete probability. Reliability theory and random graphs provided early problems of this type, followed by percolation. Important progress has been made since [2, 16] towards a general theory. The reader is referred to [10, 11] for a history and bibliography.

Let  $\Omega = \{0, 1\}^N$  where  $N < \infty$ , and let  $\mu_p$  be the product measure on  $\Omega$  with density  $p$ . Vectors in  $\Omega$  are denoted by  $\omega = (\omega(i) : 1 \leq i \leq N)$ . For any increasing subset  $A$  of  $\Omega$ , and any  $i \in \{1, 2, \dots, N\}$ , we define the *conditional influence*  $I_A(i)$  by

$$(1.1) \quad I_A(i) = \mu_p(A \mid X_i = 1) - \mu_p(A \mid X_i = 0),$$

where  $X_i : \Omega \rightarrow \mathbb{R}$  is given by  $X_i(\omega) = \omega(i)$ . It is well known (see [6, 11, 16, 22]) that there exists an absolute positive constant  $c$  such that the following holds. For all  $N$ , all  $p \in (0, 1)$ , and all increasing  $A$ , there exists  $i \in \{1, 2, \dots, N\}$  such that

$$(1.2) \quad I_A(i) \geq c \min\{\mu_p(A), 1 - \mu_p(A)\} \frac{\log N}{N}.$$

The proof uses discrete Fourier analysis and a technique known as ‘hypercontractivity’. Inequality (1.2) is usually stated for the case  $p = \frac{1}{2}$ , but it holds with the same constant  $c$  for all  $p \in (0, 1)$ .

---

*AMS 2000 subject classifications:* Primary 60E15, 60K35; secondary 82B31, 82B43.

*Keywords and phrases:* influence, sharp threshold, monotonic measure, FKG lattice condition, positive association, random-cluster model, percolation.

There is an important application to the theory of sharp thresholds for product measures, see [11]. Let  $\Pi_N$  be the set of all permutations of the index set  $I = \{1, 2, \dots, N\}$ . A subgroup  $\mathcal{A}$  of  $\Pi_N$  is said to *act transitively* on  $I$  if, for all pairs  $j, k \in I$ , there exists  $\pi \in \mathcal{A}$  with  $\pi_j = k$ . Any  $\pi \in \Pi_N$  acts on  $\Omega$  by  $\pi\omega = (\omega(\pi_i) : 1 \leq i \leq N)$ . An event  $A$  is called *symmetric* if there exists a subgroup  $\mathcal{A}$  of  $\Pi_N$  acting transitively on  $I$  such that  $A = \pi A$  for all  $\pi \in \mathcal{A}$ . If  $A$  is symmetric, then  $I_A(j) = I_A(k)$  for all  $j, k$ . By summing (1.2) over  $i$  we obtain for symmetric  $A$  that

$$(1.3) \quad \sum_{i=1}^N I_A(i) \geq c \min\{\mu_p(A), 1 - \mu_p(A)\} \log N.$$

It is standard (see the discussion of Russo's formula in [12]) that

$$(1.4) \quad \frac{d}{dp} \mu_p(A) = \sum_{i=1}^N I_A(i),$$

and it follows as in [11] that, for  $0 < \epsilon < \frac{1}{2}$ , the function  $f(p) = \mu_p(A)$  increases from  $\epsilon$  to  $1 - \epsilon$  over an interval of values of  $p$  with length smaller in order than  $1/\log N$ .

We refer to such a statement as a 'sharp-threshold theorem', and we note that such results have wide applications to problems of discrete probability. The example to be explored later in this paper is the random-cluster model on the square lattice  $\mathbb{L}^2$ . In the special case of percolation on  $\mathbb{L}^2$ , a result of the above type (with a weaker bound) was used in [20] to (re-)prove the principal duality theorem for site percolation on  $\mathbb{L}^2$ . More recently, (1.3)–(1.4) have been used in [5] to obtain a further proof that the critical probability  $p_c$  of bond percolation on the square lattice satisfies  $p_c = \frac{1}{2}$ . Using a similar argument in a second paper [4], it is proved that the critical probability of site percolation on a certain Poisson–Voronoi (random) graph in  $\mathbb{R}^2$  equals  $\frac{1}{2}$  almost surely.

The principal purpose of the current article is to extend the results above to probability measures more general than product measures. We shall prove such results for measures having a certain condition of 'monotonicity', which is equivalent to the FKG lattice condition and is described in the next section. There are many situations in the probabilistic theory of statistical mechanics where such measures are encountered, including the Ising model and the random-cluster model.

We define monotonic probability measures in Section 2, and we note there that monotonicity is equivalent to the FKG lattice condition. This is followed by an influence theorem for monotonic measures.

A monotonic measure  $\mu$  may be used as the basis of a certain parametric family of measures on  $\Omega$  indexed by a parameter  $p \in (0, 1)$ . The influence theorem for  $\mu$  may then be used to obtain a sharp-threshold theorem for this class, as described in Section 3.

The influence theorem on the discrete space  $\{0, 1\}^N$  was extended in [6] to product measures on the Euclidean cube  $[0, 1]^N$ . Using the methods of Section 2, similar results may be proved for general monotonic measures on  $[0, 1]^N$ . Unlike the discrete case, such an influence theorem does not appear to imply a corresponding sharp-threshold theorem. This is discussed in Section 4.

We turn finally to the random-cluster model, which may be viewed as an extension of percolation and a generalization of the Ising/Potts models for ferromagnetism, see [13, 14]. The random-cluster measure is defined in Section 5, and the sharp-threshold theorem is applied to the existence of box-crossings in two dimensions.

**2. Influence for monotonic measures.** We begin this section with a classification, further details of which may be found in [14]. Let  $1 \leq N < \infty$ , and write  $I = \{1, 2, \dots, N\}$  and  $\Omega = \{0, 1\}^N$ . The set of all subsets of  $\Omega$  is denoted by  $\mathcal{F}$ . A probability measure  $\mu$  on  $(\Omega, \mathcal{F})$  is said to be *positive* if  $\mu(\omega) > 0$  for all  $\omega \in \Omega$ . It is said to satisfy the *FKG lattice condition* if

$$(2.1) \quad \mu(\omega_1 \vee \omega_2)\mu(\omega_1 \wedge \omega_2) \geq \mu(\omega_1)\mu(\omega_2) \quad \text{for all } \omega_1, \omega_2 \in \Omega,$$

where  $\omega_1 \vee \omega_2$  and  $\omega_1 \wedge \omega_2$  are given by

$$\begin{aligned} \omega_1 \vee \omega_2(i) &= \max\{\omega_1(i), \omega_2(i)\}, & i \in I, \\ \omega_1 \wedge \omega_2(i) &= \min\{\omega_1(i), \omega_2(i)\}, & i \in I. \end{aligned}$$

See [9, 14].

The set  $\Omega$  is a partially ordered set with the partial order:  $\omega \geq \omega'$  if  $\omega(i) \geq \omega'(i)$  for all  $i \in I$ . A non-empty event  $A \in \mathcal{F}$  is called *increasing* if:  $\omega \in A$  whenever there exists  $\omega'$  with  $\omega \geq \omega'$  and  $\omega' \in A$ . It is called *decreasing* if its complement is increasing. For probability measures  $\mu_1, \mu_2$  on  $(\Omega, \mathcal{F})$ , we write  $\mu_1 \leq_{\text{st}} \mu_2$ , and say that  $\mu_1$  is dominated stochastically by  $\mu_2$ , if

$$\mu_1(A) \leq \mu_2(A) \quad \text{for all increasing events } A.$$

The indicator function of an event  $A$  is denoted by  $1_A$ . For  $i \in I$ , we define the random variable  $X_i$  by  $X_i(\omega) = \omega(i)$ .

A probability measure  $\mu$  on  $\Omega$  is said to be *positively associated* if

$$\mu(A \cap B) \geq \mu(A)\mu(B) \quad \text{for all increasing events } A, B.$$

The famous FKG inequality of [9] asserts that a positive probability measure  $\mu$  is positively associated if it satisfies the FKG lattice condition. It is well known that the FKG lattice condition is not necessary for positive association, and we explore this next.

We shall for simplicity restrict ourselves henceforth to positive measures. The FKG lattice condition is equivalent to a stronger property termed ‘strong positive-association’. For  $J \subseteq I$  and  $\xi \in \Omega$ , let  $\Omega_J = \{0, 1\}^J$  and

$$(2.2) \quad \Omega_J^\xi = \{\omega \in \Omega : \omega(i) = \xi(i) \text{ for } i \in I \setminus J\}.$$

The set of all subsets of  $\Omega_J$  is denoted by  $\mathcal{F}_J$ . Let  $\mu$  be a positive probability measure on  $(\Omega, \mathcal{F})$ , and define the conditional probability measure  $\mu_J^\xi$  on  $(\Omega_J, \mathcal{F}_J)$  by

$$(2.3) \quad \mu_J^\xi(\omega_J) = \mu(X_j = \omega_J(j) \text{ for } j \in J \mid X_i = \xi(i) \text{ for } i \in I \setminus J), \quad \omega_J \in \Omega_J.$$

We say that  $\mu$  is *strongly positively-associated* if: for all  $J \subseteq I$  and all  $\xi \in \Omega$ , the measure  $\mu_J^\xi$  is positively associated.

We call  $\mu$  *monotonic* if: for all  $J \subseteq I$ , all increasing subsets  $A$  of  $\Omega_J$ , and all  $\xi, \zeta \in \Omega$ ,

$$(2.4) \quad \mu_J^\xi(A) \leq \mu_J^\zeta(A) \quad \text{whenever } \xi \leq \zeta.$$

That is,  $\mu$  is monotonic if, for all  $J \subseteq I$ ,

$$(2.5) \quad \mu_J^\xi \leq_{st} \mu_J^\zeta \quad \text{whenever } \xi \leq \zeta.$$

We call  $\mu$  *1-monotonic* if (2.5) holds for all singleton sets  $J$ , which is to say that, for all  $j \in I$ ,

$$(2.6) \quad \mu(X_j = 1 \mid X_i = \xi(i) \text{ for all } i \in I \setminus \{j\})$$

is non-decreasing in  $\xi$ .

The following theorem is fairly standard, and the proof may be found in [14].

**THEOREM 2.7.** *Let  $\mu$  be a positive probability measure on  $(\Omega, \mathcal{F})$ . The following are equivalent.*

- (i)  $\mu$  is strongly positively-associated.
- (ii)  $\mu$  satisfies the FKG lattice condition.
- (iii)  $\mu$  is monotonic.

(iv)  $\mu$  is 1-monotonic.

Our principal influence theorem is as follows. For a positive probability measure  $\mu$  and an increasing event  $A$ , the *conditional influence* of the index  $i \in I$  is given as in (1.1) by

$$(2.8) \quad I_A(i) = \mu(A \mid X_i = 1) - \mu(A \mid X_i = 0).$$

For a product measure  $\mu_p$ , the influence of the index  $i$  was defined in [2, 16] as  $\mu_p(\omega^i \in A, \omega_i \notin A)$ , where  $\omega^i$  (respectively,  $\omega_i$ ) denotes the configuration obtained from  $\omega$  by setting  $\omega(i)$  equal to 1 (respectively, 0). We refer to the latter quantity as the *absolute influence* of index  $i$ . The absolute and conditional influences are equal for product measures, but one should note that

$$(2.9) \quad I_A(i) \neq \mu(\omega^i \in A, \omega_i \notin A)$$

for general probability measures  $\mu$ . Further discussion of this point is provided after the next theorem. See also [15].

**THEOREM 2.10 (Influence).** *There exists a constant  $c \in (0, \infty)$  such that the following holds. Let  $N \geq 1$  and let  $A$  be an increasing subset of  $\Omega = \{0, 1\}^N$ . Let  $\mu$  be a positive probability measure on  $(\Omega, \mathcal{F})$  that is monotonic. There exists  $i \in I$  such that*

$$(2.11) \quad I_A(i) \geq c \min\{\mu(A), 1 - \mu(A)\} \frac{\log N}{N}.$$

Since product measures are monotonic, this extends the influence theorem of [16]. In the proof of Theorem 2.10, we shall encode the measure  $\mu$  in terms of Lebesgue measure on  $[0, 1]^N$ , and we shall appeal to the influence theorem of [6]. Thus, we shall require no further arguments of discrete Fourier analysis than those already present in [6, 16].

We return briefly to the discussion of absolute and conditional influences. Suppose, for illustration, that  $P$  is chosen at random with  $\mathbb{P}(P = \frac{1}{3}) = \mathbb{P}(P = \frac{2}{3}) = \frac{1}{2}$  and that, conditional on the value of  $P$ , we are provided with independent Bernoulli random variables  $X_1, X_2, \dots, X_N$  with parameter  $P$ . It is easily checked that the law of the vector  $X_1, X_2, \dots, X_N$  satisfies the FKG lattice condition. Consider the increasing event  $A = \{S_N > \frac{1}{2}N\}$ , where  $S_N = X_1 + X_2 + \dots + X_N$ . By symmetry, the conditional influence of each index is the same, as is the absolute influence of each index. It is an easy calculation that

$$I_A(1) = \frac{1}{3} + o(1) \quad \text{as } N \rightarrow \infty.$$

On the other hand,

$$\begin{aligned} \mathbb{P}(\omega^1 \in A, \omega_1 \notin A) &= \mathbb{P}\left(\frac{1}{2}N - 1 < \sum_{i=2}^N X_i \leq \frac{1}{2}N\right) \\ &= o(e^{-\gamma N}) \quad \text{as } N \rightarrow \infty, \end{aligned}$$

for some  $\gamma > 0$ . This example indicates not only that the absolute and conditional influences can be very different, but also that the conclusion of Theorem 2.10 would be false if re-stated for absolute influences.

Definition (2.8) is well suited to measures  $\mu$  that are monotonic. When  $\mu$  is non-monotonic it can happen that  $I_A(i) = 0$  for all  $i$ . For example, consider a circular table with  $n$  places, and let  $\mu$  be the law induced on  $\{0, 1\}^n$  by picking two distinct places uniformly at random. Let  $A$  be the (increasing) event that at least two chosen places are adjacent. It is easily seen that  $\mu(A) = 2/(n-1)$ , and that  $I_A(i) = 0$  for every  $i$ . The measure  $\mu$  is not positive, but a small perturbation results in a positive measure with influences as small as required.

In the proof of Theorem 2.10 following, we see that monotonicity has the effect of increasing the influence of each coordinate in  $I$ .

PROOF OF THEOREM 2.10. Let  $A \in \mathcal{F}$  be an increasing event, and let  $\mu$  be positive and monotonic. Let  $\lambda$  denote Lebesgue measure on the cube  $[0, 1]^N$ . We propose to construct an increasing subset  $B$  of  $[0, 1]^N$  with the property that  $\lambda(B) = \mu(A)$ , to apply the influence theorem of [6] to the set  $B$ , and to deduce the claim. This will be done via a certain function  $f : [0, 1]^N \rightarrow \{0, 1\}^N$  that we construct next.

Let  $\mathbf{x} = (x_i : 1 \leq i \leq N) \in [0, 1]^N$ , and let  $f(\mathbf{x}) = (f_i(\mathbf{x}) : 1 \leq i \leq N)$  be given recursively as follows. The first coordinate  $f_1(\mathbf{x})$  is defined by:

$$(2.12) \quad \text{with } a_1 = \mu(X_1 = 1), \quad \text{set } f_1(\mathbf{x}) = \begin{cases} 1 & \text{if } x_1 > 1 - a_1, \\ 0 & \text{otherwise,} \end{cases}$$

and we note that  $f(\mathbf{x})$  depends on  $x_1$  only. Suppose we know  $f_i(\mathbf{x})$  for  $1 \leq i < k$ . Let

$$(2.13) \quad a_k = a_k(x_1, x_2, \dots, x_{k-1}) = \mu(X_k = 1 \mid X_i = f_i(\mathbf{x}) \text{ for } 1 \leq i < k),$$

and define

$$(2.14) \quad f_k(\mathbf{x}) = \begin{cases} 1 & \text{if } x_k > 1 - a_k, \\ 0 & \text{otherwise.} \end{cases}$$

Let  $\mathbf{x} \leq \mathbf{x}'$ , and write  $a_k = a_k(\mathbf{x})$  and  $a'_k = a_k(\mathbf{x}')$  for the corresponding values in (2.12)–(2.13). Clearly  $a_1 = a'_1$ , so that  $f_1(\mathbf{x}) \leq f_1(\mathbf{x}')$ . Since  $\mu$  is monotonic,  $a_2 \leq a'_2$ , so that  $f_2(\mathbf{x}) \leq f_2(\mathbf{x}')$ . Continuing inductively, we find that  $f_k(\mathbf{x}) \leq f_k(\mathbf{x}')$  for all  $k$ , which is to say that  $f(\mathbf{x}) \leq f(\mathbf{x}')$ . Therefore,  $f$  is non-decreasing on  $[0, 1]^N$ . Let  $B$  be the increasing subset of  $[0, 1]^N$  given by  $B = f^{-1}(A)$ .

We make four notes concerning the definition of  $f$ .

- (1) Each  $a_k$  depends only on  $x_1, x_2, \dots, x_{k-1}$ .
- (2) Since  $\mu$  is positive, the  $a_k$  satisfy  $0 < a_k < 1$  for all  $\mathbf{x} \in [0, 1]^N$  and  $k \in I$ .
- (3) For  $\mathbf{x} \in [0, 1]^N$  and  $k \in I$ , the values  $f_k(\mathbf{x}), f_{k+1}(\mathbf{x}), \dots, f_N(\mathbf{x})$  depend on  $x_1, x_2, \dots, x_{k-1}$  only through the values  $f_1(\mathbf{x}), f_2(\mathbf{x}), \dots, f_{k-1}(\mathbf{x})$ .
- (4) The function  $f$  and the event  $B$  depend on the ordering of the set  $I$ .

Let  $U = (U_i : 1 \leq i \leq N)$  be the identity function on  $[0, 1]^N$ , and note that  $U$  has law  $\lambda$ . By the method of construction of the function  $f$ ,  $f(U)$  has law  $\mu$ . In particular,

$$(2.15) \quad \mu(A) = \lambda(f(U) \in A) = \lambda(U \in f^{-1}(A)) = \lambda(B).$$

Let

$$J_B(i) = \lambda(B \mid U_i = 1) - \lambda(B \mid U_i = 0),$$

where the conditional probabilities are to be interpreted as

$$\lambda(B \mid U_i = u) = \lim_{\epsilon \downarrow 0} \lambda(B \mid U_i \in (u - \epsilon, u + \epsilon)).$$

Since  $B$  is an event with a certain simple structure, this is the same as  $\lambda_{N-1}(B_i^u)$  for  $u = 0, 1$ , where  $\lambda_{N-1}$  is  $(N-1)$ -dimensional Lebesgue measure and  $B_i^u$  is the set of all  $(N-1)$ -vectors  $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_N)$  such that  $(x_1, \dots, x_{i-1}, u, x_{i+1}, \dots, x_N) \in B$ .

By Theorem 1 of [6], we may find a constant  $c > 0$ , independent of the choice of  $N$  and  $A$ , such that: there exists  $i \in I$  with

$$(2.16) \quad J_B(i) \geq c \min\{\lambda(B), 1 - \lambda(B)\} \frac{\log N}{N}.$$

We choose  $i$  accordingly.

We claim that

$$(2.17) \quad I_A(j) \geq J_B(j) \quad \text{for } j \in I.$$

Once (2.17) is shown, the claim follows from (2.15) and (2.16). We prove next that

$$(2.18) \quad I_A(1) = J_B(1).$$

We have that

$$\begin{aligned}
(2.19) \quad I_A(1) &= \mu(A \mid X_1 = 1) - \mu(A \mid X_1 = 0) \\
&= \lambda(B \mid f_1(U) = 1) - \lambda(B \mid f_1(U) = 0) \\
&= \lambda(B \mid U_1 > 1 - a_1) - \lambda(B \mid U_1 \leq 1 - a_1) \\
&= \lambda(B \mid U_1 = 1) - \lambda(B \mid U_1 = 0) \\
&= J_B(1),
\end{aligned}$$

where we have used notes (2) and (3) above.

We turn our attention to (2.17) with  $j \geq 2$ , and we re-order the set  $I$  to bring the index  $j$  to the front. That is, we let  $K$  be the re-ordered index set  $K = (k_1, k_2, \dots, k_N) = (j, 1, 2, \dots, j-1, j+1, \dots, N)$ . We write  $g = (g_{k_i} : 1 \leq i \leq N)$  for the associated function given by (2.12)–(2.14) subject to the new ordering, and  $C = g^{-1}(A)$ . Thinking of (2.12)–(2.14) as an algorithm for constructing  $f$ , we are applying the same algorithm to the re-ordered set  $K$ .

We claim that

$$(2.20) \quad J_C(k_1) \geq J_B(j).$$

By (2.19) with  $I$  replaced by  $K$ ,  $J_C(k_1) = I_A(j)$ , and (2.17) follows. It remains to prove (2.20), and we shall use monotonicity again for this.

It suffices for (2.20) to prove that

$$(2.21) \quad \lambda(C \mid U_j = 1) \geq \lambda(B \mid U_j = 1),$$

together with the reversed inequality given  $U_j = 0$ . The conditioning of the left-hand side of (2.21) refers to the first coordinate encountered by the algorithm (2.12)–(2.14) when applied to the re-ordered set  $K$ . Let

$$(2.22) \quad \bar{U} = (U_1, U_2, \dots, U_{j-1}, 1, U_{j+1}, \dots, U_N).$$

The 0/1-vector  $f(\bar{U}) = (f_i(\bar{U}) : 1 \leq i \leq N)$  is constructed sequentially (as above) by considering the indices  $1, 2, \dots, N$  in turn. At stage  $k$ , we declare  $f_k(\bar{U})$  to equal 1 if  $U_k$  exceeds a certain function  $a_k$  of the variables  $f_i(\bar{U})$ ,  $1 \leq i < k$ . By the monotonicity of  $\mu$ , this function is non-increasing in these variables. The index  $j$  plays a special role in that: (i)  $f_j(\bar{U}) = 1$ , and (ii) given this fact, it is more likely than before that the variables  $f_k(\bar{U})$ ,  $j < k \leq N$ , will take the value 1. The values  $f_k(\bar{U})$ ,  $1 \leq k < j$  are unaffected by the value of  $U_j$ .

Consider now the 0/1-vector  $g(\bar{U}) = (g_{k_r}(\bar{U}) : 1 \leq r \leq N)$ , constructed in the same manner as above but with the new ordering  $K$  of the index set  $I$ .

First we examine index  $k_1 (= j)$ , and we automatically declare  $g_{k_1}(\bar{U}) = 1$  (since  $U_j = 1$ ). We then construct  $g_{k_r}(\bar{U})$ ,  $2 \leq r \leq N$ , in sequence. Since the  $a_k$  are non-decreasing in the variables constructed so far,

$$(2.23) \quad g_{k_r}(\bar{U}) \geq f_{k_r}(\bar{U}), \quad r = 2, 3, \dots, N.$$

Therefore,  $g(\bar{U}) \geq f(\bar{U})$ , implying as required that

$$(2.24) \quad \lambda(C \mid U_j = 1) = \lambda(g(\bar{U}) \in A) \geq \lambda(f(\bar{U}) \in A) = \lambda(B \mid U_j = 1).$$

Inequality (2.21) follows. The same argument implies the reversed inequality obtained from (2.21) by reversing the conditioning to  $U_j = 0$ . This implies (2.20).

A formal proof of (2.23) follows. Suppose that  $r$  is such that  $g_{k_s}(\bar{U}) \geq f_{k_s}(\bar{U})$  for  $2 \leq s < r$ . By (2.14), for  $r \leq j$ ,

$$\begin{aligned} f_{k_r}(\bar{U}) &= 1 && \text{if } U_{k_r} > \mu(X_{k_r} = 0 \mid X_{k_s} = f_{k_s}(\bar{U}) \text{ for } 2 \leq s < r), \\ g_{k_r}(\bar{U}) &= 1 && \text{if } U_{k_r} > \mu(X_{k_r} = 0 \mid X_{k_s} = g_{k_s}(\bar{U}) \text{ for } 1 \leq s < r). \end{aligned}$$

Now  $g_{k_1}(\bar{U}) = 1$  and, by the induction hypothesis and monotonicity,

$$\begin{aligned} \mu(X_{k_r} = 0 \mid X_{k_s} = f_{k_s}(\bar{U}) \text{ for } 2 \leq s < r) \\ \geq \mu(X_{k_r} = 0 \mid X_{k_s} = g_{k_s}(\bar{U}) \text{ for } 1 \leq s < r), \end{aligned}$$

whence  $g_{k_r}(\bar{U}) \geq f_{k_r}(\bar{U})$  as required.

Consider finally the case  $j < r \leq N$ . Then

$$\begin{aligned} f_{k_r}(\bar{U}) &= 1 && \text{if } U_{k_r} > \mu(X_{k_r} = 0 \mid X_{k_s} = f_{k_s}(\bar{U}) \text{ for } 1 \leq s < r), \\ g_{k_r}(\bar{U}) &= 1 && \text{if } U_{k_r} > \mu(X_{k_r} = 0 \mid X_{k_s} = g_{k_s}(\bar{U}) \text{ for } 1 \leq s < r), \end{aligned}$$

and the conclusion follows as before.  $\square$

**3. Sharp-threshold theorem.** We consider in this section a family of probability measures indexed by a parameter  $p \in (0, 1)$ , and we prove a sharp-threshold theorem for this family subject to a hypothesis of monotonicity. The motivating example is the random-cluster model, to which we return in Sections 5 and 6.

Let  $1 \leq N < \infty$ ,  $I = \{1, 2, \dots, N\}$ , and let  $\Omega = \{0, 1\}^N$  and  $\mathcal{F}$  be given as before. Let  $\mu$  be a positive probability measure on  $(\Omega, \mathcal{F})$ . For  $p \in (0, 1)$ , we define the probability measure  $\mu_p$  by

$$(3.1) \quad \mu_p(\omega) = \frac{1}{Z_p} \mu(\omega) \left\{ \prod_{i \in I} p^{\omega(i)} (1-p)^{1-\omega(i)} \right\}, \quad \omega \in \Omega,$$

where  $Z_p$  is the normalizing constant

$$(3.2) \quad Z_p = \sum_{\omega \in \Omega} \mu(\omega) \left\{ \prod_{i \in I} p^{\omega(i)} (1-p)^{1-\omega(i)} \right\}.$$

It is immediate that  $\mu_p$  is positive and that  $\mu = \mu_{\frac{1}{2}}$ . It is easy to check that  $\mu_p$  satisfies the FKG lattice condition (2.1) if and only if  $\mu$  satisfies this condition, and it follows that  $\mu$  is monotonic if and only if, for all  $p \in (0, 1)$ , [or, equivalently, for *some*  $p \in (0, 1)$ ],  $\mu_p$  is monotonic. In order to prove a sharp-threshold theorem for the family  $\mu_p$ , we present first a Russo-type formula.

**THEOREM 3.3** ([3]). *For any event  $A \in \mathcal{F}$ ,*

$$(3.4) \quad \frac{d}{dp} \mu_p(A) = \frac{1}{p(1-p)} \sum_{i \in I} \text{cov}_p(X_i, 1_A),$$

where  $\text{cov}_p$  denotes covariance with respect to the measure  $\mu_p$ .

**PROOF.** This may be obtained exactly as in [3], Proposition 4, see also Section 2.4 of [14]. The details are omitted.  $\square$

Let  $\mathcal{A}$  be a subgroup of the permutation group  $\Pi_N$ . A probability measure  $\phi$  on  $(\Omega, \mathcal{F})$  is called  $\mathcal{A}$ -invariant if  $\phi(\omega) = \phi(\alpha\omega)$  for all  $\alpha \in \mathcal{A}$ . An event  $A \in \mathcal{F}$  is called  $\mathcal{A}$ -invariant if  $A = \alpha A$  for all  $\alpha \in \mathcal{A}$ . It is easily seen that, for any subgroup  $\mathcal{A}$ ,  $\mu$  is  $\mathcal{A}$ -invariant if and only if each  $\mu_p$  is  $\mathcal{A}$ -invariant.

**THEOREM 3.5** (Sharp threshold). *There exists a constant  $c \in (0, \infty)$  such that the following holds. Let  $N \geq 1$  and let  $A \in \mathcal{F}$  be an increasing event. Let  $\mu$  be a positive probability measure on  $(\Omega, \mathcal{F})$  which is monotonic. If there exists a subgroup  $\mathcal{A}$  of  $\Pi_N$  acting transitively on  $I$  such that  $\mu$  and  $A$  are  $\mathcal{A}$ -invariant, then*

$$(3.6) \quad \frac{d}{dp} \mu_p(A) \geq \frac{c\xi_p}{p(1-p)} \min\{\mu_p(A), 1 - \mu_p(A)\} \log N, \quad p \in (0, 1),$$

where  $\xi_p = \mu_p(X_1)(1 - \mu_p(X_1))$ .

**PROOF OF THEOREM 3.5.** Let

$$I_{p,A}(i) = \mu_p(A \mid X_i = 1) - \mu_p(A \mid X_i = 0),$$

so that

$$\begin{aligned} \text{cov}_p(X_i, \mathbf{1}_A) &= \mu_p(X_i \mathbf{1}_A) - \mu_p(X_i) \mu_p(A) \\ &= \mu_p(X_i) (1 - \mu_p(X_i)) I_{p,A}(i). \end{aligned}$$

Under the given conditions,  $\mu_p(X_i) = \mu_p(X_j)$  and  $I_{p,A}(i) = I_{p,A}(j)$  for all  $i, j \in I$ . Summing over the index set  $I$  as in (3.4), we deduce (3.6) by Theorem 2.10 applied to the monotonic measure  $\mu_p$ . This is the only place we have used the assumption of monotonicity.  $\square$

**4. Probability measures on the Euclidean cube.** We have so far considered probability measures on the discrete cube  $\{0, 1\}^N$  only. The method of proof of the influence theorem, Theorem 2.10, may be applied also to probability measures on the Euclidean cube  $[0, 1]^N$  that are absolutely continuous with respect to Lebesgue measure. Any such measure  $\mu$  has a density function  $\rho$ , which is to say that

$$\mu(A) = \int_A \rho(\mathbf{x}) \lambda(d\mathbf{x}),$$

for (Lebesgue) measurable subsets  $A$  of  $[0, 1]^N$ , with  $\lambda$  denoting Lebesgue measure. Since the density function  $\rho$  is non-unique, we shall phrase the results of this section in terms of  $\rho$  rather than the associated measure  $\mu$ . Some may regard this as not entirely satisfactory, arguing that results for *measures* should be based on hypotheses for these measures, rather than for particular versions of their density functions. One may rewrite the conclusions of this section thus, but at the expense of greater measure-theoretic detail which obscures the basic argument.

Let  $N \geq 1$ , and write  $\Omega = [0, 1]^N$ . Let  $\rho : \Omega \rightarrow [0, \infty)$  be (Lebesgue) measurable. We call  $\rho$  a *density function* if

$$\int_{\Omega} \rho(\mathbf{x}) \lambda(d\mathbf{x}) = 1,$$

and in this case we denote by  $\mu_\rho$  the corresponding probability measure,

$$\mu_\rho(A) = \int_A \rho(\mathbf{x}) \lambda(d\mathbf{x}).$$

We call  $\rho$  *positive* if it is a strictly positive function on  $\Omega$ , and we say it satisfies the (*continuous*) *FKG lattice condition* if

$$(4.1) \quad \rho(\mathbf{x} \vee \mathbf{y}) \rho(\mathbf{x} \wedge \mathbf{y}) \geq \rho(\mathbf{x}) \rho(\mathbf{y}) \quad \text{for all } \mathbf{x}, \mathbf{y} \in \Omega,$$

where the operations  $\vee, \wedge$  are defined as the coordinate-wise maximum and minimum, respectively.

Let  $\rho$  be a density function. We call  $\mu_\rho$  *positively associated* if

$$\mu_\rho(A \cap B) \geq \mu_\rho(A)\mu_\rho(B),$$

for all increasing subsets of  $\Omega$ . [It is presumably well known that increasing subsets of  $\Omega$  are Lebesgue-measurable but need not be Borel-measurable; see Theorem 4.13 and the subsequent remark.]

Let  $I = \{1, 2, \dots, N\}$ . For  $J \subseteq I$ , let  $\Omega_J = [0, 1]^J$  and

$$(4.2) \quad \Omega_J^\xi = \{\mathbf{x} \in \Omega : x_j = \xi_j \text{ for } j \in I \setminus J\}, \quad \xi \in \Omega.$$

The Lebesgue  $\sigma$ -algebra of  $\Omega_J$  is denoted by  $\mathcal{F}_J$ . Let  $\rho$  be a positive density function. We define the conditional probability measure  $\mu_{\rho, J}^\xi$  on  $(\Omega_J, \mathcal{F}_J)$  by

$$(4.3) \quad \mu_{\rho, J}^\xi(E) = \int_E \rho_J^\xi(\mathbf{x}) \lambda(d(x_j : j \in J)), \quad E \in \mathcal{F}_J,$$

where  $\rho_J^\xi$  is the conditional density function

$$\rho_J^\xi(\mathbf{x}) = \frac{1}{Z_J^\xi} \rho(\mathbf{x}) 1_{\Omega_J^\xi}(\mathbf{x}), \quad Z_J^\xi = \int_{\Omega_J^\xi} \rho(\mathbf{x}) \lambda(d(x_j : j \in J)).$$

We sometimes write  $\mu_\rho(E \mid (\xi_i : i \in I \setminus J))$  for  $\mu_{\rho, J}^\xi(E)$ , and we recall the standard fact that  $\mu_\rho(\cdot \mid (\xi_i : i \in I \setminus J))$  is a version of the conditional expectation given the  $\sigma$ -field  $\mathcal{F}_{I \setminus J}$ .

We say that  $\rho$  is *strongly positively-associated* if: for all  $J \subseteq I$  and all  $\xi \in \Omega$ , the measure  $\mu_{\rho, J}^\xi$  is positively associated. We call  $\rho$  *monotonic* if: for all  $J \subseteq I$ , all increasing subsets  $A$  of  $\Omega_J$ , and all  $\xi, \zeta \in \Omega$ ,

$$(4.4) \quad \mu_{\rho, J}^\xi(A) \leq \mu_{\rho, J}^\zeta(A) \quad \text{whenever } \xi \leq \zeta,$$

which is to say that, for all  $J \subseteq I$ ,

$$(4.5) \quad \mu_{\rho, J}^\xi \leq_{\text{st}} \mu_{\rho, J}^\zeta \quad \text{whenever } \xi \leq \zeta.$$

Here is a basic result concerning stochastic ordering.

**THEOREM 4.6** ([1, 17]). *Let  $N \geq 1$ , and let  $f$  and  $g$  be density functions on  $\Omega = [0, 1]^N$ . If*

$$g(\mathbf{x} \vee \mathbf{y})f(\mathbf{x} \wedge \mathbf{y}) \geq g(\mathbf{x})f(\mathbf{y}) \quad \text{for all } \mathbf{x}, \mathbf{y} \in [0, 1]^N,$$

*then  $\mu_f \leq_{\text{st}} \mu_g$ .*

If  $\rho$  satisfies the FKG lattice condition and  $A$  is an increasing event, then

$$1_A(\mathbf{x} \vee \mathbf{y})\rho(\mathbf{x} \vee \mathbf{y})\rho(\mathbf{x} \wedge \mathbf{y}) \geq 1_A(\mathbf{x})\rho(\mathbf{x})\rho(\mathbf{y}),$$

whence, by Theorem 4.6,

$$\mu_\rho(A)\mu_\rho(B) \leq \mu_\rho(A \cap B)$$

for all increasing  $A, B$ . Therefore,  $\mu_\rho$  is positively associated.

Henceforth we restrict ourselves to *positive* density functions. Arguments similar to the above are valid with  $\rho$  (assumed positive) replaced by the conditional density function  $\rho_J^\xi$ , and one arrives thus at the following.

**THEOREM 4.7.** *Let  $N \geq 1$ , and let  $\rho$  be a positive density function on  $\Omega = [0, 1]^N$  satisfying the FKG lattice condition (4.1). Then  $\rho$  is strongly positively-associated and monotonic.*

We turn now to a ‘continuous’ version of Theorem 2.10. Let  $N \geq 1$ , and let  $\rho$  be a monotonic positive density function on  $\Omega = [0, 1]^N$ . Let  $U = (U_1, U_2, \dots, U_N)$  be the identity function on  $[0, 1]^N$ . For an increasing subset  $A$  of  $\Omega$ , we define the *conditional influences* by

$$(4.8) \quad I_A(i) = \mu_\rho(A \mid U_i = 1) - \mu_\rho(A \mid U_i = 0), \quad i \in I.$$

**THEOREM 4.9 (Influence).** *There exists a constant  $c \in (0, \infty)$  such that the following holds. Let  $N \geq 1$  and let  $A$  be an increasing subset of  $\Omega = [0, 1]^N$ . Let  $\rho$  be a positive density function on  $[0, 1]^N$  that is monotonic. There exists  $i \in I$  such that*

$$(4.10) \quad I_A(i) \geq c \min\{\mu(A), 1 - \mu(A)\} \frac{\log N}{N}.$$

**PROOF.** The proof is very similar to that of Theorem 2.10. We propose first to construct an increasing event  $B$  such that  $\lambda(B) = \mu(A)$ , by way of a function  $f : [0, 1]^N \rightarrow [0, 1]^N$ . Let  $\mathbf{x} = (x_i : 1 \leq i \leq N) \in [0, 1]^N$ , and write  $f(\mathbf{x}) = (f_1(\mathbf{x}), f_2(\mathbf{x}), \dots, f_N(\mathbf{x}))$ . The first coordinate  $f_1(\mathbf{x})$  depends on  $x_1$  only and is defined by:

$$\mu_\rho(U_1 > f_1(\mathbf{x})) = 1 - x_1.$$

Since the density function  $\rho$  is strictly positive,  $f_1(\mathbf{x})$  is a continuous and strictly increasing function of  $x_1$ . It is an elementary exercise to check that the law of  $f_1(U)$  under  $\lambda$  is the same as that of  $U_1$  under  $\mu_\rho$ .

Having defined  $f_1(\mathbf{x})$ , we define  $f_2(\mathbf{x})$  in terms of  $x_1, x_2$  only by:

$$\mu_\rho(U_2 > f_2(\mathbf{x}) \mid U_1 = f_1(\mathbf{x})) = 1 - x_2.$$

The left-hand side is defined according to (4.3). It is a standard fact that  $\mu_\rho(\cdot \mid U_1 = f_1)$  is a version of the conditional expectation  $\mu_\rho(\cdot \mid \sigma(U_1))$ , where  $\sigma(U_1)$  denotes the  $\sigma$ -field generated by  $U_1$ , and it is an exercise to check that the pair  $(f_1(U), f_2(U))$  has the same law under  $\lambda$  as does the pair  $(U_1, U_2)$  under  $\mu_\rho$ . For each given  $x_1 \in (0, 1)$ ,  $f(\mathbf{x})$  is a continuous and strictly increasing function of  $x_2$ . [We use the assumptions that  $\rho$  is positive and monotonic, respectively, here.]

We continue inductively. Suppose we know  $f_i(\mathbf{x})$  for  $1 \leq i < k$ . Then  $f_k(\mathbf{x})$  depends on  $x_1, x_2, \dots, x_k$  and is given by:

$$\mu_\rho(U_k > f_k(\mathbf{x}) \mid U_i = f_i(\mathbf{x}) \text{ for } 1 \leq i < k) = 1 - x_k.$$

As above,  $f$  is strictly increasing (using the assumption of monotonicity), and the law of  $f(U)$  under  $\lambda$  is the same as the law of  $U$  under  $\mu_\rho$ . We set  $B = f^{-1}(A)$ .

Let

$$J_B(i) = \lambda(B \mid U_i = 1) - \lambda(B \mid U_i = 0), \quad i \in I.$$

Since  $f_1$  is continuous and strictly increasing,

$$\mu_\rho(A \mid U_1 = b) = \lambda(B \mid f_1(U_1) = b) = \lambda(B \mid U_1 = b), \quad b = 0, 1,$$

implying that  $I_A(1) = J_B(1)$ . It remains to show that  $I_A(j) \geq J_B(j)$  for  $j \in I$ . Let  $j \in I, j \neq 1$ . We re-order the coordinate set as  $K = \{j, 1, 2, \dots, j-1, j+1, \dots, N\}$ , and we construct a continuous increasing function  $g$  as above but subject to the new ordering. Rather than re-work the details from the proof of Theorem 2.10, we prove only part of that necessary. We sketch a proof that  $\mu_\rho(A \mid U_j = 1) \geq \lambda(B \mid U_j = 1)$ , a similar argument being valid with 1 replaced by 0 and the inequality reversed. The main step is to show that  $f \leq g$  under the assumption that  $U_j = 1$ . Suppose that  $1 \leq r < j$ , and assume it has already been proved that  $f_i(\mathbf{x}) \leq g_i(\mathbf{x})$  for  $\mathbf{x} \in \Omega$  and  $1 \leq i < r$ . Let  $\mathbf{x} \in \Omega$ . We claim that

$$(4.11) \quad \begin{aligned} &\mu_\rho(U_r > \xi \mid U_i = f_i(\mathbf{x}) \text{ for } 1 \leq i < r) \\ &\leq \mu_\rho(U_r > \xi \mid U_j = 1, U_i = g_i(\mathbf{x}) \text{ for } 1 \leq i < r), \quad \xi \in [0, 1]. \end{aligned}$$

By monotonicity,

$$(4.12) \quad \begin{aligned} &\mu_{\rho, J}(\cdot \mid U_j = u, U_i = f_i(\mathbf{x}) \text{ for } 1 \leq i < r) \\ &\leq_{\text{st}} \mu_{\rho, J}(\cdot \mid U_j = 1, U_i = g_i(\mathbf{x}) \text{ for } 1 \leq i < r), \quad u \in [0, 1]. \end{aligned}$$

The left-hand side of (4.12) is a version of the conditional expectation of the conditional measure  $\mu_{\rho, \mathcal{J}}(\cdot \mid U_i = f_i(\mathbf{x})$  for  $1 \leq i < r$ ) given  $\sigma(U_j)$ . By averaging over the value of  $u$  in (4.12), we obtain (4.11). The other steps are proved similarly.  $\square$

Unlike the discrete setting of Section 3, Theorem 4.9 does not imply a sharp-threshold theorem of the form of Theorem 3.5. Any density function  $\rho$  on  $[0, 1]^N$  may be used to generate a parametric family  $(\rho_p : 0 < p < 1)$  of densities given by

$$\rho_p(\mathbf{x}) = \frac{1}{Z_{\rho, p}} \rho(\mathbf{x}) \prod_{i=1}^N p^{x_i} (1-p)^{1-x_i}, \quad \mathbf{x} = (x_1, x_2, \dots, x_N) \in [0, 1]^N,$$

and we write  $\mu_p = \mu_{\rho_p}$ . Let  $A$  be an increasing subset of  $[0, 1]^N$ . The proof of Theorem 3.3 may be adapted to this setting to obtain that

$$\frac{d}{dp} \mu_p(A) = \frac{1}{p(1-p)} \sum_{i=1}^N \text{cov}_p(U_i, 1_A),$$

where  $U = (U_1, U_2, \dots, U_N)$  is the identity function on  $[0, 1]^N$ , and  $\text{cov}_p$  denotes covariance with respect to  $\mu_p$ .

Let  $\rho$  be a non-zero constant function, so that  $\mu_\rho$  is Lebesgue measure. As above, let  $p \in (0, 1)$  and let  $Y_1, Y_2, \dots, Y_N$  be independent random variables taking values in  $[0, 1]$  with common density function

$$\rho_p(x) = \begin{cases} \frac{\log[p/(1-p)]}{2p-1} p^x (1-p)^{1-x} & \text{if } p \neq \frac{1}{2}, x \in (0, 1), \\ 1 & \text{if } p = \frac{1}{2}, x \in (0, 1). \end{cases}$$

It is easily checked that the joint density function

$$\rho_p(\mathbf{x}) = \prod_{i=1}^N \rho_p(x_i), \quad \mathbf{x} = (x_1, x_2, \dots, x_N) \in [0, 1]^N,$$

satisfies the FKG lattice condition, and is therefore monotonic.

Let  $A = (N^{-1}, 1]^N$ . It is an easy calculation that

$$\mu_p(A) = \begin{cases} \left(1 - \frac{\pi^{1/N} - 1}{\pi - 1}\right)^N & \text{if } p \neq \frac{1}{2}, \\ \left(1 - \frac{1}{N}\right)^N & \text{if } p = \frac{1}{2}, \end{cases}$$

where  $\pi = p/(1-p)$ . Therefore, as  $N \rightarrow \infty$ ,

$$\mu_p(A) \rightarrow \begin{cases} \pi^{-1/(\pi-1)} & \text{if } p \neq \frac{1}{2}, \\ e^{-1} & \text{if } p = \frac{1}{2}. \end{cases}$$

In addition,

$$\text{cov}_{\frac{1}{2}}(U_i, 1_A) = \frac{1}{N} \left(1 - \frac{1}{N}\right)^{N-1} \sim \frac{e^{-1}}{N}.$$

The influence theorem, Theorem 4.9, may be applied to the event  $A$ , but there is no sharp threshold for  $\mu_p(A)$ . This situation diverges from that of the discrete setting at the point where a lower bound for the conditional influence  $I_A(i)$  is used to calculate a lower bound for the covariance  $\text{cov}_p(U_i, 1_A)$ .

We return briefly to the measurability of an increasing subset of  $[0, 1]^N$ .

**THEOREM 4.13.** *Let  $N \geq 2$ . Every increasing subset of  $[0, 1]^N$  is Lebesgue-measurable.*

Increasing subsets need not be Borel-measurable, as the following example indicates. Let  $M$  be a non-Borel-measurable subset of  $[0, 1]$ . Consider the increasing subset  $A$  of  $[0, 1]^2$  given by

$$A = \{(x, y) \in [0, 1]^2 : x + y > 1\} \cup \{(x, 1 - x) : x \in M\}.$$

The function  $h : x \mapsto (x, 1 - x)$  is a continuous, and hence Borel-measurable, function from  $\mathbb{R}$  to  $\mathbb{R}^2$ . If  $A$  were Borel-measurable, then so would be

$$A' = A \cap \{(x, 1 - x) : x \in \mathbb{R}\} = \{(x, 1 - x) : x \in M\}.$$

This would imply that  $h^{-1}(A') = M$  is Borel-measurable, a contradiction.

**PROOF OF THEOREM 4.13.** The statement is trivially true when  $N = 1$ , and we prove the general case by induction on  $N$ . Suppose  $n$  is such that the result holds for  $N = n$ . Let  $A$  be an increasing subset of  $[0, 1]^{n+1}$ , and let  $g : [0, 1]^n \rightarrow [0, 1] \cup \{\infty\}$  be defined by

$$g(\mathbf{x}) = \inf\{y : (\mathbf{x}, y) \in A\}, \quad \mathbf{x} \in [0, 1]^n.$$

The function  $g$  is decreasing on  $[0, 1]^n$ , and hence, for all  $c \in \mathbb{R}$ , the subset  $H_c = \{\mathbf{x} : g(\mathbf{x}) < c\}$  is increasing. By the induction hypothesis, each  $H_c$  is Lebesgue-measurable in  $[0, 1]^n$ , and therefore  $g$  is a measurable function. Its graph  $G = \{(\mathbf{x}, g(\mathbf{x})) : \mathbf{x} \in [0, 1]^n\}$  is (by an approximation by simple

functions, or otherwise) a Lebesgue-measurable set and is also (by Fubini's Theorem) a null subset of  $[0, 1]^{n+1}$ . Furthermore, the set

$$\bar{A} = \{(\mathbf{x}, y) \in [0, 1]^{n+1} : y > g(\mathbf{x})\}$$

is Lebesgue-measurable. Now  $A$  differs from  $\bar{A}$  only on a subset of the null set  $G$ , and the claim follows.  $\square$

**5. The random-cluster model.** The sharp-threshold theorem of Section 3 may be applied as follows to the random-cluster measure. Let  $G = (V, E)$  be a finite graph, assumed for simplicity to have neither loops nor multiple edges. We take as configuration space the set  $\Omega = \{0, 1\}^E$ , and write  $\mathcal{F}$  for the set of its subsets. For  $\omega \in \Omega$ , we call an edge  $e$  *open* (in  $\omega$ ) if  $\omega(e) = 1$ , and *closed* otherwise. Let  $\eta(\omega) = \{e \in E : \omega(e) = 1\}$  be the set of open edges, and consider the open graph  $G_\omega = (V, \eta(\omega))$ . The connected components of  $G_\omega$  are termed *open clusters*, and  $k(\omega)$  denotes the number of such clusters (including any isolated vertices).

Let  $q \in (0, \infty)$ , and let  $\mu$  be the probability measure on  $(\Omega, \mathcal{F})$  given by

$$(5.1) \quad \mu(\omega) = \frac{1}{Z(q)} q^{k(\omega)}, \quad \omega \in \Omega,$$

where  $Z(q)$  is the appropriate normalizing constant. It is clear that  $\mu$  is positive, and it is easily checked that  $\mu$  satisfies the FKG lattice condition if  $q \geq 1$ . See [8, 14]. (The FKG lattice condition does not hold when  $q < 1$  and  $G$  contains a circuit.) We assume henceforth that  $q \geq 1$ . By Theorem 2.7,  $\mu$  is monotonic.

The random-cluster measure  $\phi_{p,q}$  on the graph  $G$  with parameters  $p \in (0, 1)$  and  $q \in [1, \infty)$  is given as in (3.1) by

$$(5.2) \quad \phi_{p,q}(\omega) = \frac{1}{Z(p,q)} \left\{ \prod_{e \in E} p^{\omega(e)} (1-p)^{1-\omega(e)} \right\} q^{k(\omega)}, \quad \omega \in \Omega.$$

It is standard (see [8, 14]) that

$$(5.3) \quad \frac{p}{p+q(1-p)} \leq \phi_{p,q}(X_e = 1) \leq p, \quad e \in E.$$

Let  $\mathcal{A}$  be a subgroup of the automorphism group of  $G$ . We call  $G$   $\mathcal{A}$ -*transitive* if  $\mathcal{A}$  acts transitively on  $E$ . We may apply Theorem 3.5 to obtain the following. There exists an absolute constant  $c > 0$  such that, for all  $\mathcal{A}$ -transitive graphs  $G$ , all  $p, q$ , and any increasing  $\mathcal{A}$ -invariant event  $A \in \mathcal{F}$ ,

$$\frac{d}{dp} \phi_{p,q}(A) \geq c \min \left\{ \frac{q}{\{p+q(1-p)\}^2}, 1 \right\} \min\{\phi_{p,q}(A), 1 - \phi_{p,q}(A)\} \log N,$$

whence

$$(5.4) \quad \frac{d}{dp} \phi_{p,q}(A) \geq \frac{c}{q} \min\{\phi_{p,q}(A), 1 - \phi_{p,q}(A)\} \log N.$$

This differential inequality takes the usual simpler form when  $q = 1$ , and may be integrated as follows for general  $q \geq 1$ . Let  $p_1 \in (0, 1)$  be chosen such that  $\phi_{p_1,q}(A) \geq \frac{1}{2}$ , and let  $p_1 < p_2 < 1$ . We note that  $\phi_{p,q}(A) \geq \frac{1}{2}$  for  $p \in (p_1, p_2)$ , and we integrate (5.4) over this interval to obtain that

$$(5.5) \quad \phi_{p_2,q}(A) \geq 1 - \frac{1}{2} N^{-c(p_2-p_1)/q}.$$

For  $p \geq \sqrt{q}/(1 + \sqrt{q})$ , (5.4) may be replaced by

$$(5.6) \quad \frac{d}{dp} \phi_{p,q}(A) \geq c \min\{\phi_{p,q}(A), 1 - \phi_{p,q}(A)\} \log N,$$

and (5.5) becomes

$$(5.7) \quad \phi_{p_2,q}(A) \geq 1 - \frac{1}{2} N^{-c(p_2-p_1)}, \quad \frac{\sqrt{q}}{1 + \sqrt{q}} \leq p_1 < p_2,$$

under the condition  $\phi_{p_1,q}(A) \geq \frac{1}{2}$ . As an application of this inequality, we derive next a lower bound for the probability of an open crossing of a rectangle of  $\mathbb{Z}^2$ .

Let  $\mathbb{Z} = \{\dots, -1, 0, -1, \dots\}$  be the integers, and  $\mathbb{Z}^2$  the set of all 2-vectors  $x = (x_1, x_2)$  of integers. We turn  $\mathbb{Z}^2$  into a graph by placing an edge between any two vertices  $x, y$  with  $|x - y| = 1$ , where

$$|z| = |z_1| + |z_2|, \quad z \in \mathbb{Z}^2.$$

We write  $\mathbb{E}^2$  for the set of such edges, and  $\mathbb{L}^2 = (\mathbb{Z}^2, \mathbb{E}^2)$  for the ensuing graph. We shall work on a finite torus of  $\mathbb{L}^2$ . Let  $n \geq 1$ . Consider the square  $S_n = [0, n]^2$  (this is a convenient abbreviation for  $\{0, 1, 2, \dots, n\}^2$ ) viewed as a subgraph of  $\mathbb{L}^2$ . We identify certain pairs of vertices on the boundary of  $S_n$  in order to make it symmetric. More specifically, we identify any pair of the form  $(0, m), (n, m)$  and of the form  $(m, 0), (m, n)$ , for  $0 \leq m \leq n$ , and we merge any parallel edges that ensue. Let  $T_n = (V_n, E_n)$  denote the resulting toroidal graph. Let  $\mathcal{A}_n$  be the automorphism group of the graph  $T_n$ , and note that  $\mathcal{A}_n$  acts transitively on  $E_n$ . The configuration space of the random-cluster model on  $T_n$  is denoted by  $\Omega(n) = \{0, 1\}^{E_n}$ .

Let  $p \in (0, 1)$  and  $q \in [1, \infty)$ . Write  $\phi_{n,p}$  for the random-cluster measure on  $T_n$  with parameters  $p$  and  $q$ , and note that  $\phi_{n,p}$  is  $\mathcal{A}_n$ -invariant. Let

$$p_{\text{sd}} = p_{\text{sd}}(q) = \frac{\sqrt{q}}{1 + \sqrt{q}},$$

the self-dual point of the random-cluster model on  $\mathbb{L}^2$ , see [13, 14]. We note that the (Whitney) dual of  $T_n$  is isomorphic to  $T_n$ , and the random-cluster measure on  $T_n$  is self-dual when  $p = p_{\text{sd}}$ .

Let  $\omega \in \Omega(n)$ . Any translate in  $T_n$  of a rectangle of the form  $[0, r] \times [0, s]$  is said to be of size  $r \times s$ . When  $r \neq s$ , such a translate is said to be traversed *long-ways* (respectively, traversed *short-ways*) if the two shorter sides (respectively, longer sides) of the rectangle are joined within the rectangle by an open path of  $\omega$ .

Let  $\alpha \in (1, \infty)$ , and let  $\text{SW}_{n,\alpha}$  denote the event that the rectangle  $H_{n,\alpha} = [0, \lceil n\alpha \rceil] \times [0, \lfloor n/\alpha \rfloor]$  is crossed short-ways. One would normally take  $\alpha - 1$  to be small and  $n$  to be large in the next theorem.

**THEOREM 5.8.** *Let  $\alpha \in (1, \infty)$ ,  $k, n \geq 2$ ,  $q \in [1, \infty)$ , and  $p_{\text{sd}} < p < 1$ . Suppose that  $n/(n-1) \leq \alpha < \min\{k, n\}$ . We have that*

$$(5.9) \quad \phi_{kn,p}(\text{SW}_{n,\alpha}) \geq 1 - e^{-g(p-p_{\text{sd}})}$$

where

$$g = g(k, n, \alpha) = \frac{2c}{M} \log(kn),$$

and

$$M = 2 \left( 1 + \frac{k}{\alpha - 1} \right) \left( 1 + \frac{k\alpha}{\alpha - 1} \right).$$

Note that  $M$  is of order  $2k^2\alpha/(\alpha-1)^2$  for large  $k, n$ . For  $p > p_{\text{sd}}$ , one may make  $\phi_{kn,p}(\text{SW}_{n,\alpha})$  large by holding  $k$  fixed and sending  $n \rightarrow \infty$ . It does not seem to be easy to deduce an estimate for  $\phi_{p,q}(\text{SW}_{n,\alpha})$  for a random-cluster measure  $\phi_{p,q}$  on the infinite lattice  $\mathbb{L}^2$ . Neither do we know how to use the existence of crossings short-ways to build crossings long-ways. This is in contrast to the case of product measure, see [5, 7, 12, 18, 19, 21].

**PROOF.** Assume the given conditions. Let  $R_n = [0, n+1] \times [0, n]$ , viewed as a subgraph of  $T_{kn}$ , and let  $\text{LW}_n$  be the event that  $R_n$  is traversed long-ways. By a standard duality argument,

$$(5.10) \quad \phi_{kn,p_{\text{sd}}}(\text{LW}_n) = \frac{1}{2}, \quad k \geq 2, \quad n \geq 1.$$

Let  $A_n$  be the event that there exists in  $T_{kn}$  some translate of the square  $S_n = [0, n] \times [0, n]$  that possesses either an open top-bottom crossing or an open left-right crossing. The event  $A_n$  is  $\mathcal{A}_n$ -invariant, and

$$(5.11) \quad \phi_{kn,p_{\text{sd}}}(A_n) \geq \phi_{kn,p_{\text{sd}}}(\text{LW}_n) = \frac{1}{2}.$$

We apply (5.7) to the event  $A_n$ , with  $p_1 = p_{\text{sd}}$  and with  $N = 2(kn)^2$  being the number of edges in  $T_{kn}$ . This yields that

$$(5.12) \quad \begin{aligned} \phi_{kn,p}(A_n) &\geq 1 - \frac{1}{2}[2(kn)^2]^{-c(p-p_{\text{sd}})} \\ &\geq 1 - (kn)^{-2c(p-p_{\text{sd}})}, \quad p_{\text{sd}} < p < 1. \end{aligned}$$

The event  $A_n$  is defined on the whole of the torus. We next use an argument taken from [4, 5] to obtain a more locally defined event. Let  $a = \lceil n\alpha \rceil$ ,  $b = \lfloor n/\alpha \rfloor$ , and let  $H_{n,\alpha} = [0, a] \times [0, b]$  and  $V_{n,\alpha} = [0, b] \times [0, a]$ . Let  $h_{n,\alpha}$ ,  $v_{n,\alpha}$  be the sets of vertices in  $T_{kn}$  given by

$$\begin{aligned} h_{n,\alpha} &= \left\{ (l_1(a-n), l_2(n-b)) \in V_{kn} : 0 \leq l_1 < \frac{kn}{a-n}, 0 \leq l_2 < \frac{kn}{n-b} \right\}, \\ v_{n,\alpha} &= \left\{ (l_1(n-b), l_2(a-n)) \in V_{kn} : 0 \leq l_1 < \frac{kn}{n-b}, 0 \leq l_2 < \frac{kn}{a-n} \right\}, \end{aligned}$$

where the  $l_i$  are integers. That  $n-b \geq 1$  follows by the assumption  $\alpha \geq n/(n-1)$ . Consider the set  $\mathcal{H} = H_{n,\alpha} + h_{n,\alpha}$  of translates of  $H_{n,\alpha}$  by vectors in  $h_{n,\alpha}$ , and also the set  $\mathcal{V} = V_{n,\alpha} + v_{n,\alpha}$ . If  $A_n$  occurs, then some rectangle in  $\mathcal{H} \cup \mathcal{V}$  is traversed short-ways. By positive association and symmetry,

$$(5.13) \quad \begin{aligned} \phi_{kn,p}(\overline{A_n}) &\geq \phi_{kn,p}(\text{no member of } \mathcal{H} \cup \mathcal{V} \text{ is traversed short-ways}) \\ &\geq \{1 - \phi_{kn,p}(\text{SW}_{n,\alpha})\}^R, \end{aligned}$$

where  $\text{SW}_{n,\alpha}$  is the event that  $H_n$  is traversed short-ways, and

$$(5.14) \quad R = |h_{n,\alpha}| + |v_{n,\alpha}| \leq 2 \left\lceil \frac{kn}{a-n} \right\rceil \cdot \left\lceil \frac{kn}{n-b} \right\rceil.$$

After taking into account rounding effects, we find that  $R \leq M$ . Inequality (5.9) follows from (5.12)–(5.14).  $\square$

**6. The critical point.** There is a famous conjecture that the critical point  $p_c(q)$  of the random-cluster model on  $\mathbb{L}^2$  equals  $p_{\text{sd}}(q)$ . We do not spell out the details necessary to state this conjecture properly, referring the reader instead to [13, 14]. The conjecture is known to be valid for  $q = 1$  (percolation),  $q = 2$  (a case corresponding to the Ising model), and for sufficiently large  $q$  (namely  $q \geq 25.72$ ). The conjecture would follow if one could prove a strengthening of Theorem 5.8 in which short-ways is replaced by long-ways, and with the toroidal measure replaced by the wired measure on the full lattice. We finish by explaining this.

The so-called ‘wired random-cluster measure’ on  $\mathbb{L}^2$  is denoted by  $\phi_{p,q}^1$ , and the reader is referred to the references above for a definition of  $\phi_{p,q}^1$ .

THEOREM 6.1. *Let  $q \geq 1$ . Let  $p_k$  be the  $\phi_{p,q}^1$ -probability that a  $2^k \times 2^{k+1}$  rectangle is crossed long-ways. Suppose that*

$$(6.2) \quad \prod_{k=1}^{\infty} p_k > 0, \quad p > p_{\text{sd}}(q).$$

*Then the critical point of the random-cluster model on  $\mathbb{L}^2$  equals  $p_{\text{sd}}(q)$ .*

PROOF. We use a construction that appeared in [7]. For odd  $k$ , let  $A_k$  be the event that  $[0, 2^k] \times [0, 2^{k+1}]$  is traversed long-ways. For even  $k$ , let  $A_k$  be the event that  $[0, 2^{k+1}] \times [0, 2^k]$  is traversed long-ways. By the positive-association and automorphism-invariance of  $\phi_{p,q}^1$ , under (6.2),

$$\phi_{p,q}^1 \left( \bigcap_k A_k \right) \geq \prod_{k=1}^{\infty} \phi_{p,q}^1(A_k) > 0, \quad p > p_{\text{sd}}(q).$$

On the intersection of the  $A_k$ , there exists an infinite open cluster, and therefore  $p_c(q) \leq p_{\text{sd}}(q)$ . It is standard (see [13, 14]) that  $p_{\text{sd}}(q) \leq p_c(q)$ , and therefore equality holds as claimed.  $\square$

Let  $\phi_{p,q}^0$  denote the ‘free random-cluster measure’ on the square lattice  $\mathbb{L}^2$ . By duality,  $1 - p_k = \phi_{p',q}^0(\text{SW}(k))$ , where  $\text{SW}(k)$  is the event that the rectangle  $[0, 2^{k+1} - 1] \times [0, 2^k + 1]$  is traversed short-ways, and  $p'$  is the dual value of  $p$ ,

$$\frac{p'}{1 - p'} = \frac{q(1 - p)}{p}.$$

Therefore,

$$\begin{aligned} \sum_{k=1}^{\infty} (1 - p_k) &\leq \sum_{k=1}^{\infty} 2^{k+1} \phi_{p',q}^0(\text{rad}(C) \geq 2^k + 1) \\ &\leq 4 \sum_{n=1}^{\infty} \phi_{p',q}^0(\text{rad}(C) \geq n) \\ &= 4 \phi_{p',q}^0(\text{rad}(C)), \end{aligned}$$

where  $\text{rad}(C)$  is radius of the open cluster  $C$  at the origin, that is, the maximum value of  $n$  such that 0 is joined by an open path to the boundary of the box  $[-n, n]^2$ . It follows by Theorem 6.1 that

$$\phi_{p',q}^0(\text{rad}(C)) < \infty, \quad p' < p_{\text{sd}}(q),$$

is sufficient for  $p_c(q) = p_{\text{sd}}(q)$ .

**Acknowledgment.** The work of the first author was supported by the Engineering and Physical Sciences Research Council through a PhD studentship.

## REFERENCES

- [1] BATTY, C. J. K. AND BOLLMANN, H. W. (1980). Generalised Holley–Preston inequalities on measure spaces and their products. *Z. für Wahrsch'theorie verw. Geb.* 53, 157–173.
- [2] BEN-OR, M. AND LINIAL, N. (1990). Collective coin flipping. In *Randomness and Computation*. Academic Press, New York, 91–115.
- [3] BEZUIDENHOUT, C. E., GRIMMETT, G. R., AND KESTEN, H. (1993). Strict inequality for critical values of Potts models and random-cluster processes. *Comm. Math. Phys.* 158, 1–16.
- [4] BOLLOBÁS, B. AND RIORDAN, O. (2006a). The critical probability for random Voronoi percolation in the plane is  $\frac{1}{2}$ . *Probab. Theory Related Fields*. To appear.
- [5] BOLLOBÁS, B. AND RIORDAN, O. (2006b). A short proof of the Harris–Kesten theorem. *Bull. London Math. Soc.*. To appear.
- [6] BOURGAIN, J., KAHN, J., KALAI, G., KATZNELSON, Y., AND LINIAL, N. (1992). The influence of variables in product spaces. *Israel J. Math.* 77, 55–64.
- [7] CHAYES, J. T. AND CHAYES, L. (1986). Percolation and random media. In *Critical Phenomena, Random Systems and Gauge Theories (Les Houches, Session XLIII, 1984)*, K. Osterwalder and R. Stora, Eds. Elsevier, Amsterdam, 1001–1142.
- [8] FORTUIN, C. M. (1972). On the random-cluster model. III. The simple random-cluster process. *Physica* 59, 545–570.
- [9] FORTUIN, C. M., KASTELEYN, P. W., AND GINIBRE, J. (1971). Correlation inequalities on some partially ordered sets. *Comm. Math. Phys.* 22, 89–103.
- [10] FRIEDGUT, E. (2004). Influences in product spaces: KKL and BKKKL revisited. *Combin. Probab. Comput.* 13, 17–29.
- [11] FRIEDGUT, E. AND KALAI, G. (1996). Every monotone graph property has a sharp threshold. *Proc. Amer. Math. Soc.* 124, 2993–3002.
- [12] GRIMMETT, G. R. (1999). *Percolation*. Springer, Berlin.
- [13] GRIMMETT, G. R. (2003). The random-cluster model. In *Probability on Discrete Structures*, H. Kesten, Ed. Encyclopedia of Mathematical Sciences, Vol. 110. Springer, Berlin, 73–123.
- [14] GRIMMETT, G. R. (2006). *The Random-Cluster Model*. Springer, Berlin. To appear.
- [15] HÄGGSTRÖM, O., KALAI, G., AND MOSSEL, E. (2005). A law of large numbers for weighted majority. *Adv. Appl. Math.*. To appear.
- [16] KAHN, J., KALAI, G., AND LINIAL, N. (1988). The influence of variables on Boolean functions. In *Proceedings of 29th Symposium on the Foundations of Computer Science*. Computer Science Press, 68–80.
- [17] PRESTON, C. J. (1974). A generalization of the FKG inequalities. *Comm. Math. Phys.* 36, 233–241.
- [18] RUSSO, L. (1978). A note on percolation. *Z. für Wahrsch'theorie verw. Geb.* 43, 39–48.
- [19] RUSSO, L. (1981). On the critical percolation probabilities. *Z. für Wahrsch'theorie verw. Geb.* 56, 229–237.

- [20] RUSSO, L. (1982). An approximate zero-one law. *Z. für Wahrsch'theorie verw. Geb.* 61, 129–139.
- [21] SEYMOUR, P. D. AND WELSH, D. J. A. (1978). Percolation probabilities on the square lattice. In *Advances in Graph Theory*, B. Bollobás, Ed. Annals of Discrete Mathematics 3. North-Holland, Amsterdam, 227–245.
- [22] TALAGRAND, M. (1994). On Russo's approximate zero-one law. *Ann. Probab.* 22, 1576–1587.

STATISTICAL LABORATORY,  
UNIVERSITY OF CAMBRIDGE,  
WILBERFORCE ROAD,  
CAMBRIDGE CB3 0WB, U.K.  
E-MAIL: b.t.graham@statslab.cam.ac.uk  
E-MAIL: g.r.grimmett@statslab.cam.ac.uk  
URL: <http://www.statslab.cam.ac.uk/~grg>