

Classical coding via dequantizing

Frédéric Dupuis
Aarhus University

Joint work with

Oleg Szehr (*TU München*)

Marco Tomamichel (*National University of Singapore*)

arXiv:1207.0067

January 8, 2013


Introduction

- ▶ A purely quantum idea in information theory: Given a pure state $|\rho\rangle_{ABC}$, absence of correlations between A and B means that both A and B are perfectly correlated with C .
- ▶ If $\rho_{AB} = \rho_A \otimes \rho_B$, then $|\rho\rangle_{ABC} = |\rho_1\rangle_{AC_A} \otimes |\rho_2\rangle_{BC_B}$.
- ▶ Many coding theorems for sending *quantum* information can be proven using this idea (called “decoupling”).
- ▶ Doesn't seem to work for sending classical information: the information can be present in other systems without affecting the coding scheme.
- ▶ Our goal: come up with an analogue of this method for classical information.

- ▶ Coding via decoupling: a crash course
- ▶ From decoupling to dequantizing

Decoupling: a crash course


- ▶ The problem:



- ▶ Here $|\Phi\rangle = \frac{1}{\sqrt{N}} \sum_i |ii\rangle$.
- ▶ The hope: $\rho_{\widehat{M}M'} \approx \Phi_{MM'}$.


Decoupling: a crash course

The trick: Purify everything.




Decoupling: a crash course

We then hope that we obtain:




Decoupling: a crash course



- ▶ How do we get a decoder? The key: all purifications are equivalent up to a unitary on the purifying system.


Decoupling: a crash course



- ▶ $\Phi_{M'\widehat{M}} \otimes \widehat{\rho}_{E_D E}$ and $\rho_{M'EB}$ are both purifications of $\frac{\mathbb{1}_{M'}}{|M'|} \otimes \rho_E$.
- ▶ To get a decoder, we need to get $\frac{\mathbb{1}_{M'}}{|M'|} \otimes \rho_E$ at the output

Decoupling: a crash course

How do we ensure this?



- ▶ Encoding: embed M into a larger system, and then apply U chosen uniformly at random (a 2-design works too).
- ▶ If we have a product state $\frac{\mathbb{1}_{M'}}{|M'|} \otimes \rho_E$ at the output, then we know that a decoder exists.

Decoupling theorem


How do we ensure that M' is decoupled from the environment?

Theorem

Let $\mathcal{T}_{A \rightarrow E}$ be a channel. Then,


$$\mathbb{E}_{U_A} \left\| \mathcal{T}(U_A \rho_{A M'} U_A^\dagger) - \omega_E \otimes \frac{\mathbb{1}_{M'}}{|M'|} \right\|_1 \leq \sqrt{|M'|} 2^{-\frac{1}{2} H_{\min}^\varepsilon(A'|E)_\omega} + O(\varepsilon).$$

where $\omega_{A'E}$ is a state of the form:



The iid case

One important case: $\mathcal{T} = \mathcal{N}^{\otimes n}$:



Theorem

Let $\mathcal{T}_{A \rightarrow E}$ be a channel. Then,

$$\mathbb{E}_{U_A} \left\| \mathcal{T}(U_A \rho_{AM'} U_A^\dagger) - \omega_E \otimes \frac{\mathbb{1}_{M'}}{|M'|} \right\|_1 \leq \sqrt{|M'|} 2^{-\frac{1}{2} H_{\min}^\varepsilon(A'|E)_\omega} + O(\varepsilon).$$

- ▶ Fully quantum AEP:

$$H_{\min}^\varepsilon(A'|E)_{\omega^{\otimes n}} \rightsquigarrow nH(A'|E)_\omega = nI(A'\rangle B)_\omega.$$

- ▶ Hence, need $\log |M'| \lesssim nI(A'\rangle B)_\omega$.
- ▶ We therefore recover the Lloyd-Shor-Devetak theorem this way.

The family tree


Other coding theorems that can be proven this way:

- ▶ Entanglement-assisted quantum (and classical) capacity: start with $|\Phi\rangle_{MM'} \otimes |\Phi\rangle_{\tilde{A}\tilde{B}}$.
- ▶ State transfer/“Fully Quantum Slepian-Wolf”: Use \mathcal{T} to be a partial trace.
- ▶ State merging: Choose \mathcal{T} to be a measurement.
- ▶ Coding for quantum broadcast channels
- ▶ Etc...

What about sending classical information?

From decoupling to dequantizing

Sending classical information



where

- ▶ $T_{MM'} := \frac{1}{N} \sum_{i=1}^N |ii\rangle\langle ii|_{MM'}$
- ▶ \mathcal{N} is a CQ channel: $\mathcal{N}(\xi_A) = \sum_i \text{Tr}[|i\rangle\langle i|\xi_A] \rho_B^i$.

Sending classical information

To use decoupling ideas, we need to purify everything. How do we purify a CQ channel?


- ▶ $\mathcal{N}(\xi_A) = \sum_i \text{Tr}[|i\rangle\langle i|\xi_A]\rho_B^i$.
- ▶ $U_{\mathcal{N}}|i\rangle_A = |i\rangle_X \otimes |\rho^i\rangle_{BE}$.



- ▶ The environment of the channel is now XE .

Sending classical information


If we purify the entire process, we then get:



- Note that since the encoder \mathcal{E} is classical, $U_{\mathcal{E}}$ is just a permutation π of the basis elements.

Sending classical information


If we purify the entire process, we then get:



- ▶ Note that since the encoder \mathcal{E} is classical, $U_{\mathcal{E}}$ is just a permutation π of the basis elements.
- ▶ In the end, we want to ensure that $\widehat{\rho}_{M'\widehat{M}} = \frac{1}{N} \sum_i |ii\rangle\langle ii|$.


Sending classical information

What does that entail for the environment?



- ▶ We have that $|\widehat{\rho}\rangle = \frac{1}{\sqrt{N}} \sum_i |ii\pi(i)\rangle_{\widehat{M}M'X} \otimes |\xi^i\rangle_{EE_{\mathcal{D}}}$.
- ▶ This means that $\widehat{\rho}_{M'XE} = \frac{1}{N} \sum_i |i\pi(i)\rangle\langle i\pi(i)|_{M'X} \otimes \xi_E^i$.


Sending classical information



- ▶ Only classical correlations between M' and environment XE .
- ▶ So, we need $\rho_{M'XE} = \frac{1}{N} \sum_i |i\pi(i)\rangle\langle i\pi(i)|_{M'X} \otimes \xi_E^i$.



“Dequantizing” theorem

How can we enforce this?



- ▶ Encoder π is a permutation over all computation basis elements $|i\rangle$, chosen uniformly over the symmetric group.

“Dequantizing” theorem



- ▶ Encoder π is a permutation over all computation basis elements $|i\rangle$, chosen uniformly over the symmetric group.
- ▶ Look at the difference between left and right diagrams.


“Dequantizing” theorem

Theorem

Let $\bar{\mathcal{N}}_{A \rightarrow XE}$ be a complementary CQ channel. Then,

$$\mathbb{E}_{\pi} \left\| \bar{\mathcal{N}} \left(\pi_A (\Phi_{M'A} - T_{M'A}) \pi_A^{\dagger} \right) \right\|_1 \leq \sqrt{\frac{|M'|}{|A|}} 2^{-\frac{1}{2} H_{\min}^{\varepsilon}(A'|EX)_{\omega}} + O(\varepsilon).$$

where $\omega_{A'EXB}$ is a state of the form



Theorem

Let $\bar{\mathcal{N}}_{A \rightarrow XE}$ be a complementary CQ channel. Then,

$$\mathbb{E}_{\pi_A \in \mathcal{P}(A)} \left\| \bar{\mathcal{N}} \left(\pi_A (\Phi_{M'A} - T_{M'A}) \pi_A^\dagger \right) \right\|_1 \leq \sqrt{\frac{|M'|}{|A'|}} 2^{-\frac{1}{2} H_{\min}(A'|EX)_\omega}.$$

- ▶ Suppose we apply this theorem to an iid channel.
- ▶ Choose typical subspace: $\log |A'| \rightsquigarrow nH(A')$.
- ▶ Quantum AEP:
 $H_{\min}^\varepsilon(A'|EX)_\omega \rightsquigarrow nH(A'|EX) = -nH(A'|B)$.
- ▶ Hence, $\log |A'| + H_{\min}^\varepsilon(A'|EX) \rightsquigarrow nI(A'; B)$.
- ▶ Need $\log |M'| \lesssim nI(A'; B)$

Conclusion

- ▶ We have a “decoupling-like” proof of the HSW theorem.
- ▶ Can this method be fruitfully applied to other problems?
- ▶ Can we get cleaner proofs of “mixed” problems (classical and quantum messages at the same time)?

Thank you

Thank you!