

Email corrections to: N.Berestycki@statslab.cam.ac.uk

Example Sheet 1

1. Let Q be a Q -matrix on a finite state space S , and let $f(t) = \det P(t)$ where $P(t)$ is the associated transition matrix. Show that $f(t+s) = f(t)f(s)$. Deduce that $f(t)$ is of the form e^{tq} , and identify q by taking $t \rightarrow 0$.

Show the same result directly.

2. Let $(q_k)_{k \geq 1}$ be a sequence of positive numbers such that $q = \sum_k q_k < \infty$. Let $(E_k)_{k \geq 1}$ be a sequence of independent exponential random variables where the rate (i.e., parameter) of E_k is q_k . Let $E = \inf\{E_k\}$ and let $K = \arg \min\{E_k\}$, with $K = \infty$ if the inf is not attained. Compute $\mathbb{P}(K = k, E > t)$ for all $t > 0$ and $k \geq 1$, and hence identify the joint distribution of K and E .

Let Q be a Q -matrix on a countable state space S . After recalling the first two constructions of a Markov chain based on the holding times and jump chain, deduce from the above that the two constructions are equivalent.

3. Let T_1, T_2, \dots be independent exponential random variables of parameter λ . Show that, for all $n \geq 1$, the sum $S = \sum_{i=1}^n T_i$ has the probability density function

$$f_S(x) = \frac{\lambda^n x^{n-1}}{(n-1)!} e^{-\lambda x}, x > 0.$$

This is called the Gamma (n, λ) distribution.

Let N be an independent geometric random variable with

$$\mathbb{P}(N = n) = \beta(1 - \beta)^{n-1}, \quad n = 1, 2, \dots$$

Show that $T = \sum_{i=1}^N T_i$ has exponential distribution of parameter $\lambda\beta$. Deduce another proof of the thinning property of Poisson processes.

4. Arrivals of the Number 1 bus form a Poisson process of rate 1 bus per hour, and arrivals of the Number 7 bus form an independent Poisson process of rate 7 buses per hour.

(a) What is the probability that exactly 5 buses pass by in 1 hour?

(b) What is the probability that exactly 3 Number 7 buses pass by while I am waiting for a Number 1?

(c) When the maintenance depot goes on strike half the buses break down before they reach my stop. What then is the probability that I wait for 30 minutes without seeing a single bus?

5. Customers arrive in a supermarket as a Poisson-process of rate N . There are N aisles in the supermarket and each customer selects one of them at random, independently of the other customers. Let X_t^N denote the proportion of aisles which remain empty at time t and let T^N denote the time until half the aisles are busy (have at least one customer). Show that

$$X_t^N \rightarrow e^{-t} \quad , \quad T^N \rightarrow \log 2$$

in probability as $N \rightarrow \infty$.

6. A pedestrian wishes to cross a single lane of fast-moving traffic. Suppose the number of vehicles that have passed by time t is a Poisson process of rate λ , and suppose it takes time a to walk across the lane. Assuming the pedestrian can foresee correctly the times at which vehicles will pass by, how long on average does it take to cross over safely?

Hint: Let T be the time to cross and J_1 the time at which the first car passes. Identify the contributions to $\mathbb{E} T$ from the events $\{J_1 > a\}$ and $\{J_1 < a\}$.

How long on average does it take to cross two similar lanes (a) when one must walk straight across, (b) when an island in the middle of the road makes it safe to stop half way?

7. Customers enter a supermarket as a Poisson process of rate 2. There are two salesmen near the door who offer passing customers samples of a new product. Each customer takes an exponential time of parameter 1 to think about the new product, and during this time occupies the full attention of one salesman. Having tried the product, customers proceed into the store and leave by another door. When both salesmen are occupied, customers walk straight in. Assuming that both salesmen are free at time 0, find the probability that both are busy at a later time t .

8. Let $(X_t)_{t \geq 0}$ be a Markov chain on the integers with transition rates $q_{i,i+1} = \lambda q_i$, $q_{i,i-1} = \mu q_i$, and $q_{i,j} = 0$ if $|j-i| \geq 2$, where $\lambda + \mu = 1$ and $q_i > 0$ for all i . Write down the Q -matrix of this chain and draw the diagram corresponding to this chain. Then

(a) Find the probability, starting from 0, that X_t hits i , for any $i \geq 1$.

(b) Compute the expected total time spent in state i , starting from 0.

9. Let $(X_t)_{t \geq 0}$ be a birth-and-death process with rates $\lambda_n = n\lambda$ and $\mu_n = n\mu$, and suppose $X_0 = 1$. Show that $h(t) = \mathbb{P}(X_t = 0)$ satisfies

$$h(t) = \int_0^t e^{-(\lambda+\mu)s} \{\mu + \lambda h(t-s)\}^2 ds$$

and deduce that if $\lambda \neq \mu$ then

$$h(t) = (\mu e^{\mu t} - \mu e^{\lambda t}) / (\mu e^{\mu t} - \lambda e^{\lambda t}).$$

10. Each bacterium in a colony splits into two identical bacteria after an exponential time of parameter λ , which then split in the same way but independently. Let X_t denote the size of the colony at time t , and suppose $X_0 = 1$. Show that the probability generating function $\phi(t) = \mathbb{E}(z^{X_t})$ satisfies

$$\phi(t) = ze^{-\lambda t} + \int_0^t \lambda e^{-\lambda s} \phi(t-s)^2 ds$$

and deduce that, for $q = 1 - e^{-\lambda t}$, and $n = 1, 2, \dots$,

$$\mathbb{P}(X_t = n) = q^{n-1}(1 - q).$$

11. Compute $p_{11}(t)$ for $P(t) = e^{tQ}$, where

$$Q = \begin{pmatrix} -2 & 1 & 1 \\ 4 & -4 & 0 \\ 2 & 1 & -3 \end{pmatrix}.$$

Find an invariant distribution π for Q and verify that $p_{11}(t) \rightarrow \pi_1$ as $t \rightarrow \infty$.

12. Two fleas are bound together to take part in a nine-legged race on the vertices A, B, C of a triangle. Flea 1 hops at random times in the clockwise direction; each hop takes the pair from one vertex to the next and the times between successive hops of Flea 1 are independent random variables, each with exponential distribution, mean $1/\lambda$. Flea 2 behaves similarly, but hops in the anti-clockwise direction, the times between his hops having mean $1/\mu$. Show that the probability that they are at A at a given time $t > 0$ is

$$\frac{1}{3} + \frac{2}{3} \exp \left\{ -\frac{3(\lambda + \mu)t}{2} \right\} \cos \left\{ \frac{\sqrt{3}(\lambda - \mu)t}{2} \right\}.$$