

SELF-AVOIDING WALKS AND AMENABILITY

GEOFFREY R. GRIMMETT AND ZHONGYANG LI

ABSTRACT. The connective constant $\mu(G)$ of an infinite transitive graph G is the exponential growth rate of the number of self-avoiding walks from a given origin. The relationship between connective constants and amenability is explored in the current work.

Various properties of connective constants depend on the existence of so-called ‘graph height functions’, namely: (i) whether $\mu(G)$ is a local function on certain graphs derived from G , (ii) the equality of $\mu(G)$ and the asymptotic growth rate of bridges, and (iii) whether there exists a terminating algorithm for approximating $\mu(G)$ to a given degree of accuracy.

In the context of amenable groups, it is noted that the Cayley graphs of infinite, finitely generated, elementary amenable (and, more generally, virtually indicable) groups support graph height functions, which are in addition harmonic. In contrast, the Cayley graph of the Grigorchuk group, which is amenable but not elementary amenable, does not have a graph height function.

In the context of non-amenable, transitive graphs, a lower bound is presented for the connective constant in terms of the spectral bottom of the graph. This is a strengthening of an earlier result of the same authors. Secondly, using a percolation inequality of Benjamini, Nachmias, and Peres, it is explained that the connective constant of a non-amenable, transitive graph with large girth is close to that of a regular tree. Examples are given of non-amenable groups without graph height functions, of which one is the Higman group.

The emphasis of the work is upon the structure of Cayley graphs, rather than upon the algebraic properties of the underlying groups. New methods are needed since a Cayley graph generally possesses automorphisms beyond those arising through the action of the group.

Date: 24 November 2015, revised 21 March 2016.

2010 Mathematics Subject Classification. 05C30, 20F65, 60K35, 82B20.

Key words and phrases. Self-avoiding walk, connective constant, Cayley graph, amenable group, elementary amenable group, indicable group, Grigorchuk group, Higman group, Baumslag–Solitar group, graph height function, group height function, harmonic function, spectral radius, spectral bottom.

CONTENTS

Part A. Background, and summary of results	2
1. Introduction	2
1.1. Background	2
1.2. Amenable groups	3
1.3. Non-amenable graphs	4
2. Graphs, groups, and self-avoiding walks	5
2.1. Graphs	5
2.2. Self-avoiding walks	5
2.3. Groups	6
3. Height functions	6
Part B. Results for amenable groups	8
4. Elementary amenable groups	8
5. The Grigorchuk group	9
Part C. Results for non-amenable graphs	10
6. Connective constants of non-amenable graphs	10
7. Graphs with large girth	13
8. The Higman group	13
Part D. Remaining proofs	14
9. Criteria for the absence of height functions	14
10. Proof of Theorem 5.1	15
11. Proof of Theorem 8.1	17
12. Proof of Theorem 8.2	21
Acknowledgements	22
References	22

Part A. Background, and summary of results

1. INTRODUCTION

1.1. **Background.** A *self-avoiding walk* on a graph $G = (V, E)$ is a path that visits no vertex more than once. The study of the number σ_n of self-avoiding walks of length n from a given initial vertex was initiated by Flory [10] in his work on polymerization, and this topic has acquired an iconic status in the mathematics and physics associated with lattice-graphs. Hammersley and Morton [24] proved in 1954 that, if G is vertex-transitive, there exists a constant $\mu = \mu(G)$, called the *connective constant* of G , such

that $\sigma_n = \mu^{n(1+o(1))}$ as $n \rightarrow \infty$. This result is important not only for its intrinsic value, but also because its proof contained the introduction of subadditivity to the theory of interacting systems. Subsequent work has concentrated on understanding polynomial corrections in the above asymptotic for σ_n (see, for example, [3, 37]), and on finding exact values and inequalities for connective constants (for example, [9, 17]).

There are several natural questions about connective constants whose answers depend on whether or not the underlying graph admits a so-called graph height function. The first of these is whether $\mu(G)$ is a continuous function of the graph G (see [4, 18]). This so-called *locality* question has received serious attention also in the context of percolation and other disordered systems (see [5, 38, 40]), and has been studied in recent work of the current authors on general transitive graphs, [18], and also on Cayley graphs of finitely generated groups, [21]. Secondly, when G has a graph height function, one may define *bridge* self-avoiding walks on G , and show that their numbers grow asymptotically in the same manner as σ_n (see [18]). The third such question is whether there exists a terminating algorithm to approximate $\mu(G)$ within any given (non-zero) margin of accuracy (see [18, 19]).

Roughly speaking, a graph height function on $G = (V, E)$ is a non-constant function $h : V \rightarrow \mathbb{Z}$ whose increments are invariant under the action of a finite-index subgroup of automorphisms (a formal definition may be found at Definition 3.1). It is, therefore, useful to know which transitive graphs support graph height functions.

A method for constructing graph height functions on a certain class of transitive graphs is described in [18], and the question is posed there of deciding whether all transitive graphs support graph height functions. A rich source of interesting examples of transitive graphs is provided by Cayley graphs of finitely generated groups, as studied in [21]. It is proved there that the Cayley graphs of finitely generated, virtually solvable groups support graph height functions, which are in addition harmonic. The question is posed of determining whether or not the Cayley graph of the Grigorchuk group possesses a graph height function.

This work is directed primarily at the structure of *Cayley graphs*, rather than solely that of the underlying *groups*. The difference lies in the fact that a Cayley graph generally possesses graph-automorphisms that do not arise through the action of the corresponding group.

We are concerned here with the relationship between connective constants and amenability, and we present results for both amenable and for non-amenable graphs. Since these results are fairly distinct, we summarize them here under the two headings of amenable groups and non-amenable graphs.

1.2. Amenable groups. This part of the current work has two principal results, one positive and the other negative.

- (a) (Theorem 4.1) It is proved that every Cayley graph of an infinite, finitely generated, elementary amenable (and, more generally, virtually indicable) group supports a graph height function, which is in addition harmonic. This extends [21, Thm 5.1] beyond the class of virtually solvable groups.
- (b) (Theorem 5.1) It is proved that the Cayley graph of the Grigorchuk group does not support a graph height function. This answers in the negative the above question of [18] (see also [21, Sect. 5]). Since the Grigorchuk group is amenable (but not elementary amenable), possession of a graph height function is not a characteristic of amenable groups. This is in contrast with work of Lee and Peres, [34], who have studied the existence of non-constant, Hilbert space valued, equivariant harmonic maps on amenable graphs.

1.3. Non-amenable graphs. In earlier work [20], it was shown that the connective constant μ of a transitive, simple graph with degree Δ satisfies

$$\sqrt{\Delta - 1} \leq \mu \leq \Delta - 1,$$

and it was asked whether or not the lower bound is sharp. In the first of the following three results, this is answered in the negative for non-amenable graphs.

- (a) (Theorem 6.2) It is proved that

$$(\Delta - 1)^{\frac{1}{2}(1+c\lambda)} \leq \mu,$$

where $c = c(\Delta)$ is a known constant, and λ is the spectral bottom of the simple random walk on the graph. Kesten [32, 33] and Dodziuk [8] have shown that $\lambda > 0$ if and only if the graph is non-amenable.

- (b) (Theorem 7.1) Using a percolation result of Benjamini, Nachmias, and Peres [5], it is explained that the connective constant of a non-amenable, Δ -regular graph with large girth is close to that of the Δ -regular tree.
- (c) (Theorem 8.1) It is shown that the Cayley graph of the Higman group of [28] (which is non-amenable) does not support a graph height function. This further example of a transitive graph without a graph height function complements the corresponding statement above for the (amenable) Grigorchuk group.

Relevant notation for graphs, groups, and self-avoiding walks is summarized in Section 2, and three different types of height functions are explained in Section 3. The class EG of elementary amenable groups is described in Section 4. The Grigorchuk group is defined in Section 5 and the non-existence of graph height functions thereon is given in Theorem 5.1. The improved lower bound for $\mu(G)$ for non-amenable G is presented at Theorem 6.2, and the remark about non-amenable graphs with large girth at Theorem 7.1. The Higman group is discussed in Section 8. Proofs

of theorems appear either immediately after their statements, or are deferred to self-contained sections at the end of the article.

2. GRAPHS, GROUPS, AND SELF-AVOIDING WALKS

2.1. Graphs. The graphs $G = (V, E)$ in this paper are *simple*, in that they have neither loops nor multiple edges. The *degree* $\deg(v)$ of vertex $v \in V$ is the number of edges incident to v . We write $u \sim v$ for neighbours u and v , ∂v for the neighbour set of v , and $\partial_e v$ (respectively, $\partial_e W$) for the set of edges incident to v (respectively, between W and $V \setminus W$). The graph is *locally finite* if $|\partial v| < \infty$ for $v \in V$. An edge from u to v is denoted $\langle u, v \rangle$ when undirected, and $[u, v]$ when directed from u to v . The *girth* of G is the infimum of the lengths of its circuits. The infinite Δ -regular tree T_Δ crops up periodically in this paper.

The automorphism group of G is denoted $\text{Aut}(G)$. The subgroup $\Gamma \leq \text{Aut}(G)$ is said to act *transitively* on G if, for $u, v \in V$, there exists $\alpha \in \text{Aut}(G)$ with $\alpha(u) = v$. It acts *quasi-transitively* if there exists a finite subset $W \subseteq V$ such that, for $v \in V$, there exists $\alpha \in \Gamma$ and $w \in W$ such that $\alpha(v) = w$. The graph G is said to be *(vertex-)transitive* if $\text{Aut}(G)$ acts transitively on V .

Let \mathcal{G} be the set of infinite, locally finite, connected, transitive, simple graphs, and let $G \in \mathcal{G}$. The *edge-isoperimetric constant* $\phi = \phi(G)$ is defined here as

$$(2.1) \quad \phi := \inf \left\{ \frac{|\partial_e W|}{\Delta |W|} : W \subset V, 0 < |W| < \infty \right\}.$$

We call G *amenable* if $\phi = 0$ and *non-amenable* otherwise. See [35, Sect. 6] for an account of graph amenability.

2.2. Self-avoiding walks. Let $G \in \mathcal{G}$. We choose a vertex of G and call it the *origin*, denoted $\mathbf{1}$. An *n -step self-avoiding walk* (SAW) on G is a walk containing n edges no vertex of which appears more than once. Let Σ_n be the set of n -step SAWs starting at $\mathbf{1}$, with cardinality $\sigma_n := |\Sigma_n|$. We have in the usual way (see [24, 37]) that

$$(2.2) \quad \sigma_{m+n} \leq \sigma_m \sigma_n,$$

whence the *connective constant*

$$(2.3) \quad \mu = \mu(G) := \lim_{n \rightarrow \infty} \sigma_n^{1/n}$$

exists.

A SAW is called *extendable* if it is the initial portion of an infinite SAW on G . (An extendable SAW is called ‘forward extendable’ in [16].)

2.3. Groups. Let Γ be a group with generator set S satisfying $|S| < \infty$ and $\mathbf{1} \notin S$, where $\mathbf{1} = \mathbf{1}_\Gamma$ is the identity element. We shall assume that $S^{-1} = S$, while noting that this was not assumed in [21]. We write $\Gamma = \langle S \mid R \rangle$ with R a set of relators (or relations, when convenient). Such a group is called *finitely generated*, and is called *finitely presented* if, in addition, $|R| < \infty$.

The *Cayley graph* of the presentation $\Gamma = \langle S \mid R \rangle$ is the simple graph $G = G(\Gamma, S)$ with vertex-set Γ , and an (undirected) edge $\langle \gamma_1, \gamma_2 \rangle$ if and only if $\gamma_2 = \gamma_1 s$ for some $s \in S$. Thus, our Cayley graphs are simple graphs. See [2, 35] for accounts of Cayley graphs, and [26] of geometric group theory.

The amenability of groups was introduced by von Neumann [39]. It is standard that a finitely generated group is amenable if and only if some (and hence every) Cayley graph is amenable (see, for example, [42, Chap. 12A]).

3. HEIGHT FUNCTIONS

It was shown in [18] that graphs $G \in \mathcal{G}$ supporting so-called ‘graph height functions’ have (at least) three properties:

- (i) one may define the concept of a ‘bridge’ SAW on G , as in [25],
- (ii) the exponential growth rate for counts of bridges equals the connective constant $\mu(G)$,
- (iii) there exists a terminating algorithm for determining $\mu(G)$ to within any prescribed (strictly positive) degree of accuracy.

Several natural sub-classes of \mathcal{G} contain only graphs that support graph height functions, and it was asked in [20] whether or not *every* $G \in \mathcal{G}$ supports a graph height function. This question will be answered in the negative at Theorems 5.1 and 8.1, where it is proved that neither the Grigorchuk nor Higman graphs possess a graph height function. Arguments for proving the non-existence of graph height functions may be found in Section 9.

We review the definitions of the two types of height functions, and introduce a third type. Let $G = (V, E) \in \mathcal{G}$, and let $\mathcal{H} \leq \text{Aut}(G)$. A function $F : V \rightarrow \mathbb{R}$ is said to be *\mathcal{H} -difference-invariant* if

$$(3.1) \quad F(v) - F(w) = F(\gamma v) - F(\gamma w), \quad v, w \in V, \gamma \in \mathcal{H}.$$

Definition 3.1 ([18]). *A graph height function on G is a pair (h, \mathcal{H}) , where $\mathcal{H} \leq \text{Aut}(G)$ acts quasi-transitively on G and $h : V \rightarrow \mathbb{Z}$, such that:*

- (a) $h(\mathbf{1}) = 0$,
- (b) h is \mathcal{H} -difference-invariant,
- (c) for $v \in V$, there exist $u, w \in \partial v$ such that $h(u) < h(v) < h(w)$.

Remark 3.2. *By Poincaré's Theorem for subgroups (see [27, p. 48, Exercise 20]), it is immaterial whether or not we require the subgroup \mathcal{H} to be normal in Definition 3.1.*

We turn to Cayley graphs of finitely generated groups. Let Γ be a finitely generated group with presentation $\langle S \mid R \rangle$. As in Section 2, we assume $S^{-1} = S$ and $\mathbf{1} \notin S$.

Definition 3.3. *A group height function on Γ (or on a Cayley graph of Γ) is a function $h : \Gamma \rightarrow \mathbb{Z}$ such that:*

- (a) $h(\mathbf{1}) = 0$, and h is not identically zero,
- (b) if $\gamma = s_1 s_2 \cdots s_m$ with $s_i \in S$, then $h(\gamma) = \sum_{i=1}^m h(s_i)$,
- (c) the values $(h(s) : s \in S)$ are such that, if $s_1 s_2 \cdots s_n = \mathbf{1}$ is a representation of the identity with $s_i \in S$, then $\sum_{i=1}^n h(s_i) = 0$.

A necessary and sufficient condition for the existence of a group height function is given in [21, Thm 4.1]. In the language of group theory, this condition amounts to requiring that the first Betti number is strictly positive. It was recalled in [21, Remark 4.2] that (when the non-zero $h(s)$, $s \in S$, are coprime) a group height function is simply a surjective homomorphism from Γ to \mathbb{Z} .

It is shown in [18, Example 4.4] that the Cayley graph of the infinite dihedral group has a graph height function but no group height function. We introduce a third type of height function, which may be viewed as an intermediary between a graph height function and group height function.

Definition 3.4. *For a Cayley graph G of a finitely generated group Γ , we say that the pair (h, \mathcal{H}) is a strong graph height function of the pair (Γ, G) if*

- (a) $\mathcal{H} \trianglelefteq \Gamma$ acts on Γ by left multiplication, and $[\Gamma : \mathcal{H}] < \infty$, and
- (b) (h, \mathcal{H}) is a graph height function.

It is evident that a group height function h (of Γ) is a strong graph height function of the form (h, Γ) , and a strong graph height function is a graph height function. The assumption in (a) above of the normality of \mathcal{H} is benign, as in Remark 3.2.

We recall the definition of a harmonic function. A function $h : V \rightarrow \mathbb{R}$ is called *harmonic* on the graph $G = (V, E)$ if

$$h(v) = \frac{1}{\deg(v)} \sum_{u \sim v} h(u), \quad v \in V.$$

It is an exercise to show that any group height function is harmonic.

Part B. Results for amenable groups

4. ELEMENTARY AMENABLE GROUPS

The class EG of elementary amenable groups was introduced by Day in 1957, [7], as the smallest class of groups that contains the set EG_0 of all finite and abelian groups, and is closed under the operations of taking subgroups, and of forming quotients, extensions, and directed unions. Day noted that every group in EG is amenable (see also von Neumann [39]). An important example of an amenable but not elementary amenable group was described by Grigorchuk in 1984, [12]. Grigorchuk's group is important in the study of height functions, and we return to this in Section 5.

Let EFG be the set of infinite, finitely generated members of EG.

Theorem 4.1. *Let $\Gamma \in \text{EFG}$. There exists a normal subgroup $\mathcal{H} \trianglelefteq \Gamma$ with $[\Gamma : \mathcal{H}] < \infty$ such that any locally finite Cayley graph G of Γ possesses a harmonic, strong graph height function of the form (h, \mathcal{H}) .*

Whereas every member of EFG has a proper, normal subgroup with finite index, it is proved in [31] that there exist amenable *simple* groups. The class EFG includes all virtually solvable groups, and thus Theorem 4.1 extends [21, Thm 5.1]. Since any finitely generated group with polynomial growth is virtually nilpotent, [23], and hence lies in EFG, its locally finite Cayley graphs admit harmonic graph height functions.

Theorem 4.1 is a corollary of a more general result, as follows. A group Γ is called *indicible* if there exists a surjective homomorphism $F : \Gamma \rightarrow \mathbb{Z}$, and *virtually indicible* if there exists a normal subgroup $\mathcal{H} \trianglelefteq \Gamma$, with $[\Gamma : \mathcal{H}] < \infty$, which is indicible. (By Remark 3.2, the assumption of normality is immaterial.)

Theorem 4.2. *Let Γ be finitely generated and virtually indicible. There exists a normal subgroup $\mathcal{H} \trianglelefteq \Gamma$ with $[\Gamma : \mathcal{H}] < \infty$ such that any locally finite Cayley graph G of Γ possesses a harmonic, strong graph height function of the form (h, \mathcal{H}) .*

Proof of Theorem 4.2. Let $\mathcal{H} \trianglelefteq \Gamma$ be such that $[\Gamma : \mathcal{H}] < \infty$, and there exists a surjective homomorphism $F : \mathcal{H} \rightarrow \mathbb{Z}$. With F given thus, we may apply [21, Thm 3.5] to obtain the result. \square

Proof of Theorem 4.1. This may be proved by induction using the definition of elementary amenability, but we follow instead a much shorter route using indicability. The latter is standard, but we include some details to aid the reader.

Let $\Gamma \in \text{EFG}$. By [29, Lemma 1] or [30, Lemma 1.7], Γ has normal subgroups $K \trianglelefteq \mathcal{H} \trianglelefteq \Gamma$ such that $[\Gamma : \mathcal{H}] < \infty$ and \mathcal{H}/K is free abelian with positive rank. Therefore, there exists a surjective homomorphism $F : \mathcal{H}/K \rightarrow \mathbb{Z}$. The domain of F may be extended to \mathcal{H} by $F(h) := F(Kh)$, so that \mathcal{H} is indicible, and hence Γ is virtually indicible. The theorem now follows by Theorem 4.2. \square

5. THE GRIGORCHUK GROUP

The (first) Grigorchuk group is an infinite, finitely generated, amenable group that is not elementary amenable. We show in Theorem 5.1 that there exists a locally finite Cayley graph of the Grigorchuk group with *no* graph height function. This answers in the negative Question 3.3 of [18] (see also [21, Sect. 3]).

Here is the definition of the group in question (see [11, 12, 14]). Let T be the rooted binary tree with root vertex \emptyset . The vertex-set of T can be identified with the set of finite strings u having entries 0, 1, where the empty string corresponds to the root \emptyset . Let T_u be the subtree of all vertices with root labelled u .

Let $\text{Aut}(T)$ be the automorphism group of T , and let $a \in \text{Aut}(T)$ be the automorphism that, for each string u , interchanges the two vertices $0u$ and $1u$.

Any $\gamma \in \text{Aut}(G)$ may be applied in a natural way to either subtree T_i , $i = 0, 1$. Given two elements $\gamma_0, \gamma_1 \in \text{Aut}(T)$, we define $\gamma = (\gamma_0, \gamma_1)$ to be the automorphism on T obtained by applying γ_0 to T_0 and γ_1 to T_1 . Define automorphisms b, c, d of T recursively as follows:

$$(5.1) \quad b = (a, c), \quad c = (a, d), \quad d = (e, b),$$

where e is the identity automorphism. The Grigorchuk group is defined as the subgroup of $\text{Aut}(T)$ generated by the set $\{a, b, c\}$.

Theorem 5.1. *The Cayley graph $G = (V, E)$ of the Grigorchuk group with generator set $\{a, b, c\}$ satisfies:*

- (a) G admits no graph height function,
- (b) for $\mathcal{H} \trianglelefteq \text{Aut}(G)$ with finite index, any \mathcal{H} -difference-invariant function on V is constant on each orbit of \mathcal{H} .

The proof of Theorem 5.1 is given in Section 10. In the preceding Section 9, two approaches are developed for showing the absence of a graph height function within particular classes of Cayley graph. In the case of the Grigorchuk group, two reasons combine to forbid graph height functions, namely, its Cayley group has no automorphisms beyond the action of the group itself, and the group is a torsion group in that every element has finite order.

Since the Grigorchuk group is amenable, Theorems 4.1 and 5.1 yield that: within the class of infinite, finitely generated groups, every elementary amenable group has a graph height function, but there exists an amenable group without a graph height function. The Grigorchuk group is finitely *generated* but not finitely *presented*, [12, Thm 6.2].

We ask if there exists an infinite, finitely *presented*, amenable group with a Cayley graph having no graph height function. A natural candidate might be the group

$\Gamma = \langle S \mid R \rangle$ of [13, Thm 1], with

$$S = \{a, c, d, t\},$$

$$R = \{a^2 = c^2 = d^2 = (ad)^4 = (adacac)^4 = \mathbf{1}, t^{-1}at = aca, t^{-1}ct = dc, t^{-1}dt = c\}.$$

This finitely presented, amenable HNN-extension of the Grigorchuk group is not elementary amenable. However, since it contains the free group generated by the stable letter t , it possesses a group height function. More precisely, the function

$$h(\mathbf{1}) = 0, \quad h(t) = 1, \quad h(t^{-1}) = -1, \quad h(s) = 0 \text{ for } s \in S, \quad s \neq t^{\pm 1},$$

defines a group height function.

Part C. Results for non-amenable graphs

6. CONNECTIVE CONSTANTS OF NON-AMENABLE GRAPHS

Let $G \in \mathcal{G}$ have degree Δ . It was proved in [20, Thm 4.1] that

$$\sqrt{\Delta - 1} \leq \mu(G) \leq \Delta - 1.$$

The upper bound is achieved by the Δ -regular tree T_Δ . It is unknown if the lower bound is sharp for *simple* graphs. This lower bound may however be improved for non-amenable graphs, as follows.

Let P be the transition matrix of simple random walk (SRW) on $G = (V, E)$, and let I be the identity matrix. The *spectral bottom* of $I - P$ is defined to be the largest λ with the property that, for all $f \in l^2(V)$,

$$(6.1) \quad \langle f, (I - P)f \rangle \geq \lambda \langle f, f \rangle.$$

It may be seen that $\lambda(G) = 1 - \rho(G)$ where $\rho(G)$ is the spectral radius of P (see [35, Sect. 6], and [42] for an account of the spectral radius).

Remark 6.1. *It is known that G is a non-amenable if and only if $\rho(G) < 1$, which is equivalent to $\lambda(G) > 0$. This was proved by Kesten [32, 33] for Cayley graphs of finitely-presented groups, and extended to general transitive graphs by Dodziuk [8] (see also the references in [35, Sect. 6.10]).*

Theorem 6.2. *Let $G \in \mathcal{G}$ have degree $\Delta \geq 3$. Let P be the transition matrix of SRW on G , and λ the spectral bottom of P . The connective constant $\mu(G)$ satisfies*

$$(6.2) \quad \mu(G) \geq (\Delta - 1)^{\frac{1}{2}(1+c\lambda)},$$

where $c = \Delta(\Delta - 1)/(\Delta - 2)^2$.

The improvement in the lower bound for $\mu(G)$ is strict if and only if $\lambda > 0$, which is to say that G is non-amenable. It is standard (see [35, Thm 6.7]) that

$$(6.3) \quad \frac{1}{2}\phi^2 \leq 1 - \sqrt{1 - \phi^2} \leq \lambda \leq \phi,$$

where $\phi = \phi(G)$ is the edge-isoperimetric constant of (2.1). By [1, Thm 3],

$$(6.4) \quad \lambda(G) \leq \lambda(T_\Delta) - \frac{\Delta - 2}{\Delta(\Delta - 1)^{g+2}},$$

where g is the girth of G , T_Δ is the Δ -regular tree, and

$$(6.5) \quad \lambda(T_\Delta) = 1 - \frac{2\sqrt{\Delta - 1}}{\Delta}.$$

Remark 6.3. *The spectral bottom (and therefore the spectral radius, also) is not a continuous function of G in the usual graph metric (see [18, Sect. 5]). This follows from [41, Thm 2.4], where it is proved that, for all pairs (k, l) with $k \geq 2$ and $l \geq 3$, there exists a group with polynomial growth whose Cayley graph $G_{k,l}$ is $2k$ -regular with girth exceeding l . Since $G_{k,l}$ is amenable, we have $\lambda(G_{k,l}) = 0$, whereas $\lambda(T_{2k})$ is given by (6.5).*

Proof of Theorem 6.2. This is achieved by a refinement of the argument used to prove [20, Thm 4.1], and we shall make use of the notation introduced in that proof.

Let $v_0 = \mathbf{1}$, and let $\pi = (v_0, v_1, \dots, v_{2n})$ be an extendable $2n$ -step SAW of G . For convenience, we augment π with a mid-edge incident to v_0 and not lying on the edge $\langle v_0, v_1 \rangle$. Let E_π be the set of oriented edges $[v, w]$ such that: (i) $v \in \pi$, $v \neq v_{2n}$, and (ii) the (non-oriented) edge $\langle v, w \rangle$ does not lie in π . Note that

$$(6.6) \quad |E_\pi| = 2n(\Delta - 2).$$

Each (oriented) edge in E_π is coloured either red or blue according to the following rule. For $v \in \pi$, let π_v be the sub-path of π joining v_0 to v . The edge $[v, w] \in E_\pi$ is coloured red if $\pi_v \cup [v, w]$ is not an extendable SAW, and is coloured blue otherwise. By (6.6), the number B_π (respectively, R_π) of blue edges (respectively, red edges) satisfies

$$(6.7) \quad B_\pi + R_\pi = 2n(\Delta - 2).$$

We shall make use of the following lemma.

Lemma 6.4. *The number B_π satisfies*

$$(6.8) \quad B_\pi \geq \frac{n(1 + c\lambda)}{\Delta - 2} - \frac{\Delta - 1}{2},$$

where $c = \Delta(\Delta - 1)/(\Delta - 2)^2$.

We now argue as in [20, Lemma 5.1] to deduce from Lemma 6.4 that the number of extendable $2n$ -step SAWs from v_0 is at least $C(\Delta - 1)^{n(1+c\lambda)}$ where $C = C(\Delta)$. Inequality (6.2) follows as required. \square

Proof of Lemma 6.4. An edge $[v, w] \in E_\pi$ is said to be *finite* if w lies in a finite component of $G \setminus \pi$, and *infinite* otherwise. If $[v, w] \in E_\pi$ is red, then w is necessarily finite. Blue edges, on the other hand, may be either finite or infinite.

It was explained in the proof of [20, Thm 4.1] that there exists an injection f from the set of red edges to the set of blue edges with the property that, if $e = [v, w]$ is red, and $f(e) = [v', w']$, then w and w' lie in the same component of $G \setminus \pi_v$. Since e is finite, so is $f(e)$. It follows that

$$(6.9) \quad B_\pi \geq R_\pi + B_\pi^\infty,$$

where R_π is the number of red edges, and B_π^∞ is the number of infinite blue edges.

Let $X = (X_m : m = 0, 1, 2, \dots)$ be a SRW on G , and let \mathbb{P}_v denote the law of X started at $v \in V$. For $[v, w] \in E_\pi$, let

$$\beta_{[v,w]} = \mathbb{P}_v(X_1 = w, \text{ and } \forall m > 0, X_m \notin \pi).$$

By [5, Lemma 2.1] with $A = \pi$,

$$(6.10) \quad \lambda \leq \frac{1}{2n+1} \left(\sum_{[v,w] \in E_\pi} \beta_{[v,w]} + \sum_w \beta_{[v_{2n}, w]} \right).$$

If $[v, w] \in E_\pi$ is finite, then $\beta_{[v,w]} = 0$. By (6.10),

$$(6.11) \quad \lambda \leq \left(\frac{B_\pi^\infty + \Delta - 1}{2n} \right) \mathbb{P}_v(X_1 = w, \text{ and } \forall m > 0, X_m \neq v).$$

The last probability depends on the graph G , and it is a maximum when G is the Δ -regular tree T_Δ (since T_Δ is the universal cover of G). Therefore, it is no greater than $1/\Delta$ multiplied by the probability that a random walk on \mathbb{Z} , which moves rightwards with probability $p = (\Delta - 1)/\Delta$ and leftwards with probability $q = 1 - p$, never visits 0 having started at 1. By, for example, [22, Example 12.59],

$$\mathbb{P}_v(X_1 = w, \text{ and } \forall m > 0, X_m \neq v) \leq \frac{1}{\Delta} \left(1 - \frac{q}{p} \right) = \frac{\Delta - 2}{\Delta(\Delta - 1)}.$$

By (6.9) and (6.11),

$$(6.12) \quad B_\pi \geq R_\pi + 2n\lambda \frac{\Delta(\Delta - 1)}{\Delta - 2} - \Delta + 1,$$

and (6.8) follows by (6.7). □

7. GRAPHS WITH LARGE GIRTH

Benjamini, Nachmias, and Peres showed in [5, Thm 1.1] that the critical probability $p_c(G)$ of bond percolation on a Δ -regular, non-amenable graph G with large girth is close to that of the critical probability of the Δ -regular tree T_Δ . Their main result implies the following.

Theorem 7.1. *Let $G \in \mathcal{G}$ be non-amenable with degree $\Delta \geq 3$ and girth $g \leq \infty$. There exists an absolute positive constant C such that*

$$(7.1) \quad \left[\frac{1}{\Delta - 1} + C \frac{\log(1 + \lambda^{-2})}{g\Delta} \right]^{-1} \leq \mu(G) \leq \Delta - 1,$$

where $\lambda = \lambda(G)$ is the spectral bottom of SRW on G , as in (6.1). Equality holds in the upper bound of (7.1) if and only if G is a tree, that is, $g = \infty$.

Proof. The upper bound of (7.1) is from [20, Thm 4.2]. The lower bound is an immediate consequence of [5, Thm 1.1] and the fact that $\mu(G) \geq 1/p_c(G)$ (see, for example, [6, Thm 7] and [15, eqn (1.13)], which hold for general quasi-transitive graphs). \square

We recall from Remark 6.1 that $\lambda > 0$ if and only if G is non-amenable. Theorem 7.1 does not, of itself, imply that $\mu(\cdot)$ is continuous at T_Δ , since $\lambda(\cdot)$ is not continuous at T_Δ (in the case when Δ is even, see Remark 6.3). For continuity at T_Δ , it would suffice that $\lambda(\cdot)$ is bounded away from 0 on a neighbourhood of T_Δ . By (6.3), this is valid within any class of graphs whose edge-isoperimetric constants (2.1) are bounded uniformly from 0. See also [18, Thm 5.1].

8. THE HIGMAN GROUP

The Higman group Γ of [28] is the infinite, finitely presented group with presentation $\Gamma = \langle S \mid R \rangle$ where

$$(8.1) \quad \begin{aligned} S &= \{a, b, c, d, a^{-1}, b^{-1}, c^{-1}, d^{-1}\}, \\ R &= \{a^{-1}ba = b^2, b^{-1}cb = c^2, c^{-1}dc = d^2, d^{-1}ad = a^2\}. \end{aligned}$$

This group is interesting since it has no proper normal subgroup with finite index, and the quotient of Γ by its maximal proper normal subgroup is an infinite, finitely generated, simple group. By [21, Thm 4.1(b)], Γ has no group height function. The above two reasons conspire to forbid graph height functions.

Theorem 8.1. *The Cayley graph $G = (V, E)$ of the Higman group $\Gamma = \langle S \mid R \rangle$ has no graph height function.*

A further group of Higman type is given as follows. Let S be as above, and let $\Gamma' = \langle S \mid R' \rangle$ be the finitely presented group with

$$R' = \{a^{-1}ba = b^2, b^{-2}cb^2 = c^2, c^{-3}dc^3 = d^2, d^{-4}ad^4 = a^2\}.$$

Note that Γ' is infinite and non-amenable, since the subgroup generated by the set $\{a, c, a^{-1}, c^{-1}\}$ is a free group (as in the corresponding step for the Higman group at [28, pp. 62–63]).

Theorem 8.2. *The Cayley graph $G = (V, E)$ of the above group $\Gamma' = \langle S \mid R' \rangle$ has no graph height function.*

The proofs of the above theorems are given in Sections 11 and 12, respectively.

Part D. Remaining proofs

9. CRITERIA FOR THE ABSENCE OF HEIGHT FUNCTIONS

This section contains some observations relevant to proofs in Sections 10–12 of the non-existence of graph height functions.

Let $\Gamma = \langle S \mid R \rangle$ where $|S| < \infty$, and let $G = (V, E)$ be the corresponding Cayley graph. Let Π be the set of permutations of S that preserve Γ up to isomorphism, and write $e \in \Pi$ for the identity. Thus, $\pi \in \Pi$ acts on Γ by: for $w = s_1s_2 \cdots s_m$ with $s_i \in S$, we have $\pi(w) = \pi(s_1)\pi(s_2) \cdots \pi(s_m)$. It follows that $\Pi \subseteq \text{Aut}(G)$. For $\gamma = g_1g_2 \cdots g_n \in \Gamma$ with $g_i \in S$, and $\pi \in \Pi$, we define $\gamma\pi \in \text{Aut}(G)$ by $\gamma\pi(w) = g_1g_2 \cdots g_n\pi(w)$, $w \in V$. Write $\Gamma\Pi \subseteq \text{Aut}(G)$ for the subgroup containing all such $\gamma\pi$, and note that γe operates on G in the manner of γ with left-multiplication.

The *stabilizer* Stab_v of $v \in V$ is the set of automorphisms of G that preserve v , that is,

$$\text{Stab}_v = \{\eta \in \text{Aut}(G) : \eta(v) = v\}.$$

Proposition 9.1. *Suppose $\text{Stab}_1 = \Pi$.*

- (a) $\text{Aut}(G) = \Gamma\Pi$.
- (b) *If $\mathcal{M} \trianglelefteq \text{Aut}(G)$ has finite index, the subgroup $\mathcal{N} = \mathcal{M} \cap \Gamma$ satisfies $\mathcal{N} \trianglelefteq \Gamma$ and $[\Gamma : \mathcal{N}] < \infty$.*
- (c) *If G has a graph height function, then it has a strong graph height function.*

Proof. Assume $\text{Stab}_1 = \Pi$.

(a) Let $\eta \in \text{Aut}(G)$, and write $\gamma = \eta(\mathbf{1})$. Then $\gamma^{-1}\eta \in \text{Stab}_1$, which is to say that $\gamma^{-1}\eta = \pi \in \Pi$, and thus $\eta = \gamma\pi \in \Gamma\Pi$ so that $\text{Aut}(G) = \Gamma\Pi$. Note for future use that

$$[\text{Aut}(G) : \Gamma] = |\Pi| < \infty.$$

(b) Let $\mathcal{M} \trianglelefteq \text{Aut}(G)$ be a finite-index normal subgroup, and let $\mathcal{N} = \{\gamma e : \gamma e \in \mathcal{M}\}$. Viewed as automorphisms, we have that $\gamma e = \gamma$, and hence $\mathcal{N} \leq \Gamma \leq \text{Aut}(G)$.

For $\alpha \in \Gamma$, $\nu \in \mathcal{N}$, we have that $(\alpha^{-1}\nu\alpha)e = \alpha^{-1}(\nu e)\alpha \in \mathcal{M}$, since $\mathcal{M} \trianglelefteq \text{Aut}(G)$. Therefore, $\mathcal{N} \trianglelefteq \Gamma$.

Since $\Gamma, \mathcal{M} \leq \text{Aut}(G)$ and $\mathcal{N} = \Gamma \cap \mathcal{M}$, we have that

$$[\text{Aut}(G) : \mathcal{N}] \leq [\text{Aut}(G) : \Gamma] \cdot [\text{Aut}(G) : \mathcal{M}] < \infty,$$

which implies $[\Gamma : \mathcal{N}] < \infty$, as required.

(c) Let (h, \mathcal{H}) be a graph height function of G . Since \mathcal{H} is a finite-index normal subgroup of $\text{Aut}(G)$, by part (b), there exists $\mathcal{N} \leq \mathcal{H}$ that is a finite-index normal subgroup of Γ . Since $\mathcal{N} \leq \mathcal{H}$, h acts on Γ and is \mathcal{N} -difference invariant, whence (h, \mathcal{N}) is a strong graph height function. \square

Corollary 9.2. *Let $\Gamma = \langle S \mid R \rangle$ have Cayley graph G satisfying $\text{Stab}_1 = \Pi$.*

- (a) *If Γ has no proper, normal subgroup with finite index, any graph height function of G is also a group height function of Γ .*
- (b) *If Γ is a torsion group, then G has no graph height function.*

We recall from [18, Example 4.4] that the infinite dihedral group is a torsion group, whereas its Cayley graph has a graph height function.

Proof. (a) Let (h, \mathcal{M}) be a graph height function of G . If Γ satisfies the given condition then, by Proposition 9.1(b), $\mathcal{M} \supseteq \Gamma$. Therefore, (h, Γ) is a graph height function and hence a group height function.

(b) If G has a graph height function, by Proposition 9.1(c), G has a strong graph height function (h, \mathcal{N}) . Assume every element of Γ has finite order. For $\gamma \in \mathcal{N}$ with $\gamma^n = \mathbf{1}$, we have that $h(\gamma^n) = nh(\gamma) = 0$, whence $h \equiv 0$ on \mathcal{N} .

For $\gamma \in \Gamma$, find α_γ such that $\gamma \in \alpha_\gamma \mathcal{N}$. Since h is \mathcal{N} -difference-invariant, there exists $\nu \in \mathcal{N}$ such that

$$(9.1) \quad h(\gamma) = h(\alpha_\gamma) + h(\nu) = h(\alpha_\gamma).$$

Now $[\Gamma : \mathcal{N}] < \infty$, so we may consider only finitely many choices for α_γ . Therefore, h is bounded on Γ , in contradiction of the assumption that it is a graph height function. \square

10. PROOF OF THEOREM 5.1

The main step is to show that

$$(10.1) \quad \text{Stab}_1 = \{e\},$$

where e is the identity of $\text{Aut}(G)$. Once this is shown, claim (a) follows from Corollary 9.2(b) and the fact that every element of the Grigorchuk group has finite order, [26]. It therefore suffices for (a) to show (10.1), and to this end we study the structure of the Cayley graph $G = (V, E)$.

It was shown in [36] (see also [14, eqn (4.7)]) that $\Gamma = \langle S \mid R \rangle$ where $S = \{a, b, c, d\}$, R is the following set of relations

$$(10.2) \quad \begin{aligned} \mathbf{1} &= a^2 = b^2 = c^2 = d^2 = bcd \\ &= \sigma^k((ad)^4) = \sigma^k((adacac)^4), \quad k = 0, 1, 2, \dots, \end{aligned}$$

and σ is the substitution

$$\sigma : \begin{cases} a \mapsto aca, \\ b \mapsto d, \\ c \mapsto b, \\ d \mapsto c. \end{cases}$$

It follows that the following, written in terms of the reduced generator set $\{a, b, c\}$ after elimination of d , are valid relations:

$$(10.3) \quad \mathbf{1} = a^2 = b^2 = c^2 = (bc)^2 = (abc)^4 = (ac)^8 = (abcacac)^4 = (acab)^8 = (ab)^{16},$$

(see also [14, Sect. 1]). Note the asymmetry between b and c in that ab (respectively, ac) has order 16 (respectively, 8).

Let

$$V_n = \{v \in \Gamma : \text{dist}(v, \mathbf{1}) = n\},$$

where dist denotes graph-distance on G . Since G is locally finite, $|V_n| < \infty$. For $\eta \in \text{Stab}_{\mathbf{1}}$, η restricted to V_n is a permutation of V_n . As illustrated in Figure 10.1,

$$V_0 = \{\mathbf{1}\}, \quad V_1 = \{a, b, c\}, \quad V_2 = \{ab, ac, ba, bc = cb, ca\}.$$




FIGURE 10.1. The subgraph of G on $V_0 \cup V_1 \cup V_2$.

Let $\eta \in \text{Stab}_{\mathbf{1}}$, so that $\eta(a) \in V_1$. Since the shortest cycles using the edges $\langle \mathbf{1}, b \rangle$ and $\langle \mathbf{1}, c \rangle$ have length 4, and using $\langle \mathbf{1}, a \rangle$ greater than 4 (see Figure 10.1), we have that $\eta(a) = a$. By a similar argument, we obtain that, for $n \geq 1$,

$$(10.4) \quad \eta(va) = \eta(v)a, \quad v \in V_n, \quad va \in V_{n+1},$$

which we express by saying that η maps a -type edges to a -type edges.

We show next that

$$(10.5) \quad \eta(vc) = \eta(v)c, \quad v \in V, \eta \in \text{Stab}_{\mathbf{1}},$$

which is to say that η maps c -type edges to c -type edges. By (10.4)–(10.5), $\eta \in \text{Stab}_{\mathbf{1}}$ maps b -type edges to b -type edges also, whence $\eta = e$ as required. It remains to prove (10.5).

Assume, in contradiction of (10.5), that there exists $v \in V$, $\eta \in \text{Stab}_{\mathbf{1}}$ such that $\eta(vc) = \eta(v)b$. Since ac has order 8, we have that $(ca)^8 = \mathbf{1}$. Let C be the directed cycle corresponding to the word $v(ca)^8$; thus, C includes the edge $[v, vc]$. Then $\eta(C)$ is a cycle of length 16 including the edge $[\eta(v), \eta(v)b]$. Since C contains exactly 8 a -type edges at alternating positions, by (10.4), so does $\eta(C)$. Therefore, $\eta(C)$ has the form $\eta(v)ba \prod_{i=2}^8 (x_i a)$, where $x_i \in \{b, c\}$ for $i = 2, 3, \dots, 8$. In particular,

$$(10.6) \quad ba \prod_{i=2}^8 (x_i a) = \mathbf{1}, \quad x_i \in \{b, c\}, \quad i = 2, 3, \dots, 8.$$

The word problem of the Grigorchuk group is solvable (see [12] and [14, Sect. 4]), in that there exists an algorithm to determine whether or not $w = \mathbf{1}$ for any given word $w \in \{a, b, c\}^*$ (where S^* denotes the set of finite ordered sequences of elements of S). By applying this algorithm (see below), we deduce that (10.6) has no solution. Equation (10.5) follows, and the proof of part (a) is complete.

Finally, here is a short amplification of the analysis of (10.6). The word in (10.6) has the form $b(ay_1a)z_1(ay_2a)z_2(ay_3a)z_3(ay_4a)$, where $y_i, z_j \in \{b, c\}$. By (5.1), the effect of such a word on the right sub-tree T_1 is $\gamma_1 := ca(c/d)a(c/d)a(c/d)a$, where each term of the form (y/z) is to be interpreted as ‘either y or z ’. The effect of γ_1 on the left sub-tree T_{10} of T_1 is $\gamma_{10} := a(d/b)(a/e)(d/b)$. If there is an odd number of appearances of a in γ_{10} , then γ_{10} is not the identity, and thus we may assume $\gamma_{10} := a(d/b)a(d/b)$. It is immediate that none of the four possibilities is the identity, and the claim follows.

Part (b) holds as follows. Suppose there exists $\mathcal{H} \trianglelefteq \Gamma$, $\gamma \in \Gamma$, and a non-constant \mathcal{H} -difference-invariant function $F : \gamma\mathcal{H} \rightarrow \mathbb{Z}$. It is elementary that \mathcal{H} is unimodular and symmetric (see, for example, [19, Sect. 4]). By [21, Thm 3.5] and the comment near the beginning of [21, Sect. 8], G has a graph height function, in contradiction of part (a).

11. PROOF OF THEOREM 8.1

We shall prove three statements:

- (i) Γ has no group height function,

- (ii) Π is the cyclic group generated by the permutation $(abcd)$, with the convention that $\eta(x^{-1}) = \eta(x)^{-1}$, for $\eta \in \Pi$, $x \in \{a, b, c, d\}$,
- (iii) $\text{Stab}_{\mathbf{1}} = \Pi$.

It is proved in [28] that the Higman group has no proper, finite-index, normal subgroup, and the result follows from the above statements by Corollary 9.2(a).

Proof of (i). The absence of a group height function is immediate by [21, Example 6.3].

Proof of (ii). Evidently, Π contains the given cyclic group, and we turn to the converse. Since elements of Π preserve Γ up to isomorphism,

$$(11.1) \quad \eta(x^{-1}) = \eta(x)^{-1}, \quad x \in S.$$

We next rule out the possibility that $\eta(x) = y^{-1}$ for some $x, y \in \{a, b, c, d\}$. Suppose, for illustration, that $\eta(a) = b^{-1}$. By (11.1), the relation $a^{-1}ba = b^2$ becomes $b\beta b^{-1} = \beta^2$ where $\beta = \eta(b)$. The Higman group has no such relation with $\beta \in S$. In summary,

$$(11.2) \quad \eta(x) \in \{a, b, c, d\}, \quad \eta(x^{-1}) = \eta(x)^{-1}, \quad x \in \{a, b, c, d\}.$$

The shortest cycles containing the edge $\langle \mathbf{1}, a \rangle$, modulo rotation and reversal, arise from the relations $ab^2a^{-1}b^{-1} = \mathbf{1}$ and $ada^{-2}d^{-1} = \mathbf{1}$ (see Figure 11.2). The first uses $a^{\pm 1}$ twice and $b^{\pm 1}$ thrice, and the second uses $a^{\pm 1}$ thrice and $d^{\pm 1}$ twice. Let $\eta \in \Pi$, and suppose for illustration that $\eta(a) = b$ (the same argument is valid for any $\eta(x)$, $x \in \{a, b, c, d\}$). By considering the cycles starting $\langle \mathbf{1}, b \rangle$, $\langle \mathbf{1}, c \rangle$, $\langle \mathbf{1}, d \rangle$, and using (11.2), we deduce that

$$\eta(b) = c, \quad \eta(c) = d, \quad \eta(d) = a,$$

and the claim is proved.

Proof of (iii). We begin with some observations concerning the Baumslag–Solitar (BS) group $\text{BS}(x, y)$ with presentation $\langle x, y, x^{-1}, y^{-1} \mid x^{-1}yx = y^2 \rangle$, of which the Cayley graph is sketched in Figure 11.1. Edges of the form $\langle \gamma, \gamma x^{\pm 1} \rangle$ have *type* x , and of the form $\langle \gamma, \gamma y^{\pm 1} \rangle$ *type* y . By inspection, the shortest cycles have length 5 (see Figure 11.2), and, for $\gamma \in \text{BS}(x, y)$,

$$(11.3) \quad \text{for } p, q = \pm 1, \text{ the edges } \langle \gamma, \gamma x^p \rangle \text{ and } \langle \gamma, \gamma y^q \rangle \text{ lie in a common 5-cycle,}$$

$$(11.4) \quad \text{the third edge of any directed 5-cycle beginning } [\gamma, \gamma x] \text{ has type } y,$$

$$(11.5) \quad \text{the third edge of any directed 5-cycle beginning } [\gamma, \gamma x^{-1}] \text{ has type } x,$$

$$(11.6) \quad \text{every 5-cycle contains two consecutive edges of type } y, \text{ and not of type } x,$$

$$(11.7) \quad \text{a type } x \text{ (respectively, type } y) \text{ edge lies in 2 (respectively, 3) 5-cycles.}$$

Returning to the Higman group, for convenience, we relabel the vector (a, b, c, d) as (s_0, s_1, s_2, s_3) , with addition and subtraction of indices modulo 4. Let G be the Cayley graph of the Higman group $\Gamma = \langle S \mid R \rangle$, rooted at $\mathbf{1}$. An edge of G is said




FIGURE 11.1. Part of the Cayley graph of the Baumslag–Solitar group $BS(x, y)$.




FIGURE 11.2. Part of one ‘sheet’ of the Cayley graph of $BS(x, y)$.

to be of *type* s_i if it has the form $\langle \gamma, \gamma s_i^{\pm 1} \rangle$ with $\gamma \in \Gamma$. We explain next how to obtain information about the types of the edges of G , by examination of G only, and without further information about the vertex-labellings as elements of Γ .

We consider first the set $\partial_e \mathbf{1}$ of edges of G incident to $\mathbf{1}$. Let $e_1 = \langle \mathbf{1}, v \rangle$, $t \in \{0, 1, 2, 3\}$, and $p \in \{-1, 1\}$. Assume that

$$(11.8) \quad v = s_t^p,$$

so that, in particular, e_1 has type s_t . By (11.3), for $j = \pm 1$, e_1 lies in a 5-cycle of $BS(s_{t-1}, s_t)$ (respectively, $BS(s_t, s_{t+1})$) containing $\langle \mathbf{1}, s_{t-1}^j \rangle$ (respectively, $\langle \mathbf{1}, s_{t+1}^j \rangle$). On the other hand, by consideration of the relator set R , e_1 lies in no 5-cycle including an edge of type s_{t+2} . Therefore, the edges of the form $\langle \mathbf{1}, s_{t+2}^{\pm 1} \rangle$ may be identified by

examination of G , and we denote these as g_1, g_2 . There is exactly one further edge of $\partial_e \mathbf{1}$ that lies in no 5-cycle containing either g_1 or g_2 , and we denote this edge as e_2 . In summary,

$$\{e_1, e_2\} = \{\langle \mathbf{1}, s_t^{-1} \rangle, \langle \mathbf{1}, s_t \rangle\}, \quad \{g_1, g_2\} = \{\langle \mathbf{1}, s_{t+2}^{-1} \rangle, \langle \mathbf{1}, s_{t+2} \rangle\}.$$

Having identified the edges of $\partial_e \mathbf{1}$ with types s_t and s_{t+2} , we move to the other endpoint $v = s_t^p$ of e_1 , and apply the same argument. Let e_1, e_1' be the two type- s_t edges incident to v .

We turn next to the remaining four edges of $\partial_e \mathbf{1}$. Let k be such an edge, and consider the property: k lies in a 5-cycle of G containing both e_1 and e_1' . By (11.6) and examination of the Cayley graphs of the four groups $\text{BS}(s_i, s_{i+1})$, $0 \leq i < 4$, we see that k has this property if it has type $t-1$, and not if it has type $t+1$. Thus we may identify the types of the four remaining edges of $\partial_e \mathbf{1}$, which we write as

$$\{f_1, f_2\} = \{\langle \mathbf{1}, s_{t+1}^{-1} \rangle, \langle \mathbf{1}, s_{t+1} \rangle\}, \quad \{h_1, h_2\} = \{\langle \mathbf{1}, s_{t+3}^{-1} \rangle, \langle \mathbf{1}, s_{t+3} \rangle\}.$$

Having determined the types of edges in $\partial_e \mathbf{1}$ (relative to the type t of the initial edge e_1), we move to an endpoint of such an edge other than $\mathbf{1}$, and apply the same argument. By iteration, we deduce the types of all edges of G . Let $T(k)$ denote the type of edge k . It follows from the above that

$$(11.9) \quad T(k) - T(e_1) \text{ is independent of } t = T(e_1),$$

with arithmetic on indices, modulo 4.

We explain next how to identify the value of $p = p(v)$ in (11.8) from the graphical structure of G . Let S_i be the subgraph of G containing all edges with type either s_i or s_{i+1} , so that each component of S_i is isomorphic to the Cayley graph of $\text{BS}(s_i, s_{i+1})$. By (11.4)–(11.5), every directed 5-cycle of $\text{BS}(s_t, s_{t+1})$ starting with the edge $[\mathbf{1}, s_t]$ has third edge with type s_{t+1} , whereas every directed 5-cycle starting with $[\mathbf{1}, s_t^{-1}]$ has third edge with type s_t . We examine S_t to determine which of these two cases holds, and the outcome determines the value of $p = p(v)$.

The above argument is applied to each directed edge $[\gamma, \gamma s_i^{\pm 1}]$ of G , and the power of s_i is thus determined from the graphical structure of G .

Let $\eta \in \text{Stab}_{\mathbf{1}}$. By (11.9), the effect of η is to change the edge-types by

$$T(k) \mapsto T(k) + T(\eta(e_1)) - t.$$

Now, $\eta(v)$ is adjacent to $\mathbf{1}$ and, by the above, once $\eta(v)$ is known, the action of η on the rest of G is determined. Since $\eta \in \text{Aut}(G)$, $\eta(v)$ may be any neighbour w of $\mathbf{1}$ with the property that $p(w) = p(v)$. There are exactly four such neighbours (including v) and we deduce from (11.9) that η lies in the cyclic group generated by the permutation $(s_0 s_1 s_2 s_3)$.

12. PROOF OF THEOREM 8.2

We shall prove three statements:

- (i) Γ has no group height function,
- (ii) $\text{Stab}_1 = \Pi$ where $\Pi = \{e\}$,
- (iii) Γ has no proper normal subgroup with finite index.

The result follows from these statements by Corollary 9.2(a), and we turn to their proofs.

Proof of (i). The absence of a group height function is immediate by [21, Thm 4.1(b)].

Proof of (ii). Let $\eta \in \text{Stab}_1$ and $\gamma \in \Gamma$. We consider the action of η on directed edges of G . By inspection of the set R' of relations, an edge of the type $\langle \gamma, \gamma x \rangle$ lies in shortest cycles of length

$$\begin{cases} 5, 8 & \text{if } x = a^{\pm 1}, \\ 5, 7 & \text{if } x = b^{\pm 1}, \\ 7, 8 & \text{if } x = c^{\pm 1}, \\ 9, 11 & \text{if } x = d^{\pm 1}. \end{cases}$$

Since the four combinations are distinct, it must be that

$$(12.1) \quad \eta([\gamma, \gamma x]) = [\gamma', \gamma' x^{\pm 1}], \quad \gamma \in \Gamma, \quad x \in S,$$

where $\gamma' = \eta(\gamma)$. We show next that

$$(12.2) \quad \eta([\gamma, \gamma x]) \neq [\gamma', \gamma' x^{-1}], \quad \gamma \in \Gamma, \quad x \in S,$$

which combines with (12.1) to imply $\eta = e$ as required.

It suffices to consider the case $x = a$ in (12.2), since a similar proof holds in the other cases. Suppose $\eta([\gamma, \gamma a]) = [\gamma', \gamma' a^{-1}]$, and consider the cycle corresponding to $\gamma a b^{-2} a^{-1} b^{-1}$, that is $(\gamma, \gamma a, \gamma a b^{-1}, \gamma a b^{-2}, \gamma a b^{-2} a^{-1}, \gamma a b^{-2} a^{-1} b^{-1} = \gamma)$. By (12.1), this is mapped under η to the cycle corresponding to $\gamma' a^{-1} b^{\pm 2} a^{\pm 1} b^{\pm 1}$. By examining the relation set R' , the only cycles beginning $\gamma' a^{-1} b^{\pm 1}$ with length not exceeding 5 are $\gamma' a^{-1} b a b^{-2}$ and $\gamma' a^{-1} b^{-1} a b^2$, in contradiction of the above (since the third step of these two cycles is a rather than the required $b^{\pm 1}$).

Proof of (iii). Suppose \mathcal{N} is a proper normal subgroup of Γ with finite index. The quotient group Γ/\mathcal{N} is non-trivial and finite with generators $\bar{s} = s\mathcal{N}$, $s \in S$, satisfying

$$(12.3) \quad \begin{aligned} \bar{a}^{-1} \bar{b} \bar{a} &= \bar{b}^2, & \bar{b}^{-2} \bar{c} \bar{b}^2 &= \bar{c}^2, \\ \bar{c}^{-3} \bar{d} \bar{c}^3 &= \bar{d}^2, & \bar{d}^{-4} \bar{a} \bar{d}^4 &= \bar{a}^2. \end{aligned}$$

Since Γ/\mathcal{N} is finite, each \bar{s} has finite order, denoted $\text{ord}(\bar{s})$. It follows from (12.3) that

$$(12.4) \quad \text{ord}(\bar{s}) > 1, \quad s = a, b, c, d.$$

To see this, suppose for illustration that $\text{ord}(\bar{c}) = 1$, so that $\bar{c} = \bar{\mathbf{1}}$. By the third equation of (12.3), $\text{ord}(\bar{d}) = 1$, so that $\bar{d} = \bar{\mathbf{1}}$, and similarly for \bar{a} and \bar{b} , implying that Γ/\mathcal{N} is trivial, a contradiction.

By induction, for $n \geq 1$,

$$\begin{aligned} \bar{a}^{-n}\bar{b}\bar{a}^n &= \bar{b}^{2^n}, & \bar{b}^{-2n}\bar{c}\bar{b}^{2n} &= \bar{c}^{2^n}, \\ \bar{c}^{-3n}\bar{d}\bar{c}^{3n} &= \bar{d}^{2^n}, & \bar{d}^{-4n}\bar{a}\bar{d}^{4n} &= \bar{a}^{2^n}, \end{aligned}$$

whence, by setting $n = \text{ord}(\bar{a})$, etc,

$$(12.5) \quad \begin{aligned} \text{ord}(\bar{b}) &| (2^{\text{ord}(\bar{a})} - 1), & \text{ord}(\bar{c}) &| (2^{\text{ord}(\bar{b})} - 1), \\ \text{ord}(\bar{d}) &| (2^{\text{ord}(\bar{c})} - 1), & \text{ord}(\bar{a}) &| (2^{\text{ord}(\bar{d})} - 1), \end{aligned}$$

where $u | v$ means that v is a multiple of u . We shall deduce a contradiction from (12.4) and (12.5). This is done as in [28], of which we reproduce the proof for completeness.

Let p be the least prime factor of the four integers $\text{ord}(\bar{s})$, $s \in \{a, b, c, d\}$. By (12.4), $p > 1$. Suppose that $p | \text{ord}(\bar{a})$ (with a similar argument if $p | \text{ord}(\bar{s})$ for some other parameter s). Then $p | 2^{\text{ord}(\bar{d})} - 1$ by (12.5), and in particular p is odd and therefore coprime with 2. Let r be the multiplicative order of 2 mod p , that is, the least positive integer r such that $p | 2^r - 1$. In particular, $r > 1$, so that r has a prime factor q . By Fermat's little theorem, $r | p - 1$ so that $q < p$. Furthermore, $r | \text{ord}(\bar{d})$ so that $q | \text{ord}(\bar{d})$, in contradiction of the minimality of p . We deduce that Γ has no proper, normal subgroup with finite index.

ACKNOWLEDGEMENTS

GRG was supported in part by the Engineering and Physical Sciences Research Council under grant EP/I03372X/1. ZL acknowledges support from the Simons Foundation under grant #351813. The authors thank a reader for proposing indicability as a route to the rationalization of an earlier proof. GRG thanks Jack Button for valuable advice.

REFERENCES

- [1] M. Abért, Y. Glasner, and B. Virág, *The measurable Kesten theorem*, Ann. Probab. (2012), <http://arxiv.org/abs/1111.2080>.
- [2] L. Babai, *Automorphism groups, isomorphism, reconstruction*, Handbook of Combinatorics, vol. II, Elsevier, Amsterdam, 1995, pp. 1447–1540.

- [3] R. Bauerschmidt, H. Duminil-Copin, J. Goodman, and G. Slade, *Lectures on self-avoiding walks*, Probability and Statistical Physics in Two and More Dimensions (D. Ellwood, C. M. Newman, V. Sidoravicius, and W. Werner, eds.), Clay Mathematics Institute Proceedings, vol. 15, CMI/AMS publication, 2012, pp. 395–476.
- [4] I. Benjamini, *Euclidean vs graph metric*, Erdős Centennial (L. Lovász, I. Ruzsa, and V. T. Sós, eds.), Springer Verlag, Berlin, 2013, pp. 35–57.
- [5] I. Benjamini, A. Nachmias, and Y. Peres, *Is the critical percolation probability local?*, Probab. Th. Rel. Fields **149** (2011), 261–269.
- [6] S. R. Broadbent and J. M. Hammersley, *Percolation processes. I. Crystals and mazes*, Proc. Cambridge Philos. Soc. **53** (1957), 629–641.
- [7] M. M. Day, *Amenable semigroups*, Illinois J. Math. **1** (1957), 509–544.
- [8] J. Dodziuk, *Difference equations, isoperimetric inequality and transience of certain random walks*, Trans. Amer. Math. Soc. **284** (1984), 787–794.
- [9] H. Duminil-Copin and S. Smirnov, *The connective constant of the honeycomb lattice equals $\sqrt{2 + \sqrt{2}}$* , Ann. Math. **175** (2012), 1653–1665.
- [10] P. Flory, *Principles of Polymer Chemistry*, Cornell University Press, 1953.
- [11] R. I. Grigorchuk, *On Burnside’s problem on periodic groups*, Funktsional. Anal. i Prilozhen. **14** (1980), 53–54.
- [12] ———, *Degrees of growth of finitely generated groups and the theory of invariant means*, Izv. Akad. Nauk SSSR Ser. Mat. **48** (1984), 939–985.
- [13] ———, *An example of a finitely presented amenable group that does not belong to the class EG*, Mat. Sbornik **189** (1998), 79–100, (transl., Sbornik Math. **189** (1998), 75–95).
- [14] ———, *Solved and unsolved problems around one group*, Infinite Groups: Geometric, Combinatorial and Dynamic Aspects, Progress in Mathematics, vol. 248, Springer Verlag, Berlin, 2005, pp. 117–218.
- [15] G. R. Grimmett, *Percolation*, 2nd ed., Springer Verlag, Berlin.
- [16] G. R. Grimmett, A. E. Holroyd, and Y. Peres, *Extendable self-avoiding walks*, Ann. de l’Inst. Henri Poincaré D **1** (2014), 61–75.
- [17] G. R. Grimmett and Z. Li, *Counting self-avoiding walks*, (2013), <http://arxiv.org/abs/1304.7216>.
- [18] ———, *Locality of connective constants, I. Transitive graphs*, (2014), <http://arxiv.org/abs/1412.0150>.
- [19] ———, *Strict inequalities for connective constants of regular graphs*, SIAM J. Disc. Math. **28** (2014), 1306–1333.
- [20] ———, *Bounds on connective constants of regular graphs*, Combinatorica **35** (2015), 279–294.
- [21] ———, *Locality of connective constants, II. Cayley graphs*, (2015), <http://arxiv.org/abs/1501.00476>.
- [22] G. R. Grimmett and D. J. A. Welsh, *Probability, an Introduction*, 2nd ed., Oxford University Press, Oxford, 2014.
- [23] M. Gromov, *Groups of polynomial growth and expanding maps*, Inst. Hautes Études Sci. Publ. Math. **53** (1981), 53–73.
- [24] J. M. Hammersley and W. Morton, *Poor man’s Monte Carlo*, J. Roy. Statist. Soc. B **16** (1954), 23–38.
- [25] J. M. Hammersley and D. J. A. Welsh, *Further results on the rate of convergence to the connective constant of the hypercubical lattice*, Quart. J. Math. Oxford **13** (1962), 108–110.

- [26] P. de la Harpe, *Topics in Geometric Group Theory*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, 2000.
- [27] I. N. Herstein, *Topics in Algebra*, 2nd ed., Xerox Corporation, Lexington, Mass., 1975.
- [28] G. Higman, *A finitely generated infinite simple group*, J. London Math. Soc. **26** (1951), 61–64.
- [29] J. A. Hillman, *The Algebraic Characterization of Geometric 4-Manifolds*, London Mathematical Society Lecture Note Series, vol. 198, Cambridge University Press, Cambridge, 1994.
- [30] ———, *Four-Manifolds, Geometries and Knots*, Geometry and Topology Monographs, vol. 5, 2002, available at <http://arxiv.org/abs/math/0212142>.
- [31] K. Juschenko and N. Monod, *Cantor systems, piecewise translations and simple amenable groups*, Ann. Math. **178** (2013), 775–787.
- [32] H. Kesten, *Full Banach mean values on countable groups*, Math. Scand. **7** (1959), 146–156.
- [33] ———, *Symmetric random walks on groups*, Trans. Amer. Math. Soc. **92** (1959), 336–354.
- [34] J. R. Lee and Y. Peres, *Harmonic maps on amenable groups and a diffusive lower bound for random walks*, Ann. Probab. **41** (2013), 3392–3419.
- [35] R. Lyons and Y. Peres, *Probability on Trees and Networks*, Cambridge University Press, Cambridge, 2016, <http://mypage.iu.edu/~rdlyons/>.
- [36] I. G. Lysënok, *A set of defining relations for the Grigorchuk group*, Mat. Zametki **38** (1985), 503–516, 634, (transl., Math. Notes **38** (1985), 784–792).
- [37] N. Madras and G. Slade, *Self-Avoiding Walks*, Birkhäuser, Boston, 1993.
- [38] S. Martineau and V. Tassion, *Locality of percolation for abelian Cayley graphs*, Ann. Probab. (2013), <http://arxiv.org/abs/1312.1946>.
- [39] J. von Neumann, *Zur allgemeinen Theorie des Masses*, Fund. Math. **13** (1929), 73–116.
- [40] I. Pak and T. Smirnova-Nagnibeda, *On non-uniqueness of percolation on non-amenable Cayley graphs*, C. R. Acad. Sci. Paris Sér. I Math. **330** (2000), 495–500.
- [41] N. Seifert, *On the girth of infinite graphs*, Discr. Math. **118** (1993), 279–287.
- [42] W. Woess, *Random Walks on Infinite Graphs and Groups*, Cambridge University Press, Cambridge, 2000.

STATISTICAL LABORATORY, CENTRE FOR MATHEMATICAL SCIENCES, CAMBRIDGE UNIVERSITY, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB, UK

E-mail address: g.r.grimmett@statslab.cam.ac.uk,

URL: <http://www.statslab.cam.ac.uk/~grg/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CONNECTICUT 06269-3009, USA

E-mail address: zhongyang.li@uconn.edu

URL: <http://www.math.uconn.edu/~zhongyang/>