Probability and Measure 3

Non-measurable sets. Several have asked how to prove the existence of non-Lebesgue sets. Let $I = [0,1]$, Define the equivalence relation $\sim$ by $x \sim y$ if $x - y$ is rational. Consider the equivalence classes $(C_j : j \in J)$. By the Axiom of Choice, we may find a set of representatives, say $R = (c_j : j \in J)$ where $c_j \in C_j$. Assume R is measurable (Borel or Lebesgue, say). Now $I = \bigcup_{q \in \mathbb{Q}} (q + R)$ where $\mathbb{Q}$ is the rationals and addition is done modulo 1. This is a countably-infinite disjoint union of sets with the same measure. By the countable additivity of Lebesgue measure λ, and the fact that $\lambda(I) = 1$, we can have neither $\lambda(R) = 0$ nor $\lambda(R) > 0$. Note the use of the Axiom of Choice.

Note. Unless otherwise specified, $(\Omega, \mathcal{F}, \mu)$ denotes a measure space, and $(\Omega, \mathcal{F}, \mathbb{P})$ a probability space.

1. Let $g : \mathbb{R} \to \mathbb{R}$ be convex in the sense that:

$$g\big(tx + (1-t)y\big) \leq tg(x) + (1-t)g(y), \qquad x, y \in \mathbb{R},\ 0 \leq t \leq 1.$$

Show that g has supporting tangents in the sense that: for $x \in \mathbb{R}$, there exists $\lambda \in \mathbb{R}$ such that $g(y) \geq g(x) + \lambda(y - x)$ for all y.

2. Prove that the space $L^\infty(\Omega, \mathcal{F}, \mu)$ is complete.

3. Let $p \geq 1$, and let f_n, f be measurable functions. Show that f_n converges to f in measure whenever $f_n \to f$ in L^p, but that the converse is not true.

4. Let X be a random variable, $p \geq 1$, and write $Y = |X|$. Show that

$$\mathbb{E}(Y^p) = \int_0^\infty p x^{p-1} \mathbb{P}(Y \geq x)\, dx.$$

(a) If $\mathbb{E}(Y^p) < \infty$, show that $\mathbb{P}(Y \geq x) = \mathrm{o}(x^{-p})$ as $x \to \infty$.
(b) If $\mathbb{P}(Y \geq x) = \mathrm{o}(x^{-p})$, show that $\mathbb{E}(Y^q) < \infty$ for $1 \leq q < p$.

5. Show that random variables X and Y on $(\Omega, \mathcal{F}, \mathbb{P})$ are independent if and only if, for all measurable $g, h : \mathbb{R} \to \mathbb{R}$ such that $g(X)$, $h(Y)$ are integrable,

$$\mathbb{E}(g(X)h(Y)) = \mathbb{E}(g(X))\mathbb{E}(h(Y)).$$

6. Let X be a random variable whose moment generating function $M(t) = \mathbb{E}(e^{tX})$ is finite on some neighbourhood of the origin. Show that

$$\mathbb{P}(X > x) \leq \inf_{t \in \mathbb{R}} \big\{ e^{-tx} M(t) \big\}, \qquad x \in \mathbb{R}.$$

7. (Continuation) Let $X_1, X_2, \ldots$ be independent identically distributed random variables with moment generating function $M(t)$ and mean value m. Assume that M is finite on some neighbourhood of the origin. Show that $S_n = \sum_{i=1}^n X_i$ satisfies

$$\mathbb{P}(S_n > na) \leq \Lambda(a)^n, \qquad a > m,$$

where $\Lambda(a) = \inf_{t\in\mathbb{R}}\{e^{-ta}M(t)\}$. Compute $\Lambda(a)$ when the X_i have the exponential distribution with parameter 1, and $a > 1$.

8. Let $(X_n : n \geq 1)$ be a sequence of identically distributed random variables in $L^2(\mathbb{P})$. Let $\epsilon > 0$. Show that $\mathbb{P}(|X_1| \geq \epsilon\sqrt{n}) = \mathrm{o}(n^{-1})$ as $n \to \infty$, and that

$$\frac{1}{\sqrt{n}} \max_{1\leq k\leq n} |X_k| \to 0 \qquad \text{in probability } \textbf{and} \text{ in } L^1.$$

9. Let $\mathcal{H}$ be a closed linear space of $L^2(\Omega, \mathcal{F}, \mu)$ with the L^2 norm $\|\cdot\| = \|\cdot\|_2$ and the inner product $\langle\cdot,\cdot\rangle$. Show the 'Pythagoras' and 'parallelogram rules',

$$\|f+g\|^2 = \|f\|^2 + 2\langle f, g\rangle + \|g\|^2,$$
$$\|f+g\|^2 + \|f-g\|^2 = 2(\|f\|^2 + \|g\|^2),$$

valid for $g, f \in \mathcal{H}$.

10. Let $V_1, V_2, \ldots$ be an increasing sequence of closed subspaces of $L^2(\Omega, \mathcal{F}, \mu)$ such that $V_i \subseteq V_{i+1}$ for all i. For $f \in L^2$, write f_n for the orthogonal projection of f onto V_n. Show that f_n converges in L^2.

11. Let V, W be closed subspaces of $L^2(\Omega, \mathcal{F}, \mathbb{P})$ such that $V \subseteq W$. Show that $\mathbb{E}(\mathbb{E}(X \mid W) \mid V) = \mathbb{E}(X \mid V)$ for $X \in L^2$.

12. Find a uniformly integrable sequence $(X_n : n \geq 1)$ such that $X_n \to 0$ a.s. and $\mathbb{E}(\sup_n |X_n|) = \infty$.

13. Let $g : [0,\infty) \to [0,\infty)$ be an increasing function such that $g(x)/x \to \infty$ as $x \to \infty$. Show that $(X_n : n \geq 1)$ is uniformly integrable if $\sup_n \mathbb{E}(g(|X_n|)) < \infty$.

14. Show that the sum $X_n + Y_n$ of two uniformly integrable sequences forms a uniformly integrable sequence.

15. (a) Suppose that $X_n \to X$ in L^r where $r \geq 1$. Show that $(|X_n|^r : n \geq 1)$ is uniformly integrable, and deduce that $\mathbb{E}(X_n^r) \to \mathbb{E}(X^r)$ if r is an integer.

(b) Conversely, suppose that $(|X_n|^r : n \geq 1)$ is uniformly integrable where $r \geq 1$, and show that $X_n \to X$ in L^r whenever $X_n \to X$ in probability.