The Mathematical Work of Evarist Giné

Vladimir Koltchinskii, Richard Nickl Sara van de Geer, Jon A. Wellner

Georgia Institute of Technology, University of Cambridge ETH Zürich, University of Washington

October 29, 2015

1 Introduction

Evarist Giné, or Evarist Giné-Masdéu, 1944-2015, was an influential, brilliant and prolific contributor to modern probability theory and mathematical statistics, with a focus on problems arising in infinite-dimensional settings. His work has had a profound impact on modern probability theory, mathematical statistics and recently also machine learning. This article constitutes an attempt to describe Evarist's major mathematical achievements, which we divide into separate subsections each of which are focused on a major area of work, including *probability in Banach spaces, empirical processes, the bootstrap, U-statistics and processes, and mathematical statistics.* We refer to the editorial of this memorial volume for a biographical summary of Evarist's life. A list of all of Evarist's publications, including his three books, can be found at the end of this article.

A unifying aspect of most of Evarist's work was the masterly combination of techniques from real and functional analysis with fundamental ideas from probability theory: Evarist had a deep knowledge of analytic techniques that he often applied with ingenious simplicity in probabilistic problems. At the same time Evarist was a versatile and classically trained probabilist who mastered several core areas ranging from limit theorems and inequalities for sums of independent random variables to Gaussian processes and martingale arguments.

2 The main building blocks of Evarist's work

2.1 The PhD thesis

Evarist wrote his PhD thesis under the supervision of Richard M. Dudley at the Massachusetts Institute of Technology (MIT).

Five remarkable papers came out of this thesis: Two of them ([1,2]) were published in the *Annals of Probability* and were side-products of Evarist's time as a graduate student. Paper [1] led Evarist into the area described in the next subsection, and paper [2] with R. Klein established limiting properties of the quadratic variation of processes with Gaussian increments generalising results by Dudley (1973) for Brownian motion.

But the main topic of Evarist's thesis was the construction of computable statistical tests for observations that take values in a general compact Riemannian manifold, published in the Annals of Statistics in 1975, see [4]. The handling editor for this substantial paper was Lucien Le Cam, who found the mathematical level so high that the only referee he could think of was Richard Dudley! This paper basically initiated the whole area of 'Sobolev tests' that has been important ever since in the area of directional statistics, and required the development of some important mathematical results of independent interest, such as a proof of the fact that a Sobolev ball defined on any compact Riemannian manifold satisfies the empirical central limit theorem (i.e., is a *P*-Donsker class – see Subsection 2.3 for more on this). At that time the general empirical process machinery was not yet available but Evarist succeeded with a clever reduction of the problem to the central limit theorem in C(S) (the space of continuous functions on a compact metric space S) using duality theory for Sobolev spaces. The thesis also required the proof of some exact identities in geometric analysis that were published in a separate paper [3], and led to a (and also Evarist's only) genuinely applied paper [5], published in the Journal of Geology. Overall it can be said that this thesis was absolutely outstanding in its breadth and depth.

2.2 Probability in Banach spaces

Departing from the paper [1] that was part of the PhD thesis, Evarist embarked on one of the hot topics in probability theory in the 1970s: the problem of deriving the classical limit theorems for sums $\sum_{i=1}^{n} X_i$ of centred i.i.d. random variables X_i that take values in an infinite-dimensional Banach space B: For instance the *central limit theorem* (CLT): for a suitable Gaussian random variable G taking values in B one wants to show the distributional limit theorem

$$\frac{1}{\sqrt{n}}\sum_{i=1}^{n} X_i \to^d G \quad \text{as} \quad n \to \infty.$$

While in finite-dimensional spaces a necessary and sufficient condition for the CLT to hold is finiteness of $E||X_1||^2$, in infinite dimensions this is not the case, and geometric properties of the Banach space start to play a key role. The mathematical techniques developed in this area during that time were of fundamental importance to many areas in modern mathematics, including geometric functional analysis, concentration of measure, statistical learning and mathematical statistics.

Evarist's contributions to the area were substantial, including about 20 papers and culminating in Evarist's first book *The central limit theorem for real and Banach-valued random variables* published in 1980. His work in the area, which was carried out jointly with several co-authors including Alejandro de Acosta, Aloisio Araujo and Joel Zinn, included the derivation of the Lévy-Khintchine formula for infinitely divisible laws in Banach spaces and related characterisations of the domain of attraction of a normal law in Banach spaces [12,14], convergence of moments in the CLT in Banach spaces [15], and CLTs for some specific spaces of functions [7,10, 16, 28, 29].

An elegant application of these techniques to the theory of *random sets* (see Kendall (1974), Matheron (1975)) is given in the paper [33] jointly with Marjorie

Hahn and Joel Zinn: If B is a separable Banach space, we can define the set K(B) of all non-empty compact subsets of B, which is a complete metric space when endowed with the Hausdorff distance δ . Moreover (Minkowski-) addition is a well-defined operation on K(B) and for $A \in K(B)$ we can even define a norm $||A|| = \sup\{||a||_B : a \in A\}$. A random compact set is any Borel random variable X taking values in K(B), and its expectation EX can be defined in a suitable ('Bochner'-) sense. Using a clever reduction to the CLT in C(S) with an appropriate choice of S, the paper [33] proved the central limit theorem for random sets: For instance, if $B = \mathbb{R}^d$, $E||X|| < \infty$ and if the X_i 's are i.i.d. random compact sets, then

$$\sqrt{n}\delta\left(\frac{1}{n}\sum_{i=1}^{n}X_{i}, EX\right)$$

converges in distribution to a suitable norm of a Gaussian process. A result for the case where B is infinite-dimensional is also proved.

2.3 Empirical Processes

Techniques for probability in Banach spaces were mostly developed for separable spaces, which is quite reasonable as probability distributions on complete metric spaces necessarily concentrate most of their mass on compact sets (i.e., they are Radon measures). At the same time, a key open problem in the late 1970s was the central limit theorem for *empirical processes* indexed by abstract classes \mathcal{F} of functions $f : S \to \mathbb{R}$, and where S is an arbitrary sample space where i.i.d. random variables X_1, \ldots, X_n of law P take their values.

i.i.d. random variables X_1, \ldots, X_n of law P take their values. The empirical measure $P_n := n^{-1} \sum_{j=1}^n \delta_{X_j}$ is a natural statistical estimator of the unknown law P and it is of importance to understand how close the sample means $P_n f = n^{-1} \sum_{j=1}^n f(X_j)$ are to the true means $Pf = \mathbb{E}f(X)$, uniformly over a large class \mathcal{F} of functions f. In the late 60s - early 70s, Vapnik and Chervonenkis, motivated by applications in statistical learning theory) obtained striking necessary and sufficient conditions for the uniform law of large numbers for empirical measures (the Glivenko-Cantelli problem)

$$\sup_{f \in \mathcal{F}} |P_n f - P f| \to 0 \text{ as } n \to \infty \text{ a.s.}$$

in the case when $\mathcal{F} = \{I_C : C \in \mathcal{C}\}\)$, where \mathcal{C} is a class of measurable subsets of S. They later extended these results to the classes of uniformly bounded functions (Vapnik and Chervonenkis (1981)). However, the extension of the celebrated 'Donsker theorem', that is, the central limit theorem for classical empirical processes on the real line, to the same general framework remained open. In today's notation the general empirical processes are written as

$$f \mapsto \nu_n(f) = n^{1/2}(P_n f - Pf) = \frac{1}{\sqrt{n}} \sum_{i=1}^n (f(X_i) - Ef(X)), \quad f \in \mathcal{F}_i$$

and the question is whether ν_n converges to some Gaussian process $(G_P(f) : f \in \mathcal{F})$ uniformly in $f \in \mathcal{F}$. Even if this looks at first like a very abstract problem, the techniques required for its solution have had the most profound impact on

modern theoretical statistics and learning theory. In a seminal paper Dudley (1978) studied such limit theorems, and showcased that even in the simplest cases where \mathcal{F} consists of indicators of a class \mathcal{C} of subsets of Euclidean space, techniques very different from those used in probability in Banach spaces are required. This is partly due to the fact that empirical processes typically do not concentrate on some Banach space of *continuous* (for some metric) functions on \mathcal{F} – think of the empirical distribution function where $\mathcal{F} = \{1_{(-\infty,t]} : t \in \mathbb{R}^d\}$ – and the resulting lack of separability of the Banach space $\ell^{\infty}(\mathcal{F})$ of bounded functions on \mathcal{F} in which ν_n takes values. Dudley (1978) studied the central limit theorem for empirical processes indexed by classes of sets calling such a class $\mathcal C$ Donsker for the law P of X_1, \ldots, X_n if the CLT holds on the class. He obtained sufficient conditions for Donsker property in terms of the bracketing metric entropy of \mathcal{C} and proved that Vapnik-Chervonenkis classes of sets (the notion also being introduced by Dudley) are Donsker for any law P subject only to proper measurability assumptions. After Dudley's key contribution some useful sets of sufficient conditions for the CLT for empirical processes to hold were given by V. Koltchinskii (1981) in terms of random entropies of function classes and by D. Pollard (1982) in terms of their uniform entropies. Le Cam (1983) obtained sufficient conditions for the CLT on classes of sets in terms of random entropies. These conditions were similar in spirit to Vapnik-Chervonenkis conditions for the laws of large numbers. The proofs of these results were based on what is now called symmetrization inequalities for tail probabilities. By a conditioning argument, they allowed to reduce the bounds on the sup-norms of empirical processes to the bounds for a subgaussian (conditionally on X_1, \ldots, X_n) process of the following type: $\mathcal{F} \ni f \mapsto n^{-1} \sum_{j=1}^n \varepsilon_j f(X_j)$, where $\{\varepsilon_j\}$ are i.i.d. Rademacher random variables independent of $\{X_j\}$. This subgaussian process (called nowadays the Rademacher process) was controlled in terms of entropies of random sets $\{(f(X_1), \ldots, f(X_n)) : f \in \mathcal{F}\} \subset \mathbb{R}^n$. Evarist's entrance to the stage of empirical processes could not have been more impressive, in a 70 page special invited paper [34] in The Annals of Probability in 1984, jointly written with Joel Zinn. This paper not only gave some very final results about the CLT for empirical processes, but also introduced powerful new techniques to the area and developed to perfection the techniques used before. In particular, the symmetrization method used by Koltchinskii (1981), Pollard (1982) and Le Cam (1983) has reached its final form in two beautiful Giné-Zinn symmetrization inequalities used ever since, and the reduction of bounding of the empirical process to bounding the Rademacher process has been studied to its full extent. Moreover, this technique was combined with other tools from probability in Banach spaces, such as the multiplier inequality of Pisier, and the paper established connections of the emerging theory of empirical processes with a large body of literature on probability in Banach spaces. The idea of Gaussian or Rademacher symmetrization via the multiplier inequality of Pisier was to play an important role in Evarist's later work, also with Zinn, on bootstrapping the empirical process; see subsection 2.4 below. Specific important results obtained in the Giné and Zinn (1984) paper include the final versions of random entropy conditions for the central limit theorem for empirical processes, the proof of necessity of such conditions for the classes of sets (the necessity of conditions in terms of Vapnik-Chervonenkis shattering numbers was proved later by Talagrand) and the extension of Vapnik-Chervonenkis necessary and sufficient condition for the law of large numbers (Glivenko-Cantelli theorem) to the case of unbounded classes of functions.

Another deep and beautiful result that Evarist obtained in empirical process theory is the Gaussian characterisation of uniform Donsker classes. Again in joint work with Joel Zinn, published in The Annals of Probability [53], Evarist asked the question: when does the CLT for the empirical process holds uniformly in the distribution P of the X_i 's? More precisely, if β is a metric for weak convergence in $\ell^{\infty}(\mathcal{F})$ and G the limiting Gaussian process, when is it true that

$$\sup_{P} \beta(\nu_n^P, G_P) \to 0 \text{ as } n \to \infty$$

where the supremum extends over all probability measures in P? Such a result is of key importance for the statistical interpretation of the CLT, as in its typical applications it is used to infer unknown properties of P and hence no prior knowledge of P should be required for its validity. In essence this is a question about the structure of the class \mathcal{F} , and \mathcal{F} is called uniform Donsker if the last limit holds true. The striking result of [53] is that a necessary and sufficient condition for the uniform Donsker property is that the limit G_P is pregaussian uniformly in P (which effectively means that the sample-continuity of the Gaussian process G_P in its argument $f \in \mathcal{F}$ for the intrinsic covariance metric holds uniformly in P). Therefore, to check whether a class \mathcal{F} is a uniform Donsker class is a problem that can be decided entirely in terms of properties of Gaussian processes, which is in remarkable contrast to the otherwise existing gap between the Donsker and the pregaussian property. If much of Evarist's work in empirical process theory was inspired by the idea of establishing powerful connections between empirical and (sub-)Gaussian processes, this result may be considered as a high-point of that program.

2.4 The Bootstrap

A fundamental idea in statistics, due to Efron (1979), is the resampling methodology known as the *bootstrap*. In can be used for inference and confidence sets in situations where limit distributions exist but are not accessible for the statistician (because they are complex or depend on unknown parameters). This can be illustrated in the situation where X_1, \ldots, X_n are i.i.d. random variables from law P with mean μ . Then we can draw at random from the sample values to create a bootstrap sample: Let X_{ni}^b , i = 1, ..., n, be i.i.d. draws of the random variable X_n^b with law $\mathbb{P}_n(X_n^b = X_i) = 1/n$ for every i = 1, ..., n. If $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n$ is the sample mean and $\bar{X}_n^b = \frac{1}{n} \sum_{i=1}^n X_{ni}^b$ is the mean of the resampled values, then the idea is that the (known, given the X_i 's) distribution of $\bar{X}_n^b = \bar{X}_n^b$. tion of $\bar{X}_n^b - \bar{X}_n$ is pivotal for the (unknown) distribution $\bar{X}_n - \mu$. Thus, when computing quantiles for the latter, we can resort to computing (approximate) quantiles of the former. That this works in many situations is fundamentally linked to the central limit theorem, and Evarist (together with Joel Zinn) made several substantial contributions to the field, among them [46] showing necessity of the sufficient conditions found in Bickel and Freeman (1981) and providing a general bootstrap CLT for empirical processes, that basically completely settles the question of 'consistency' of the (nonparametric) bootstrap. The key result of the 1990 Annals of Probability paper [51] with Joel Zinn is that, for ν_n^b the bootstrap empirical process,

$$\nu_n^b \to^d G_P$$
 (in probability) if and only if $\nu_n \to^d G_P$.

The 1990 paper remains a tour de force of techniques from empirical process theory and probability theory in Banach spaces in several respects, including: (a) providing another use of the "multiplier inequality" of Pisier which played a role with Gaussian multipliers in their 1984 paper, this time with (symmetrized) Poisson multipliers after applying Poissonization and symmetrization to the multinomial weights of Efron's bootstrap; (b) connecting the striking results of Ledoux and Talagrand (1986, 1988) concerning multiplier and conditional multiplier CLT's in Banach spaces to an important set of statistical questions.

Evarist, in collaboration with several co-authors (notably his late student, Miguel Arcones), also addressed a number of other issues in connection with bootstrap resampling methods. Arcones and Giné [47] show that the bootstrap of the sample mean of i.i.d. random variables with finite variance "works" if both the bootstrap sample size m_n and the original sample size go to infinity. In [54] they studied several bootstrap test of symmetry, and in [57] they showed how to bootstrap U- and V- statistics. The important issue of bootstrapping M-estimators and other smooth statistical functionals was addressed in [59]. This whole set of research directions and problems culminated in Evarist's St. Flour Lecture lotes on asymptotic theory for bootstrap resampling. These notes continue to serve an an important reference and touchstone for current research.

2.5 U-statistics and U-processes

The notion of U-statistic is a natural extension of the most classical object in probability, the sum of independent random variables. Given an i.i.d. sequence X_1, \ldots, X_n, \ldots of random variables in a measurable space S and a measurable function $h: S \times \cdots \times S \mapsto \mathbb{R}$ (a kernel), the U-statistic of order k is defined as

$$U_n(h) := \frac{(n-k)!}{n!} \sum_{i_1, \dots, i_k} h(X_{i_1}, \dots, X_{i_k})$$

with the sum being extended to all $1 \leq i_1, \ldots, i_k \leq n$ such that $i_l \neq i_{l'}, l \neq l'$. This notion originated in the work of Halmos (1946) on unbiased estimation and von Mises (1947) on expansions of smooth statistical functionals in the late 40s. U-statistics were formally introduced and studied by Hoeffding in 1948. Evarist started being interested in the asymptotic theory of U-statistics in the early 90s, when this theory was already relatively well developed, both in the case of U-statistics based on i.i.d. observations and in more general cases. Moreover, Nolan and Pollard (1987, 1988) initiated the study of U-processes (empirical processes of U-statistic structure indexed by their kernels). However, by the early 90s, many of these results had not reached the same degree of finality as the classical limit theorems for the sums of independent random variables and a number of hard and challenging problems remained open. Many of these problems were solved in the 90s in a series of papers by Evarist with several co-authors, including Joel Zinn, Miguel Arcones, Stanislaw Kwapien and Rafal Latala. The results obtained by Evarist and his collaborators included Marcinkiewicz type laws of large numbers for U-statistics (Giné and Zinn [61]) and the necessity of finiteness of the second moment and degeneracy of the kernel for the Central Limit Theorem for U-statistics (Giné and Zinn [65]). They also included important results on the central limit theorem for U-processes indexed by Vapnik-Chervonenkis classes of functions (Arcones and Giné [64])

and striking applications of these results to asymptotics of M-estimators based on U-statistics, in particular, a beautiful proof of asymptotic normality of the simplicial median (Arcones, Chen, and Giné [67]). These developments were based on new and powerful technical tools, such as the Hoffmann-Jørgensen inequality for U-processes (Giné and Zinn [61]) and decoupling inequalities by de la Peña (1992), de la Peña and Montgomery-Smith (1994, 1995). The method of decoupling became particularly important and it gave the name to the (1999) book 'Decoupling' written by Evarist with Victor de la Peña that remains to be the most important reference on the modern theory of U-statistics. However, one of the most spectacular results of this theory was obtained after this book was published. In several papers written in the later 90s, Evarist and his coauthors were trying to find a necessary and sufficient condition for the law of iterated logarithms (LILs) for degenerate U-statistics, a problem that happened to be extremely hard. The sufficiency of finiteness of the second moment of the kernel for the LIL was known since the late 80s (Dehling (1989)). Giné and Zhang [71] showed that there are degenerate kernels with infinite second moment for which the LIL does hold. They provided sufficient conditions on the kernel that did not imply the finiteness of the second moment, but these conditions were still not necessary. This challenging problem was solved for the second order U-statistics in a remarkable paper by Giné, Kwapien, Latala and Zinn [82] who proved the following result. Suppose X, Y, X_1, X_2, \ldots are i.i.d. random variables with values in a measurable space (S, \mathcal{A}) and let $h: S \times S \mapsto \mathbb{R}$ be a measurable symmetric kernel. Then,

$$\limsup_{n} \frac{1}{n \log \log n} \left| \sum_{1 \le i \ne j \le n} h(X_i, X_j) \right| < \infty \text{ a.s.}$$

if and only if the following conditions hold for some constant $C < \infty$:

(a) h is canonical (degenerate) for the law of X (that is, $\mathbb{E}h(X, y) = 0$ for almost all y)

(b) for all $u \ge 10$,

$$\mathbb{E}(h^2(X,Y) \wedge u) \le C \log \log u$$

(c) for some C > 0,

$$\sup \mathbb{E}\Big\{h(X,Y)f(X)g(Y): \max(\mathbb{E}f^2(X), \mathbb{E}g^2(X)) \le 1; f,g \in L^\infty\Big\} \le C.$$

The proof of this completely unexpected result was a masterpiece of technique and it relied on a variety of tools of U-statistics theory (many of them developed by the authors such as exponential bounds for Rademacher chaos due to Latala) and on rather sophisticated truncation arguments. A related result is a new and final version of a Bernstein type concentration inequality for Ustatistics (Giné, Latala and Zinn [84]) that is one of the most important and useful inequalities in this area of probability. This inequality was proved for Ustatistics of order 2, but later extended to higher orders by Adamczak (2006). Adamczak and Latala (2008) obtained necessary and sufficient conditions for bounded LIL for higher order U-statistics.

2.6 The asymptotic distribution of the *t*-statistic

Student's one-sample t-statistic

$$T_n = \frac{\sum_{i=1}^n X_i / n^{1/2}}{\left\{\sum_{i=1}^n (X_i - \overline{X}_n)^2 / (n-1)\right\}^{1/2}} = \frac{S_n / V_n}{\sqrt{\frac{n - (S_n / V_n)^2}{n-1}}}$$

where $S_n = \sum_{i=1}^n X_i$, $V_n = \sum_{i=1}^n X_i^2$ and X_1, \ldots, X_n are i.i.d. random variables, plays a key role in basic applied statistics. While its exact distribution is well-known under Gaussian sampling theory, it is important to understand the properties of T_n under non-Gaussian (or other non-standard) assumptions. Efron (1969) reviewed early studies of T_n under non-standard conditions (including Hotelling (1961), Hoeffding (1963), and others), and studied the limiting behavior of T_n and self-normalized sums. Logan, Mallows, Rice, and Sheep (1973) showed that if X is in the domain of attraction of an α -stable law, $0 < \alpha \leq 2$, centered if $\alpha > 1$ and symmetric if $\alpha = 1$, then $S_n/V_n \rightarrow_d Z_\alpha$ where Z_{α} is sub-Gaussian. They conjectured that " S_n/V_n is asymptotically normal if [and perhaps only if] X is in the domain of attraction of the normal law" and X is centered. The direct part of this conjecture follows fairly easily from standard results, see Maller (1981). The converse or "only if" part was proved in [72] in 1997 by Evarist in collaboration with David Mason and Friedrich Götze. This beautiful paper shows Evarist's complete mastery of the Paley-Zygmund inequality and the use thereof to show that if $\{S_n/V_n\}$ is stochastically bounded, then it is L_1 -bounded as well.

Evarist returned to this theme in at least two other papers: in [73] (with David Mason) he studied laws of the iterated logarithm for self-normalized sums; in [95] (with Friedrich Götze) he established asymptotic normality of multivariate t-statistics under non-standard conditions.

2.7 Nonparametric Statistics

In the 21st century, Evarist started to work on problems in nonparametric statistics, an area within mathematical statistics of high activity since the mid 90s. Evarist's interest was sparked by the broad applicability of empirical process tools to the area. A main insight was that Talagrand's (1996) deep inequality for empirical processes could be used with great effect, particularly to deal with problems that involve risk bounds in supremum-norm loss in density estimation, see the papers [87, 107]. For example consider a kernel estimator of the form

$$f_n(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - X_i}{h}\right), \quad X_i \sim i.i.d. \quad P,$$

with K a suitable kernel function. If we ignore the estimation 'bias', the uniform risk can be viewed as the empirical process

$$||f_n - Ef_n||_{\infty} = \frac{1}{h} \sup_{g \in \mathcal{G}} |(P_n - P)g|, \quad \mathcal{G} = \left\{g = K\left(\frac{x - \cdot}{h}\right) : x \in \mathbb{R}^d\right\}.$$

A first key idea used in the paper [87] is that under simple conditions on K, such as that it is of bounded variation, the class \mathcal{G} can be shown to be a Vapnik-Chervonenkis type class and hence, by the usual chaining argument and if P

has a bounded density,

$$E \sup_{g \in \mathcal{G}} |(P_n - P)g| \lesssim \sqrt{\frac{h \log(1/h)}{n}}$$

for relevant choices of h. Moreover, by Talagrand's inequality the concentration of $||f_n - Ef_n||_{\infty}$ around its expectation is effectively Gaussian (again for relevant choices of h), and this can be used for various things: Initially Evarist derived from it the exact almost sure limiting constant of $\sqrt{nh/\log(1/h)} \|f_n - Ef_n\|_{\infty}$ as $n \to \infty, h \to 0$, both for the kernel estimator [87] and for wavelet estimators [107], where a slightly different scaling is required. In later work [106,108]these exponential inequalities were shown to be of great use also to construct adaptive estimators of densities that successfully deal with the bias $||Ef_n - f||_{\infty}$ too (by applying Lepski's (1990) method). Related techniques were also used in the paper [97] to give empirical approximations (using the graph Laplacian) of the true Laplace operator on a Riemannian manifold (a result that has been used in the machine learning community), and also in the paper [113], where these concentration inequalities were used to give a new approach to derive contraction rates in Bayesian nonparametric function estimation, ideas which have since been used in Bayesian non-parametrics in the recent papers Ray (2013) and Nickl and Söhl (2015), among others.

In another influential paper [110] Evarist constructed adaptive confidence bands for unknown densities by deriving the exact Gumbel limit distribution of $||f_n - f||_{\infty}$, suitable scaled and centred, where f_n is a fully adaptive estimator (again based on Lepski's (1990) method). This was the first exact limiting distribution result for any adaptive estimator, and required a subtle use of Gaussian approximation techniques and limit theory for non-stationary Gaussian processes. Next to the probabilistic challenges this required the introduction of some new qualitative assumptions on f, which have now become known under the name of 'self-similarity', and which were shown in [110] to be effectively generic for Hölder spaces. These 'self-similarity' conditions have turned out to be more or less the 'right' conditions for the existence of adaptive nonparametric confidence sets, and have been further studied in several recent papers including Hoffmann and Nickl (2011), Chetverikov, Chernozhukov and Kato (2014) and the recent discussion paper by Szabó, van der Vaart and van Zanten (2015), all in the Annals of Statistics.

A further result that deserves mentioning here, and that is related to some techniques dating back to the famous 1984 paper [34] of Evarist with Joel Zinn, is the paper [103], where it is shown that certain pre-gaussian classes \mathcal{F} of functions exist that are a) not P-Donsker for certain P but for which b) the smoothed empirical process $\sqrt{n}(P_n * K_h - P)$ corresponding to a kernel density estimator does converge in distribution in $\ell_{\infty}(\mathcal{F})$ to the generalised Brownian bridge G_P – thus $P_n * K_h$ is strictly better than P_n in this case. These results have been instrumental in the recent study of statistical inference for the distribution function of Lévy measures and infinitely divisible distributions, see the articles Nickl and Reiß (2012) and Nickl, Reiß, Söhl and Trabs (2015).

The key relevance of probabilistic techniques for the foundations of nonparametric statistics have led to Evarist's third monograph, which is currently in press, of the title *Mathematical Foundations of Infinite-Dimensional Statistical Models*. It summarises and contains much of Evarist's work in the area, and demonstrates yet again the deep insight Evarist had into the mathematical foundations that underpin modern probability theory and statistics. In particular in the chapters on Gaussian and empirical processes in this book Evarist has left us a great intellectual monument that will be a reference for generations to come.

3 Publications of Evarist Giné I: Books

- Aloisio Araujo and Evarist Giné. The Central Limit Theorem for Real and Banach Valued Random Variables. John Wiley & Sons, New York-Chichester-Brisbane, 1980. Wiley Series in Probability and Mathematical Statistics.
- [2] Evarist Giné, Geoffrey R. Grimmett, and Laurent Saloff-Coste. Lectures on probability theory and statistics, volume 1665 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1997. Lectures from the 26th Summer School on Probability Theory held in Saint-Flour, August 19–September 4, 1996, Edited by P. Bernard.
- [3] Víctor H. de la Peña and Evarist Giné. Decoupling. From Dependence to Independence, Randomly Stopped Processes. U-statistics and Processes. Martingales and Beyond. Probability and its Applications (New York). Springer-Verlag, New York, 1999.
- [4] Evarist Giné and Richard Nickl. Mathematical Foundations of Infinite-Dimensional Statistical Models. Cambridge University Press, Cambridge UK, 2016.

4 Publications of Evarist Giné II: Articles

- Evarist Giné M. On the central limit theorem for sample continuous processes. Ann. Probability, 2:629–641, 1974.
- [2] Ruben Klein and Evarist Giné. On quadratic variation of processes with Gaussian increments. Ann. Probability, 3(4):716–721, 1975.
- [3] Evarist Giné M. The addition formula for the eigenfunctions of the Laplacian. Advances in Math., 18(1):102–107, 1975.
- [4] Evarist Giné M. Invariant tests for uniformity on compact Riemannian manifolds based on Sobolev norms. Ann. Statist., 3(6):1243–1266, 1975.
- [5] Richard Dudley, Priscilla C Perkins and Evarist Giné M. Statistical tests for preferred orientation. J. Geology, 83:685–705, 1975.

- [6] Evarist Giné M. Bounds for the speed of convergence in the central limit theorem in C(S). Z. Wahrscheinlichkeitstheorie und Verw. Gebiete, 36(4):317–331, 1976
- [7] Evarist Giné M. Some remarks on the central limit theorem in C(S). In Probability in Banach spaces (Proc. First Internat. Conf., Oberwolfach, 1975), pages 101–106. Lecture Notes in Math., Vol. 526. Springer, Berlin, 1976.
- [8] Ubaldo García-Palomares and Evarist Giné M. On the linear programming approach to the optimality property of Prokhorov's distance. J. Math. Anal. Appl., 60(3):596–600, 1977.
- [9] Evarist Giné. A survey on the general central limit problem in Banach spaces. In Séminaire sur la Géométrie des Espaces de Banach (1977– 1978), pages Exp. No. 24, 17. École Polytech., Palaiseau, 1978.
- [10] Evarist Giné and José R. León. On the central limit theorem in Hilbert space. Stochastica, 4(1):43–71, 1980.
- [11] Evarist Giné M. and José R. León. On the central limit theorem in Hilbert space. In Proceedings of the First World Conference on Mathematics at the Service of Man (Barcelona, 1977), Vol. I, pages 620–625. Univ. Politec., Barcelona, 1980.
- [12] Aloisio Araujo and Evarist Giné M. Type, cotype and Lévy measures in Banach spaces. Ann. Probab., 6(4):637–643, 1978.
- [13] Alejandro de Acosta, Aloisio Araujo, and Evarist Giné. On Poisson measures, Gaussian measures and the central limit theorem in Banach spaces. Advances in Probability IV, 1–68 (Probability on Banach Spaces). 1978.
- [14] Aloisio Araujo and Evarist Giné. On tails and domains of attraction of stable measures in Banach spaces. *Trans. Amer. Math. Soc.*, 248(1):105– 119, 1979.
- [15] Alejandro de Acosta and Evarist Giné. Convergence of moments and related functionals in the general central limit theorem in Banach spaces. Z. Wahrsch. Verw. Gebiete, 48(2):213–231, 1979.
- [16] Evarist Giné, V. Mandrekar, and Joel Zinn. On sums of independent random variables with values in L_p ($2 \le p < \infty$). In Probability in Banach spaces, II (Proc. Second Internat. Conf., Oberwolfach, 1978), volume 709 of Lecture Notes in Math., pages 111–124. Springer, Berlin, 1979.
- [17] Evarist Giné. Domains of attraction in Banach spaces. In Séminaire de Probabilités, XIII (Univ. Strasbourg, Strasbourg, 1977/78), volume 721 of Lecture Notes in Math., pages 22–40. Springer, Berlin, 1979.
- [18] Evarist Giné. Corrections to: "Domains of attraction in Banach spaces" [Séminaire de Probabilités, XIII (Univ. Strasbourg, Strasbourg, 1977/1978), pp. 24–40, Lecture Notes in Math., Vol. 721, Springer, Berlin, 1979; MR 80j:60018]. In Seminar on Probability, XIV (Paris, 1978/1979) (French), volume 784 of Lecture Notes in Math., page p. 17. Springer, Berlin, 1980.

- [19] Evarist Giné. Sums of independent random variables and sums of their squares. In Proceedings of the seventh Spanish-Portuguese conference on mathematics, Part III (Sant Feliu de Guíxois, 1980), number 22, pages 127–132, 1980.
- [20] Evarist Giné. Domains of partial attraction in several dimensions. Ann. Inst. H. Poincaré Sect. B (N.S.), 16(2):87–100, 1980.
- [21] Evarist Giné. Correction to: "Domains of partial attraction in several dimensions". Ann. Inst. H. Poincaré Sect. B (N.S.), 17(1):143–145, 1981.
- [22] Evarist Giné. The central limit problem: three less typical aspects. In Proceedings of the seventh Spanish-Portuguese conference on mathematics, Part III (Sant Feliu de Guíxois, 1980), number 22, pages 133–138, 1980.
- [23] Evarist Giné and Michael B. Marcus. On the central limit theorem in C(K). In Statistical and physical aspects of Gaussian processes (Saint-Flour, 1980), volume 307 of Colloq. Internat. CNRS, pages 361–383. CNRS, Paris, 1981.
- [24] Evarist Giné. Central limit theorems in Banach spaces: a survey. In Probability in Banach spaces, III (Medford, Mass., 1980), volume 860 of Lecture Notes in Math., pages 138–152. Springer, Berlin-New York, 1981.
- [25] Aloisio Araujo, Evarist Giné, V. Mandrekar, and Joel Zinn. On the accompanying laws theorem in Banach spaces. Ann. Probab., 9(2):202–210, 1981.
- [26] Evarist Giné and Michael B. Marcus. Some results on the domain of attraction of stable measures on C(K). Probab. Math. Statist., 2(2):125–147 (1983), 1982.
- [27] Evarist Giné and Michel B. Marcus. The central limit theorem for stochastic integrals with respect to Lévy processes. Ann. Probab., 11(1):58–77, 1983.
- [28] Evarist Giné and Joel Zinn. Central limit theorems and weak laws of large numbers in certain Banach spaces. Z. Wahrsch. Verw. Gebiete, 62(3):323– 354, 1983.
- [29] Evarist Giné. The Lévy-Lindeberg central limit theorem in L_p , 0 .Proc. Amer. Math. Soc., 88(1):147–153, 1983.
- [30] Evarist Giné and Marjorie G. Hahn. On stability of probability laws with univariate stable marginals. Z. Wahrsch. Verw. Gebiete, 64(2):157–165, 1983.
- [31] Evarist Giné. Large deviations in spaces of stable type. Ann. Inst. H. Poincaré Sect. B (N.S.), 19(3):267–279, 1983.
- [32] Evarist Giné. A counterexample on domains of partial attraction in Banach spaces. In Probability in Banach spaces, IV (Oberwolfach, 1982), volume 990 of Lecture Notes in Math., pages 102–111. Springer, Berlin-New York, 1983.

- [33] Evarist Giné, Marjorie G. Hahn, and Joel Zinn. Limit theorems for random sets: an application of probability in Banach space results. In *Probability in Banach spaces, IV (Oberwolfach, 1982)*, volume 990 of *Lecture Notes in Math.*, pages 112–135. Springer, Berlin, 1983.
- [34] Evarist Giné and Joel Zinn. Some limit theorems for empirical processes. Ann. Probab., 12(4):929–998, 1984. With discussion.
- [35] Evarist Giné and Marjorie G. Hahn. Characterization and domains of attraction of *p*-stable random compact sets. Ann. Probab., 13(2):447–468, 1985.
- [36] Evarist Giné, Michael B. Marcus, and Joel Zinn. A version of Chevet's theorem for stable processes. J. Funct. Anal., 63(1):47–73, 1985.
- [37] Evarist Giné and Marjorie G. Hahn. The Lévy-Hinčin representation for random compact convex subsets which are infinitely divisible under Minkowski addition. Z. Wahrsch. Verw. Gebiete, 70(2):271–287, 1985.
- [38] Evarist Giné and Joel Zinn. On the approximation of p-homogeneous functions in Banach spaces. In *Homenatge a Francesc Sales*, pp 78–87. Universitat de Barcelona, Barcelona 1985.
- [39] Evarist Giné and Marjorie G. Hahn. M-infinitely divisible random compact convex sets. In *Probability in Banach spaces*, V (Medford, Mass., 1984), volume 1153 of Lecture Notes in Math., pages 226–248. Springer, Berlin, 1985.
- [40] Evarist Giné and Joel Zinn. Lectures on the central limit theorem for empirical processes. In *Probability and Banach spaces (Zaragoza, 1985)*, volume 1221 of *Lecture Notes in Math.*, pages 50–113. Springer, Berlin, 1986.
- [41] Evarist Giné and Joel Zinn. Empirical processes indexed by Lipschitz functions. Ann. Probab., 14(4):1329–1338, 1986.
- [42] Evarist Giné and Joel Zinn. A remark on the central limit theorem for random measures and processes. In *Probability theory and mathematical* statistics, Vol. I (Vilnius, 1985), pages 483–487. VNU Sci. Press, Utrecht, 1987.
- [43] Evarist Giné and Joel Zinn. The law of large numbers for partial sum processes indexed by sets. Ann. Probab., 15(1):154–163, 1987.
- [44] Niels T. Andersen, Evarist Giné, Mina Ossiander, and Joel Zinn. The central limit theorem and the law of iterated logarithm for empirical processes under local conditions. *Probab. Theory Related Fields*, 77(2):271– 305, 1988.
- [45] Niels T. Andersen, Evarist Giné, and Joel Zinn. The central limit theorem for empirical processes under local conditions: the case of Radon infinitely divisible limits without Gaussian component. *Trans. Amer. Math. Soc.*, 308(2):603–635, 1988.

- [46] Evarist Giné and Joel Zinn. Necessary conditions for the bootstrap of the mean. Ann. Statist., 17(2):684–691, 1989.
- [47] Miguel A. Arcones and Evarist Giné. The bootstrap of the mean with arbitrary bootstrap sample size. Ann. Inst. H. Poincaré Probab. Statist., 25(4):457–481, 1989.
- [48] Evarist Giné and Joel Zinn. Discussion of D. Pollard's paper, Asymptotics via empirical processes. Statistical Science, 4:355-356. 1989.
- [49] Evarist Giné and Joel Zinn. L_p multipliers in the central limit theorem with p-stable limit. In Probability theory on vector spaces, IV (Lańcut, 1987), volume 1391 of Lecture Notes in Math., pages 74–81. Springer, Berlin, 1989.
- [50] Evarist Giné, Michael B. Marcus, and Joel Zinn. On random multipliers in the central limit theorem with *p*-stable limit, 0 . In*Probability* in Banach spaces 6 (Sandbjerg, 1986), volume 20 of Progr. Probab., pages120–149. Birkhäuser Boston, Boston, MA, 1990.
- [51] Evarist Giné and Joel Zinn. Bootstrapping general empirical measures. Ann. Probab., 18(2):851–869, 1990.
- [52] Evarist Giné, Marjorie G. Hahn, and Pirooz Vatan. Max-infinitely divisible and max-stable sample continuous processes. *Probab. Theory Related Fields*, 87(2):139–165, 1990.
- [53] Evarist Giné and Joel Zinn. Gaussian characterization of uniform Donsker classes of functions. Ann. Probab., 19(2):758–782, 1991.
- [54] Miguel A. Arcones and Evarist Giné. Some bootstrap tests of symmetry for univariate continuous distributions. Ann. Statist., 19(3):1496–1511, 1991.
- [55] R. M. Dudley, E. Giné, and J. Zinn. Uniform and universal Glivenko-Cantelli classes. J. Theoret. Probab., 4(3):485–510, 1991.
- [56] Miguel A. Arcones and Evarist Giné. Additions and correction to: "The bootstrap of the mean with arbitrary bootstrap sample size" [Ann. Inst. H. Poincaré Probab. Statist. 25 (1989), no. 4, 457–481; MR1045246 (91d:62019)]. Ann. Inst. H. Poincaré Probab. Statist., 27(4):583–595, 1991.
- [57] Miguel A. Arcones and Evarist Giné. On the bootstrap of U and V statistics. Ann. Statist., 20(2):655–674, 1992.
- [58] Miguel A. Arcones and Evarist Giné. On decoupling, series expansions, and tail behavior of chaos processes. J. Theoret. Probab., 6(1):101–122, 1993.
- [59] Miguel A. Arcones and Evarist Giné. On the bootstrap of *M*-estimators and other statistical functionals. In *Exploring the limits of bootstrap (East Lansing, MI, 1990)*, Wiley Ser. Probab. Math. Statist. Probab. Math. Statist., pages 13–47. Wiley, New York, 1992.

- [60] Evarist Giné and Joel Zinn. On Hoffmann-Jørgensen's inequality for Uprocesses. In Probability in Banach spaces, 8 (Brunswick, ME, 1991), volume 30 of Progr. Probab., pages 80–91. Birkhäuser Boston, Boston, MA, 1992.
- [61] Evarist Giné and Joel Zinn. Marcinkiewicz type laws of large numbers and convergence of moments for U-statistics. In Probability in Banach spaces, 8 (Brunswick, ME, 1991), volume 30 of Progr. Probab., pages 273–291. Birkhäuser Boston, Boston, MA, 1992.
- [62] Miguel A. Arcones and Evarist Giné. Limit theorems for U-processes. Ann. Probab., 21(3):1494–1542, 1993.
- [63] D. Alamayehu, Victor de la Peña and Evarist Giné. Bootstrap goodness of fit tests based on the empirical distribution function. Proc. Conf. on the Interface (M. Tarter and M. Look, editors): 228-233, 1993
- [64] Miguel A. Arcones and Evarist Giné. U-processes indexed by Vapnik-Červonenkis classes of functions with applications to asymptotics and bootstrap of U-statistics with estimated parameters. Stochastic Process. Appl., 52(1):17–38, 1994.
- [65] Evarist Giné and Joel Zinn. A remark on convergence in distribution of U-statistics. Ann. Probab., 22(1):117–125, 1994.
- [66] Evarist Giné-Masdeu. Empirical processes and applications: a general, biased view. (in Catalan) Bul. Soc. Catalana Mat., (9):7–34, 1994.
- [67] Miguel A. Arcones, Zhiqiang Chen, and Evarist Giné. Estimators related to U-processes with applications to multivariate medians: asymptotic normality. Ann. Statist., 22(3):1460–1477, 1994.
- [68] Jack Cuzick, Evarist Giné, and Joel Zinn. Laws of large numbers for quadratic forms, maxima of products and truncated sums of i.i.d. random variables. Ann. Probab., 23(1):292–333, 1995.
- [69] Miguel A. Arcones and Evarist Giné. On the law of the iterated logarithm for canonical U-statistics and processes. Stochastic Process. Appl., 58(2):217–245, 1995.
- [70] Evarist Giné. Empirical processes and applications: an overview. *Bernoulli*, 2(1):1–38, 1996. With a discussion by Jon A. Wellner and a rejoinder by the author.
- [71] Evarist Giné and Cun-Hui Zhang. On integrability in the LIL for degenerate U-statistics. J. Theoret. Probab., 9(2):385–412, 1996.
- [72] Evarist Giné, Friedrich Götze, and David M. Mason. When is the Student t-statistic asymptotically standard normal? Ann. Probab., 25(3):1514– 1531, 1997.
- [73] Evarist Giné and David M. Mason. On the LIL for self-normalized sums of IID random variables. J. Theoret. Probab., 11(2):351–370, 1998.

- [74] Evarist Gine. Lectures on some aspects of the bootstrap. In Lectures on probability theory and statistics (Saint-Flour, 1996), volume 1665 of Lecture Notes in Math., pages 37–151. Springer, Berlin, 1997.
- [75] Evarist Gine. Decoupling and limit theorems for U-statistics and U-processes. In Lectures on probability theory and statistics (Saint-Flour, 1996), volume 1665 of Lecture Notes in Math., pages 1–35. Springer, Berlin, 1997.
- [76] Vladimir Koltchinskii and Evarist Giné. Random matrix approximation of spectra of integral operators. *Bernoulli*, 6(1):113–167, 2000.
- [77] Evarist Giné and Jon A. Wellner. Uniform convergence in some limit theorems for multiple particle systems. *Stochastic Process. Appl.*, 72(1):47–72, 1997.
- [78] Evarist Giné. A consequence for random polynomials of a result of de la Peña and Montgomery-Smith. In *High dimensional probability (Oberwolfach, 1996)*, volume 43 of *Progr. Probab.*, pages 103–110. Birkhäuser, Basel, 1998.
- [79] Eustasio del Barrio, Evarist Giné, and Carlos Matrán. Central limit theorems for the Wasserstein distance between the empirical and the true distributions. Ann. Probab., 27(2):1009–1071, 1999.
- [80] Eustasio del Barrio, Evarist Giné, and Carlos Matrán. Correction: "Central limit theorems for the Wasserstein distance between the empirical and the true distributions" [Ann. Probab. 27 (1999), no. 2, 1009–1071; MR1698999 (2000g:60034)]. Ann. Probab., 31(2):1142–1143, 2003.
- [81] Evarist Giné and Armelle Guillou. Laws of the iterated logarithm for censored data. Ann. Probab., 27(4):2042–2067, 1999.
- [82] Evarist Giné, Stanislaw Kwapien, Rafal Latala, and Joel Zinn. The LIL for canonical U-statistics of order 2. Ann. Probab., 29(1):520–557, 2001.
- [83] Evarist Giné and Armelle Guillou. On consistency of kernel density estimators for randomly censored data: rates holding uniformly over adaptive intervals. Ann. Inst. H. Poincaré Probab. Statist., 37(4):503–522, 2001.
- [84] Evarist Giné, Rafal Latala, and Joel Zinn. Exponential and moment inequalities for U-statistics. In High dimensional probability, II (Seattle, WA, 1999), volume 47 of Progr. Probab., pages 13–38. Birkhäuser Boston, Boston, MA, 2000.
- [98] Evarist Giné, David Mason, and Jon A. Wellner (co-editors). High dimensional probability, II (Seattle, WA, 1999). Birkhauser Series Progress in Probability No. 47, 2000.
- [86] Evarist Giné, David M. Mason, and Andrei Yu. Zaitsev. The L₁-norm density estimator process. Ann. Probab., 31(2):719–768, 2003.

- [87] Evarist Giné and Armelle Guillou. Rates of strong uniform consistency for multivariate kernel density estimators. Ann. Inst. H. Poincaré Probab. Statist., 38(6):907–921, 2002. En l'honneur de J. Bretagnolle, D. Dacunha-Castelle, I. Ibragimov.
- [88] Evarist Giné, Vladimir Koltchinskii, and Joel Zinn. Weighted uniform consistency of kernel density estimators. Ann. Probab., 32(3B):2570–2605, 2004.
- [89] Evarist Giné and David M. Mason. The law of the iterated logarithm for the integrated squared deviation of a kernel density estimator. *Bernoulli*, 10(4):721–752, 2004.
- [90] Evarist Giné, Vladimir Koltchinskii, and Lyudmila Sakhanenko. Kernel density estimators: convergence in distribution for weighted sup-norms. *Probab. Theory Related Fields*, 130(2):167–198, 2004.
- [91] Evarist Giné, Vladimir Koltchinskii, and Lyudmila Sakhanenko. Convergence in distribution of self-normalized sup-norms of kernel density estimators. In *High dimensional probability*, *III (Sandjberg, 2002)*, volume 55 of *Progr. Probab.*, pages 241–253. Birkhäuser, Basel, 2003.
- [92] Evarist Giné, Vladimir Koltchinskii, and Jon A. Wellner. Ratio limit theorems for empirical processes. In *Stochastic inequalities and applications*, volume 56 of *Progr. Probab.*, pages 249–278. Birkhäuser, Basel, 2003.
- [93] Zhiqiang Chen and Evarist Giné. Another approach to asymptotics and bootstrap of randomly trimmed means. Ann. Inst. Statist. Math., 56(4):771–790, 2004.
- [98] Evarist Giné, Christian Houdré, and D. Nualart (co-editors). Stochastic Inequalities and Applications. Birkhauser, Basel, 2003.
- [95] Evarist Giné and Friedrich Götze. On standard normal convergence of the multivariate Student t-statistic for symmetric random vectors. *Electron. Comm. Probab.*, 9:162–171 (electronic), 2004.
- [96] Eustasio del Barrio, Evarist Giné, and Frederic Utzet. Asymptotics for L_2 functionals of the empirical quantile process, with applications to tests of fit based on weighted Wasserstein distances. *Bernoulli*, 11(1):131–189, 2005.
- [97] Evarist Giné and Vladimir Koltchinskii. Empirical graph Laplacian approximation of Laplace-Beltrami operators: large sample results. In *High dimensional probability*, volume 51 of *IMS Lecture Notes Monogr. Ser.*, pages 238–259. Inst. Math. Statist., Beachwood, OH, 2006.
- [98] Evarist Giné, Vladimir Koltchinskii, and Wenbo Li, and Joel Zinn (coeditors). High dimensional probability, Proceedings of the Fourth International Conference. IMS LNMS, Vol 51, Beachwood, Ohio. 2006.
- [99] Evarist Giné and Vladimir Koltchinskii. Concentration inequalities and asymptotic results for ratio type empirical processes. Ann. Probab., 34(3):1143–1216, 2006.

- [100] Evarist Giné and David M. Mason. On local U-statistic processes and the estimation of densities of functions of several sample variables. Ann. Statist., 35(3):1105–1145, 2007.
- [101] Evarist Giné and David M. Mason. Laws of the iterated logarithm for the local U-statistic process. J. Theoret. Probab., 20(3):457–485, 2007.
- [102] Evarist Giné and David M. Mason. Uniform in bandwidth estimation of integral functionals of the density function. Scand. J. Statist., 35(4):739– 761, 2008.
- [103] Evarist Giné and Richard Nickl. Uniform central limit theorems for kernel density estimators. Probab. Theory Related Fields, 141(3-4):333–387, 2008.
- [104] Evarist Giné and Richard Nickl. A simple adaptive estimator of the integrated square of a density. *Bernoulli*, 14(1):47–61, 2008.
- [105] Evarist Giné and Richard Nickl. Adaptation on the space of finite signed measures. Math. Methods Statist., 17(2):113–122, 2008.
- [106] Evarist Giné and Richard Nickl. An exponential inequality for the distribution function of the kernel density estimator, with applications to adaptive estimation. *Probab. Theory Related Fields*, 143(3-4):569–596, 2009.
- [107] Evarist Giné and Richard Nickl. Uniform limit theorems for wavelet density estimators. Ann. Probab., 37(4):1605–1646, 2009.
- [108] Evarist Giné and Richard Nickl. Adaptive estimation of a distribution function and its density in sup-norm loss by wavelet and spline projections. *Bernoulli*, 16(4):1137–1163, 2010.
- [109] Evarist Giné and Hailin Sang. Uniform asymptotics for kernel density estimators with variable bandwidths. J. Nonparametr. Stat., 22(5-6):773– 795, 2010.
- [110] Evarist Giné and Richard Nickl. Confidence bands in density estimation. Ann. Statist., 38(2):1122–1170, 2010.
- [111] Evarist Giné, C. S. Güntürk, and W. R. Madych. On the periodized square of L² cardinal splines. *Exp. Math.*, 20(2):177–188, 2011.
- [112] Evarist Giné, Vladimir Koltchinskii, and Rimas Norvaisa (co-editors). Selected Works of R. M. Dudley. Springer. 2010.
- [113] Evarist Giné and Richard Nickl. Rates of contraction for posterior distributions in L^r -metrics, $1 \leq r \leq \infty$. Ann. Statist., 39(6):2883–2911, 2011.
- [114] Evarist Giné and Hailin Sang. On the estimation of smooth densities by strict probability densities at optimal rates in sup-norm. In From probability to statistics and back: high-dimensional models and processes, volume 9 of Inst. Math. Stat. (IMS) Collect., pages 128–149. Inst. Math. Statist., Beachwood, OH, 2013.

[115] Evarist Giné and W. R. Madych. On wavelet projection kernels and the integrated squared error in density estimation. *Statist. Probab. Lett.*, 91:32– 40, 2014.

5 References

- ADAMCZAK, R. (2006). Moment inequalities for U-statistics. Ann. Probab. 34 2288–2314.
- [2] ADAMCZAK, R. and LATAŁA, R. (2008). The LIL for canonical U-statistics. Ann. Probab. 36 1023–1058.
- [3] BICKEL, P. J. and FREEDMAN, D. A. (1981). Some asymptotic theory for the bootstrap. *Ann. Statist.* **9** 1196–1217.
- [4] CHERNOZHUKOV, V., CHETVERIKOV, D. and KATO, K. (2014). Anticoncentration and honest, adaptive confidence bands. Ann. Statist. 42 1787–1818.
- [5] DE LA PEÑA, V. H. (1992). Decoupling and Khintchine's inequalities for U-statistics. Ann. Probab. 20 1877–1892.
- [6] DE LA PEÑA, V. H. and MONTGOMERY-SMITH, S. J. (1994). Bounds on the tail probability of U-statistics and quadratic forms. Bull. Amer. Math. Soc. (N.S.) 31 223–227.
- [7] DE LA PEÑA, V. H., MONTGOMERY-SMITH, S. J. and SZULGA, J. (1994). Contraction and decoupling inequalities for multilinear forms and U-statistics. Ann. Probab. 22 1745–1765.
- [8] DEHLING, H. (1989). The functional law of the iterated logarithm for von Mises functionals and multiple Wiener integrals. J. Multivariate Anal. 28 177–189.
- [9] DUDLEY, R. M. (1973). Sample functions of the Gaussian process. Ann. Probability 1 66–103.
- [10] DUDLEY, R. M. (1978). Central limit theorems for empirical measures. Ann. Probab. 6 899–929 (1979).
- [11] EFRON, B. (1969). Student's t-test under symmetry conditions. J. Amer. Statist. Assoc. 64 1278–1302.
- [12] EFRON, B. (1979). Bootstrap methods: another look at the jackknife. Ann. Statist. 7 1–26.
- [13] HALMOS, P. R. (1946). The theory of unbiased estimation. Ann. Math. Statistics 17 34–43.
- [14] HOEFFDING, W. (1963). Probability inequalities for sums of bounded random variables. J. Amer. Statist. Assoc. 58 13–30.

- [15] HOFFMANN, M. and NICKL, R. (2011). On adaptive inference and confidence bands. Ann. Statist. 39 2383–2409.
- [16] HOTELLING, H. (1961). The behavior of some standard statistical tests under nonstandard conditions. In Proc. 4th Berkeley Sympos. Math. Statist. and Prob., Vol. I. Univ. California Press, Berkeley, Calif., 319–359.
- [17] KENDALL, D. G. (1974). Foundations of a theory of random sets. In Stochastic geometry (a tribute to the memory of Rollo Davidson). Wiley, London, 322–376.
- [18] KOLTCHINSKII, V. I. (1981). On the central limit theorem for empirical measures. *Theory of Probability and Mathematical Statistics* 24 71–82.
- [19] LE CAM, L. (1983). A remark on empirical measures. In A Festschrift for Erich L. Lehmann. Wadsworth Statist./Probab. Ser., Wadsworth, Belmont, CA, 305–327.
- [20] LEDOUX, M. and TALAGRAND, M. (1986). Conditions d'intégrabilité pour les multiplicateurs dans le TLC banachique. Ann. Probab. 14 916–921.
- [21] LEDOUX, M. and TALAGRAND, M. (1988). Un critère sur les petites boules dans le théorème limite central. Probab. Theory Related Fields 77 29–47.
- [22] LEPSKII, O. V. (1990). A problem of adaptive estimation in Gaussian white noise. *Teor. Veroyatnost. i Primenen.* 35 459–470.
- [23] LOGAN, B. F., MALLOWS, C. L., RICE, S. O. and SHEPP, L. A. (1973). Limit distributions of self-normalized sums. Ann. Probability 1 788–809.
- [24] MALLER, R. A. (1981). A theorem on products of random variables, with application to regression. Austral. J. Statist. 23 177–185.
- [25] MATHERON, G. (1975). Random sets and integral geometry. John Wiley & Sons, New York-London-Sydney.
- [26] NICKL, R. and REISS, M. (2012). A Donsker theorem for Lévy measures. J. Funct. Anal. 263 3306–3332.
- [27] NICKL, R., REISS, M., SÖHL, J. and TRABS, M. (2015). High frequency Donsker theorems for Lévy measures. *Probab. Theory Related Fields* to appear.
- [28] NICKL, R. and SÖHL, J. (2015). Nonparametric bayesian posterior contraction rates for discretely observed scalar diffusions. arXiv:1510.05526
- [29] NOLAN, D. and POLLARD, D. (1987). U-processes: rates of convergence. Ann. Statist. 15 780–799.
- [30] NOLAN, D. and POLLARD, D. (1988). Functional limit theorems for Uprocesses. Ann. Probab. 16 1291–1298.
- [31] POLLARD, D. (1982). A central limit theorem for empirical processes. J. Austral. Math. Soc. Ser. A 33 235–248.

- [32] RAY, K. (2013). Bayesian inverse problems with non-conjugate priors. *Electron. J. Stat.* 7 2516–2549.
- [33] SZABÓ, B., VAN DER VAART, A. W. and VAN ZANTEN, J. H. (2015). Frequentist coverage of adaptive nonparametric Bayesian credible sets (with discussion). Ann. Statist. 43 1391–1428.
- [34] TALAGRAND, M. (1996). New concentration inequalities in product spaces. Invent. Math. 126 505–563.
- [35] V. MISES, R. (1947). On the asymptotic distribution of differentiable statistical functions. Ann. Math. Statistics 18 309–348.
- [36] VAPNIK, V. N. and CHERVONENKIS, A. Y. (1981). Necessary and sufficient conditions for the uniform convergence of empirical means to their true values. *Teor. Veroyatnost. i Primenen.* 26 543–563.