

An isoperimetric inequality for the Wiener sausage

Yuval Peres* Perla Sousi†

September 9, 2011

Abstract

Let $(\xi(s))_{s \geq 0}$ be a standard Brownian motion in $d \geq 1$ dimensions and let $(D_s)_{s \geq 0}$ be a collection of open sets in \mathbb{R}^d . For each s , let B_s be a ball centered at 0 with $\text{vol}(B_s) = \text{vol}(D_s)$. We show that $\mathbb{E}[\text{vol}(\cup_{s \leq t}(\xi(s) + D_s))] \geq \mathbb{E}[\text{vol}(\cup_{s \leq t}(\xi(s) + B_s))]$, for all t . In particular, this implies that the expected volume of the Wiener sausage increases when a drift is added to the Brownian motion.

Keywords and phrases. Poisson point process, random walk, Brownian motion, coupling, rearrangement inequality.

MSC 2010 subject classifications. Primary 26D15, 60J65, Secondary 60D05, 60G55, 60G50.

1 Introduction

Let $(\xi(t))_{t \geq 0}$ be a standard Brownian motion in \mathbb{R}^d . For any open set $A \subset \mathbb{R}^d$, the corresponding Wiener sausage at time t is $\cup_{s \leq t}(\xi(s) + A)$.

Kesten, Spitzer and Whitman(1964) (see [4, p. 252]) proved that for $d \geq 3$, if $A \subset \mathbb{R}^d$ is an open set with finite volume, then the Wiener sausage satisfies

$$\frac{\mathbb{E}[\text{vol}(\cup_{s \leq t}(\xi(s) + A))]}{t} \rightarrow \text{Cap}(A) \text{ as } t \rightarrow \infty,$$

where $\text{Cap}(A)$ is the Newtonian capacity of the set A .

Pólya and Szëgo (see [10]) proved that for $d \geq 3$, among all open sets of fixed volume, the ball has the smallest Newtonian capacity. Thus, for any open set $A \subset \mathbb{R}^d$ of finite volume,

$$\mathbb{E}[\text{vol}(\cup_{s \leq t}(\xi(s) + A))] \geq (1 - o(1))\mathbb{E}[\text{vol}(\cup_{s \leq t}(\xi(s) + B))], \text{ as } t \rightarrow \infty, \quad (1.1)$$

where B is a ball with $\text{vol}(B) = \text{vol}(A)$.

This naturally raises the question whether (1.1) holds for fixed t without the $1 - o(1)$ factor. Our main result gives a positive answer in the more general setting where the set A is allowed to vary with time.

*Microsoft Research, Redmond, Washington, USA; peres@microsoft.com

†University of Cambridge, Cambridge, UK; p.sousi@statslab.cam.ac.uk

Theorem 1.1. *Let $(\xi(s))_{s \geq 0}$ be a standard Brownian motion in $d \geq 1$ dimensions and let $(D_s)_{s \geq 0}$ be open sets in \mathbb{R}^d . For each s , let $r_s > 0$ be such that $\text{vol}(\mathcal{B}(0, r_s)) = \text{vol}(D_s)$. Then for all t we have that*

$$\mathbb{E}[\text{vol}(\cup_{s \leq t} (\xi(s) + D_s))] \geq \mathbb{E}[\text{vol}(\cup_{s \leq t} \mathcal{B}(\xi(s), r_s))].$$

Our original motivation for Theorem 1.1 came from our joint work with A. Sinclair and A. Stauffer [8]. In [8] it was proved that in dimension 2 for any continuous function $f : \mathbb{R}_+ \rightarrow \mathbb{R}^2$,

$$\mathbb{E}[\text{vol}(\cup_{s \leq t} \mathcal{B}(\xi(s) + f(s), r))] \geq (1 - o(1))\mathbb{E}[\text{vol}(\cup_{s \leq t} \mathcal{B}(\xi(s), r))], \text{ as } t \rightarrow \infty$$

and it was conjectured that for all $d \geq 1$ and for any continuous function $f : \mathbb{R}_+ \rightarrow \mathbb{R}^d$,

$$\mathbb{E}[\text{vol}(\cup_{s \leq t} \mathcal{B}(\xi(s) + f(s), r))] \geq \mathbb{E}[\text{vol}(\cup_{s \leq t} \mathcal{B}(\xi(s), r))], \quad (1.2)$$

where $\mathcal{B}(x, r)$ stands for the open ball centered at x of radius r .

In dimension 1 this conjecture was shown in [8] to follow from the reflection principle. Theorem (1.1) above establishes (1.2) without requiring f to be continuous or even measurable, by taking $D_s = \mathcal{B}(f(s), r)$.

Remark 1.2. For any collection of open sets (D_s) the set $\cup_{s \leq t} (\xi(s) + D_s)$ is open, and hence Lebesgue measurable. Its volume, namely $\text{vol}(\cup_{s \leq t} (\xi(s) + D_s))$, is a random variable. This is explained at the end of the proof of Theorem 1.1 in Remark 3.2.

Remark 1.3. The Wiener sausage determined by a ball, i.e. $W(t) = \cup_{s \leq t} \mathcal{B}(\xi(s), r)$, has been studied extensively. Spitzer [11] obtained exact asymptotics as $t \rightarrow \infty$ for the expected volume of the Wiener sausage in 2 and 3 dimensions and Donsker and Varadhan [2] obtained exact asymptotics for the exponential moments of the volume. Kesten, Spitzer and Whitman (see [11] and [12]) proved laws of large numbers type results and Le Gall in [5] proved fluctuation results corresponding to these laws of large numbers.


Remark 1.4. In [6] (see also [3, Corollary 2.1]) a result analogous to (1.2) for random walks is proved, namely that the expected range of a lattice walk always increases when a drift is added to the walk. The proof of that result and our proof of Theorem 1.1 do not seem to yield each other.

Remark 1.5. A result analogous to (1.2) was proved in [9] for the Hausdorff dimension of the image and the graph of $\xi + f$, where f is a continuous function. Namely, in [9] it is proved that


$$\dim(\xi + f)[0, 1] \geq \max\{\dim \xi[0, 1], \dim f[0, 1]\} \text{ a.s.}$$

and similarly for the dimension of the graph.

To prove Theorem 1.1, we establish an analogous statement for a certain random walk, which we refer to as the ε -ball walk.



(a) Random walk sausage with squares



(b) Random walk sausage with discs

Proposition 1.6. *Let $\varepsilon > 0$ and let $(z_*(k))_{k \geq 0}$ be a random walk in \mathbb{R}^d starting from 0 and with transition kernel given by*

$$p(x, y) = \frac{\mathbf{1}(\|x - y\| < \varepsilon)}{\varepsilon^d \omega(d)}, \quad (1.3)$$

where $\omega(d)$ is the volume of the unit ball in \mathbb{R}^d . For any integer n and any collection of open sets $(U_k)_{k \geq 0}$ in \mathbb{R}^d we have that

$$\mathbb{E}[\text{vol}(\cup_{k=0}^n (z_*(k) + U_k))] \geq \mathbb{E}[\text{vol}(\cup_{k=0}^n \mathcal{B}(z_*(k), r_k))], \quad (1.4)$$

where r_k is such that $\text{vol}(\mathcal{B}(0, r_k)) = \text{vol}(U_k)$.

Remark 1.7. Proposition 1.6 can be extended to any random walk with transition density $p(x, y) = f(|x - y|)$ with f decreasing. Using that one can deduce an analog of Theorem 1.1 for Lévy processes with a radially decreasing transition density.

In the next section we first state a rearrangement inequality in Theorem 2.1, taken from [1, Theorem 2], and apply it to random walks on the sphere. Then we prove Proposition 1.6. In Section 3, using Proposition 1.6 and Donsker's invariance principle, we give the proof of Theorem 1.1.

In Section 4 we collect some easy convergence lemmas.

Finally, in Section 5 we conclude with some open questions and remarks.

2 Rearrangement inequalities and proof of Proposition 1.6

Let \mathbb{S} denote a sphere in d dimensions. We fix $x^* \in \mathbb{S}$. For a subset A of \mathbb{S} we define A^* to be a geodesic cap centered at x^* such that $\mu(A^*) = \mu(A)$, where μ is the surface area measure on the sphere. We call A^* the symmetric rearrangement of A .

The following theorem is a special case of Burchard and Schmuckenschläger [1, Theorem 2].

Theorem 2.1 ([1]). *Let A_1, \dots, A_n be Borel subsets of \mathbb{S} and let $\psi_{ij} : \mathbb{S} \times \mathbb{S} \rightarrow \mathbb{R}_+$ be nonincreasing functions of distance. Then we have the following inequality*

$$\begin{aligned} \int_{\mathbb{S}} \dots \int_{\mathbb{S}} \prod_{1 \leq i \leq n} \mathbf{1}(x_i \in A_i) \prod_{1 \leq i < j \leq n} \psi_{ij}(x_i, x_j) d\mu(x_1) \dots d\mu(x_n) \\ \leq \int_{\mathbb{S}} \dots \int_{\mathbb{S}} \prod_{1 \leq i \leq n} \mathbf{1}(x_i \in A_i^*) \prod_{1 \leq i < j \leq n} \psi_{ij}(x_i, x_j) d\mu(x_1) \dots d\mu(x_n). \end{aligned}$$

Let $\mathbb{S}_R \subset \mathbb{R}^{d+1}$ be the sphere of radius R in $d+1$ dimensions centered at 0, i.e.

$$\mathbb{S}_R = \{(x_1, \dots, x_{d+1}) \in \mathbb{R}^{d+1} : \sum_{i=1}^{d+1} x_i^2 = R^2\}. \quad (2.1)$$

Let $\varepsilon > 0$ and let $\tilde{\zeta}$ be a random walk on the sphere that starts from a uniform point on the surface of the sphere and has transition kernel given by

$$\psi(x, y) = \frac{\mathbf{1}(\rho(x, y) < \varepsilon)}{\mu(\mathcal{C}(x, \varepsilon))}. \quad (2.2)$$

Again μ stands for the surface area measure on the sphere, $\rho(x, y)$ stands for the geodesic distance between x and y and

$$\mathcal{C}(x, \varepsilon) = \{z \in \mathbb{S}_R : \rho(x, z) < \varepsilon\}.$$

We call $\tilde{\zeta}$ the ε -cap walk. For a collection $(\Theta_k)_{k \geq 0}$ of Borel subsets of \mathbb{S}_R we define

$$\tau^\Theta = \inf\{k \geq 0 : \tilde{\zeta}(k) \in \Theta_k\}.$$

Lemma 2.2. *Let $(\Theta_k)_{k \geq 0}$ be Borel subsets of the sphere, \mathbb{S}_R . Then for all n we have that*

$$\mathbb{P}(\tau^\Theta > n) \leq \mathbb{P}(\tau^{\mathcal{C}} > n), \quad (2.3)$$

where for each k we define \mathcal{C}_k to be a geodesic cap centered at $(0, \dots, 0, -R)$ such that $\mu(\mathcal{C}_k) = \mu(\Theta_k)$.

Proof. Using the Markov property, we write $\mathbb{P}(\tau^\Theta > n)$ as

$$\frac{1}{\mu(\mathbb{S})} \int_{\mathbb{S}_R} \dots \int_{\mathbb{S}_R} \prod_{i=1}^n \psi(x_{i-1}, x_i) \prod_{i=0}^n \mathbf{1}(x_i \in \Theta_i^c) d\mu(x_0) \dots d\mu(x_n).$$

Let $x^* = (0, \dots, 0, R)$. Since the transition kernel ψ is a nonincreasing function of distance, applying Theorem 2.1, we obtain that this last integral is bounded from above by

$$\int_{\mathbb{S}_R} \dots \int_{\mathbb{S}_R} \prod_{i=1}^n \psi(x_{i-1}, x_i) \prod_{i=0}^n \mathbf{1}(x_i \in A_i) d\mu(x_0) \dots d\mu(x_n),$$

where for each i , we have that A_i is a geodesic cap centered at x^* with $\mu(A_i) = \mu(\Theta_i^c)$. Writing $\mathcal{C}_i = A_i^c$, we have that \mathcal{C}_i is a geodesic cap centered at $(0, \dots, 0, -R)$ such that $\mu(\mathcal{C}_i) = \mu(\Theta_i)$ and hence proving the lemma. \square

It is intuitive and standard that as $R \rightarrow \infty$ the sphere \mathbb{S}_R tends to Euclidean space. The following two lemmas make this precise in the situation we need for the proof of Proposition 1.6. These lemmas are established in Section 4.

We define the projection mapping $\pi : \mathbb{S}_R \rightarrow \mathbb{R}^d$ via

$$\pi(x_1, \dots, x_{d+1}) = (x_1, \dots, x_d) \quad (2.4)$$

and the inverse map $\pi^{-1}(x_1, \dots, x_d) = (x_1, \dots, x_d, 0) \in \mathbb{S}_R$ such that $x_{d+1} \leq 0$.

Lemma 2.3. *Let $r, K > 0$. Then for all $\delta > 0$, there exists R_0 such that for all $x \in \mathcal{B}(0, K)$ we have that*

$$\mathcal{B}(x, r - \delta) \subset \pi(\mathcal{C}(\pi^{-1}(x), r)) \subset \mathcal{B}(x, r + \delta), \text{ for all } R > R_0. \quad (2.5)$$

Also for all $A \subset \pi^{-1}(\mathcal{B}(0, K))$ we have that

$$\mu(A) - \text{vol}(\pi(A)) \rightarrow 0, \text{ as } R \rightarrow \infty.$$

Lemma 2.4. *Let $L, n > 0$. Let z be an ε -ball walk in \mathbb{R}^d started from a uniform point in $\mathcal{B}(0, L + n\varepsilon)$. Let ζ be an ε -cap walk on \mathbb{S}_R started from a uniform point in the cap $\mathcal{C}(L) = \pi^{-1}(\mathcal{B}(0, L + n\varepsilon))$. Then there exists a coupling of z and $\pi(\zeta)$ such that*

$$\mathbb{P}(\pi(\zeta(k)) = z(k), \forall k = 0, \dots, n) \rightarrow 1 \text{ as } R \rightarrow \infty.$$

The rest of this section is devoted to the proof of Proposition 1.6.

Proof of Proposition 1.6. First note that if the sets (U_k) have infinite volume, then the inequality is trivially true. We will prove the theorem under the assumption that they are all bounded, since then by truncation we can get the result for any collection of unbounded open sets.

Let $L > 0$ be sufficiently large such that

$$\cup_{i=0}^n U_i \subset \mathcal{B}(0, L).$$

We now consider the ball $\mathcal{B}(0, L + n\varepsilon)$ in \mathbb{R}^d and we take R big enough so that we can ensure that

$$\mathcal{C}(L) = \pi^{-1}(\mathcal{B}(0, L + n\varepsilon)) \quad (2.6)$$

is a cap (centered at $(0, \dots, 0, -R)$) of geodesic radius bigger than $L + n\varepsilon$.

From Lemma 2.2 we have that

$$\mathbb{P}(\tau^{\pi^{-1}(U)} > n) \leq \mathbb{P}(\tau^{\mathcal{C}} > n), \quad (2.7)$$

where for each k we define \mathcal{C}_k to be a geodesic cap centered at $(0, \dots, 0, -R)$ such that $\mu(\mathcal{C}_k) = \mu(\pi^{-1}(U_k))$.

It is clear that

$$\mathbb{P}(\tau^{\pi^{-1}(U)} \leq n) = \mathbb{P}(\tilde{\zeta}(0) \in \mathcal{C}(L)) \mathbb{P}_{\mathcal{C}(L)}(\tau^{\pi^{-1}(U)} \leq n),$$

where under $\mathbb{P}_{\mathcal{C}(L)}$ the starting point of the random walk is uniform on $\mathcal{C}(L)$. Similarly for the collection of caps \mathcal{C} we have

$$\mathbb{P}(\tau^{\mathcal{C}} \leq n) = \mathbb{P}(\tilde{\zeta}(0) \in \mathcal{C}(L))\mathbb{P}_{\mathcal{C}(L)}(\tau^{\mathcal{C}} \leq n).$$

Hence, using the above equalities together with (2.7) we obtain that

$$\mathbb{P}(\forall k = 0, \dots, n, \pi(\zeta(k)) \notin U_k) \leq \mathbb{P}(\forall k = 0, \dots, n, \pi(\zeta(k)) \notin \pi(\mathcal{C}_k)), \quad (2.8)$$

where ζ is an ε -cap walk started from a uniform point in $\mathcal{C}(L)$, i.e., it has transition kernel given by (2.2). Now, we will go back to \mathbb{R}^d . Let $(z(k))_k$ be an ε -ball walk that starts from a uniform point in $\mathcal{B}(0, L + n\varepsilon)$, i.e., it has transition kernel given by (1.3). Thus for each k we can write $z(k) = z(0) - z_*(k)$, where $z(0)$ is uniform on $\mathcal{B}(0, L + n\varepsilon)$ and $(z_*(k))$ is an ε -ball walk that starts from 0. We define

$$T^U = \min\{k \geq 0 : z(k) \in U_k\}. \quad (2.9)$$

Since $z(0)$ is uniform on $\mathcal{B}(0, L + n\varepsilon)$, we can write

$$\begin{aligned} \mathbb{P}(T^U > n) &= \mathbb{P}(z(0) \notin U_0, z(1) \notin U_1, \dots, z(n) \notin U_n) \\ &= \mathbb{P}(z(0) \notin (\cup_{k=0}^n (z_*(k) + U_k))) = 1 - \frac{\mathbb{E}[\text{vol}(\cup_{k=0}^n (z_*(k) + U_k))]}{\text{vol}(\mathcal{B}(0, L + n\varepsilon))}. \end{aligned}$$

It is easy to check that for each k the projection $\pi(\mathcal{C}_k)$ is a ball in \mathbb{R}^d centered at 0. Let r_k^R be its radius and let r_k be such that $\text{vol}(\mathcal{B}(0, r_k)) = \text{vol}(U_k)$. Then by Lemma 2.3 we have that

$$r_k^R \rightarrow r_k, \text{ as } R \rightarrow \infty. \quad (2.10)$$

From Lemma 2.4, there exists a coupling of $(\pi(\zeta(k)))_{k=0, \dots, n}$ with $(z(k))_{k=0, \dots, n}$, so that

$$\mathbb{P}(\forall k = 0, \dots, n : \pi(\zeta(k)) = z(k)) \rightarrow 1 \text{ as } R \rightarrow \infty. \quad (2.11)$$

By the union bound we have that

$$\mathbb{P}(\forall k = 0, \dots, n : z(k) \notin U_k) \leq \mathbb{P}(\forall k = 0, \dots, n : \pi(\zeta(k)) \notin U_k) + \mathbb{P}(\text{coupling fails}), \quad (2.12)$$

where *coupling fails* means that there exists $k \in \{0, \dots, n\}$ such that $\pi(\zeta(k)) \neq z(k)$. Thus, from (2.11) we have

$$\mathbb{P}(\text{coupling fails}) = 1 - \mathbb{P}(\forall k = 0, \dots, n : \pi(\zeta(k)) = z(k)) \rightarrow 0 \text{ as } R \rightarrow \infty. \quad (2.13)$$

Similarly,

$$\mathbb{P}(\forall k = 0, \dots, n : z(k) \notin \pi(\mathcal{C}_k)) \geq \mathbb{P}(\forall k = 0, \dots, n : \pi(\zeta(k)) \notin \pi(\mathcal{C}_k)) - \mathbb{P}(\text{coupling fails}). \quad (2.14)$$

Hence, (2.12) and (2.14) together with (2.8) give that

$$\mathbb{P}(T^{\pi(\mathcal{C})} > n) \geq \mathbb{P}(\forall k, z(k) \notin U_k) - 2\mathbb{P}(\text{coupling fails}).$$

Therefore, using that $\pi(\mathcal{C}_k) = \mathcal{B}(0, r_k^R)$, we deduce that

$$1 - \frac{\mathbb{E}[\text{vol}(\cup_{k=0}^n (z_*(k) + \mathcal{B}(0, r_k^R)))]}{\text{vol}(\mathcal{B}(0, L + n\varepsilon))} \geq 1 - \frac{\mathbb{E}[\text{vol}(\cup_{k=0}^n (z_*(k) + U_k))]}{\text{vol}(\mathcal{B}(0, L + n\varepsilon))} - 2\mathbb{P}(\text{coupling fails}).$$

Using (2.10) and (2.13) and letting $R \rightarrow \infty$ in the inequality above concludes the proof of the proposition. \square

3 Proof of Theorem 1.1

For $n \in \mathbb{N}$ and $t > 0$ we define

$$\mathcal{D}_{n,t} = \left\{ \frac{kt}{2^n} : k = 0, \dots, 2^n \right\}.$$

Lemma 3.1. *Let $(\xi(s))_{s \geq 0}$ be a standard Brownian motion in $d \geq 1$ dimensions and let $(V_s)_{s \geq 0}$ be open sets in \mathbb{R}^d . For each s , let $r_s > 0$ be such that $\text{vol}(V_s) = \text{vol}(\mathcal{B}(0, r_s))$. Then for all $n \in \mathbb{N}$ and $t > 0$ we have that*

$$\mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_{n,t}} (\xi(\ell) + V_\ell) \right) \right] \geq \mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_{n,t}} \mathcal{B}(\xi(\ell), r_\ell) \right) \right].$$

Proof. First note that if the sets (V_s) have infinite volume, then the inequality is trivially true. We will prove the theorem under the assumption that they are all bounded, since then by truncation we can get the result for any collection of unbounded open sets.

Let X_1, X_2, \dots be i.i.d. random variables with uniform distribution on $\mathcal{B}(0, 1)$. Write $S_n = \sum_{i=1}^n X_i$. Let $N \in \mathbb{N}$ and let σ^2 be the variance of X_1 . Then for any collection of bounded open sets (U_s) we have from Proposition 1.6 that

$$\mathbb{E} \left[\text{vol} \left(\cup_{s \in \{0, 1/N, \dots, [t]/N\}} \left(\frac{S_{[Ns]}}{\sqrt{\sigma^2 N}} + U_s \right) \right) \right] \geq \mathbb{E} \left[\text{vol} \left(\cup_{s \in \{0, 1/N, \dots, [t]/N\}} \left(\frac{S_{[Ns]}}{\sqrt{\sigma^2 N}} + \mathcal{B}(0, r_s) \right) \right) \right], \quad (3.1)$$

where r_s is such that $\text{vol}(U_s) = \text{vol}(\mathcal{B}(0, r_s))$.

We will drop the dependence on t from $\mathcal{D}_{n,t}$ to simplify the notation. Since (3.1) holds true for any collection of bounded sets, we have that for N large enough

$$\mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_n} \left(\frac{S_{[N\ell]}}{\sqrt{\sigma^2 N}} + V_\ell \right) \right) \right] \geq \mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_n} \left(\frac{S_{[N\ell]}}{\sqrt{\sigma^2 N}} + \mathcal{B}(0, r_\ell) \right) \right) \right],$$

since in (3.1) we can take $U_\ell = V_\ell$, for $\ell \in \mathcal{D}_n$ and empty otherwise. As before, r_ℓ is such that $\text{vol}(\mathcal{B}(0, r_\ell)) = \text{vol}(V_\ell)$.

By Donsker's invariance principle (see for instance [7, Theorem 5.22]), for a fixed n , we have that

$$\left(\frac{S_{[N\ell]}}{\sqrt{\sigma^2 N}} \right)_{\ell \in \mathcal{D}_n} \xrightarrow{(w)} (\xi(\ell))_{\ell \in \mathcal{D}_n} \text{ as } N \rightarrow \infty. \quad (3.2)$$

If for all $\ell \in \mathcal{D}_n$ we have that $\text{vol}(\partial V_\ell) = 0$, then by (3.2) and Lemma 4.1 we deduce that

$$\mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + V_\ell) \right) \right] \geq \mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + \mathcal{B}(0, r_\ell)) \right) \right].$$

If $\exists k$ such that $\text{vol}(\partial V_k) > 0$, then for each $\ell \in \mathcal{D}_n$ we write $V_\ell = \cup_{j=1}^\infty A_{j,\ell}$, where $(A_{j,\ell})_j$ are all the dyadic cubes that are contained in V_ℓ .

Then for every finite K , we have that

$$\mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + \cup_{j=1}^K A_{j,\ell}) \right) \right] \geq \mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + \mathcal{B}(0, r(\ell, K))) \right) \right], \quad (3.3)$$

where for each ℓ we have that $r(\ell, K)$ satisfies $\text{vol}(\mathcal{B}(0, r(\ell, K))) = \text{vol}(\cup_{j=1}^K A_{j,\ell})$, and hence $r(\ell, K) \nearrow r_\ell$ as $K \rightarrow \infty$. Thus, letting $K \rightarrow \infty$, by monotone convergence we conclude that

$$\mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + V_\ell) \right) \right] \geq \mathbb{E} \left[\text{vol} \left(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + \mathcal{B}(0, r_\ell)) \right) \right].$$

□

Proof of Theorem 1.1. Let $M > 0$. We will show the theorem under the assumption that $D_s \subset \mathcal{B}(0, M)$, for all $s \leq t$, since then by monotone convergence we can get it for any collection of open sets.

For each s , we define the set

$$D_{s,n} = \{z \in D_s : d(z, D_s^c) > (t/2^n)^{1/3}\}, \quad (3.4)$$

which is open. For every $\ell \in \mathcal{D}_n$, we define

$$Z_\ell = \cup_{\ell \leq s < \ell + t/2^n} D_{s,n}, \quad (3.5)$$

which is again open as a union of open sets. For each s we let $r_{s,n}$ be such that $\text{vol}(\mathcal{B}(0, r_{s,n})) = \text{vol}(D_{s,n})$. From Proposition 1.6 we then get that

$$\mathbb{E} [\text{vol}(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + Z_\ell))] \geq \mathbb{E} [\text{vol}(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + \mathcal{B}(0, r_{\ell,n}^*)))] , \quad (3.6)$$

where $r_{\ell,n}^*$ satisfies

$$r_{\ell,n}^* = \left(\sup_{\ell \leq s < \ell + t/2^n} r_{s,n} - (t/2^n)^{1/3} \right)^+.$$

We now define the event

$$\Omega_n = \left\{ \forall h \leq t/2^n : \sup_{s,u:|s-u| \leq h} \|\xi(s) - \xi(u)\| \leq (2^{1/3} - 1)h^{1/3} \right\}.$$

We will now show that on Ω_n we have that

$$\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + Z_\ell) \subset \cup_{s \leq t} (\xi(s) + D_s).$$

Let $y \in \xi(\ell) + Z_\ell$, for some $\ell \in \mathcal{D}_n$. Then there exists $\ell \leq s < \ell + t/2^n$ such that $d(y - \xi(\ell), D_s^c) > (t/2^n)^{1/3}$. Also, since $\|\xi(s) - \xi(\ell)\| \leq (t/2^n)^{1/3}$ on Ω_n , we get by the triangle inequality that

$$d(y - \xi(s), D_s^c) \geq d(y - \xi(\ell), D_s^c) - \|\xi(s) - \xi(\ell)\| > (t/2^n)^{1/3} - (2^{1/3} - 1)(t/2^n)^{1/3} > 0,$$

and hence we deduce that $y \in \xi(s) + D_s$.

Therefore, from (3.6) we obtain

$$\mathbb{E} [\text{vol}(\cup_{s \leq t} (\xi(s) + D_s))] + \mathbb{E} [\text{vol}(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + Z_\ell)) \mathbf{1}(\Omega_n^c)] \geq \mathbb{E} [\text{vol}(\cup_{\ell \in \mathcal{D}_n} \mathcal{B}(\xi(\ell), r_{\ell,n}^*)) \mathbf{1}(\Omega_n)]. \quad (3.7)$$

Claim 3.1. *As $n \rightarrow \infty$ we have that a.s.*

$$\text{vol}(\cup_{\ell \in \mathcal{D}_n} \mathcal{B}(\xi(\ell), r_{\ell,n}^*)) \mathbf{1}(\Omega_n) \nearrow \text{vol}(\cup_{s \leq t} \mathcal{B}(\xi(s), r_s)).$$

Proof. Indeed it is clear that $\Omega_n \subset \Omega_{n+1}$ and on Ω_n we have that

$$\cup_{\ell \in \mathcal{D}_n} \mathcal{B}(\xi(\ell), r_{\ell,n}^*) \subset \cup_{\ell \in \mathcal{D}_{n+1}} \mathcal{B}(\xi(\ell), r_{\ell,n+1}^*). \quad (3.8)$$

To prove that, we need to show that for each $\ell \in \mathcal{D}_n$, the ball $\mathcal{B}(\xi(\ell), r_{\ell,n}^*)$ is contained in the right hand side of (3.8). We may assume that $r_{\ell,n}^* > 0$. Let y be such that $\|y - \xi(\ell)\| < r_{\ell,n}^*$. We set $\ell' = \ell + t/2^{n+1}$ and $\ell'' = \ell + t/2^n$. Then there are two cases:

- If $\sup_{\ell \leq s < \ell'} r_{s,n} \geq \sup_{\ell' \leq s < \ell''} r_{s,n}$, then $\|y - \xi(\ell)\| < r_{\ell,n+1}^*$.
- If $\sup_{\ell \leq s < \ell'} r_{s,n} < \sup_{\ell' \leq s < \ell''} r_{s,n}$, then

$$\begin{aligned} \|y - \xi(\ell')\| &\leq \|y - \xi(\ell)\| + \|\xi(\ell) - \xi(\ell')\| < r_{\ell,n}^* + (2^{1/3} - 1)(t/2^{n+1})^{1/3} \\ &= \sup_{\ell' \leq s < \ell + t/2^n} r_{s,n} - (t/2^n)^{1/3} + (t/2^{n+1})^{1/3}(2^{1/3} - 1) \leq r_{\ell,n+1}^*. \end{aligned}$$

For each n we define the set A_n as follows:

$$A_n = \begin{cases} \cup_{\ell \in \mathcal{D}_n} \mathcal{B}(\xi(\ell), r_{\ell,n}^*), & \text{on } \Omega_n \\ \emptyset, & \text{otherwise.} \end{cases}$$

We will now show that

$$\cup_n A_n = \cup_{s \leq t} \mathcal{B}(\xi(s), r_s) \quad \text{a.s.} \quad (3.9)$$

To prove that, we will first show that for each n and $\ell \in \mathcal{D}_n$ on Ω_n we have that $\mathcal{B}(\xi(\ell), r_{\ell,n}^*) \subset \cup_{s \leq t} \mathcal{B}(\xi(s), r_s)$. Again we may assume that $r_{\ell,n}^* > 0$. Let y be such that $\|y - \xi(\ell)\| < r_{\ell,n}^*$. For each $\varepsilon > 0$, there exists u such that $\ell \leq u < \ell + t/2^n$ and $r_{u,n} > \sup_{\ell \leq s < \ell + t/2^n} r_{s,n} - \varepsilon$. We have

$$\begin{aligned} \|y - \xi(u)\| &\leq \|y - \xi(\ell)\| + \|\xi(\ell) - \xi(u)\| \\ &< r_{\ell,n}^* + (2^{1/3} - 1)(t/2^n)^{1/3} < r_{u,n} + \varepsilon - (t/2^n)^{1/3} + (2^{1/3} - 1)(t/2^n)^{1/3} \end{aligned}$$

and by choosing ε small enough, we can make $\|y - \xi(u)\|$ smaller than $r_{u,n}$.

For the other inclusion we will show that for each m fixed on Ω_m we have that $\cup_{s \leq t} \mathcal{B}(\xi(s), r_s) \subset \cup_n A_n$. This will be enough since by Lévy's modulus of continuity theorem (see for instance [7, Theorem 1.14]) we have that $\mathbb{P}(\cup_m \Omega_m) = 1$. Let y be such that $\|y - \xi(s)\| < r_s$. Take $\varepsilon > 0$ small enough so that $\|y - \xi(s)\| < r_s - \varepsilon$. The sets $D_{s,n}$ defined in (3.4) are increasing in n . Since D_s is open, $D_s = \cup_n D_{s,n}$. Hence by monotone convergence, we get that $r_{s,n} \nearrow r_s$ as $n \rightarrow \infty$. Thus for n large enough, we get that $r_s \leq r_{s,n} + \varepsilon$. We now take n even larger (also larger than m) so that $\|y - \xi(s)\| < r_s - \varepsilon - 2^{1/3}(t/2^n)^{1/3}$. If we take $\ell \in \mathcal{D}_n$ such that $\ell \leq s < \ell + t/2^n$, then, using that the events (Ω_n) are increasing, we deduce that

$$\begin{aligned} \|y - \xi(\ell)\| &\leq \|y - \xi(s)\| + \|\xi(s) - \xi(\ell)\| \\ &\leq r_s - \varepsilon - 2^{1/3}(t/2^n)^{1/3} + (2^{1/3} - 1)(t/2^n)^{1/3} \leq r_{s,n} - (t/2^n)^{1/3} \leq r_{\ell,n}^*. \end{aligned}$$

Therefore we showed that $y \in \mathcal{B}(\xi(\ell), r_{\ell,n}^*)$ and this proves (3.9). Monotone convergence theorem completes the proof of Claim 3.1.

Finally by the dominated convergence theorem we get that

$$\mathbb{E} [\text{vol}(\cup_{\ell \in \mathcal{D}_n} (\xi(\ell) + Z_\ell)) \mathbf{1}(\Omega_n^c)] \rightarrow 0,$$

since $\mathbb{P}(\Omega_n^c) \rightarrow 0$ as $n \rightarrow \infty$, by Lévy's modulus of continuity theorem and also

$$\text{vol}(\cup_{\ell \in \mathcal{D}_n} (\xi_\ell + Z_\ell)) \leq \text{vol}(\mathcal{B}(0, \max_{s \leq t} |\xi(s)| + M)),$$

since we assumed that the sets (D_s) are all contained in $\mathcal{B}(0, M)$.

Taking the limit as $n \rightarrow \infty$ in (3.7) concludes the proof. \square

Remark 3.2. The proofs of Lemma 3.1 and Theorem 1.1 also give that for any collection of open sets (D_s) , the volume of $\cup_{s \leq t} (\xi(s) + D_s)$ is a random variable. Indeed, on $\cup_n \Omega_n$ (which has $\mathbb{P}(\cup_n \Omega_n) = 1$) we have that

$$\text{vol}(\cup_{s \leq t} (\xi(s) + D_s)) = \lim_{n \rightarrow \infty} \text{vol}(\cup_{k=1}^n \cup_{\ell \in \mathcal{D}_k} (\xi(\ell) + Z_\ell)),$$

where Z_ℓ is as defined in (3.5). If for all n and all $\ell \in \mathcal{D}_n$ we have that $\text{vol}(\partial Z_\ell) = 0$, then from Lemma 4.1 we get the measurability. Otherwise, we write Z_ℓ as a countable union of the dyadic subcubes contained in it and then use the monotonicity property together with Lemma 4.1 again, like in the last part of the proof of Lemma 3.1.

4 Relating the sphere to its tangent plane

In this section we give the proofs of Lemmas 2.3 and 2.4. We also establish a useful lemma on continuity of volumes.

Proof of Lemma 2.3. Let c be a sufficiently large positive constant so that for R sufficiently large we have for all $x \in \mathcal{B}(0, K)$ that

$$\pi(\mathcal{C}(\pi^{-1}(x), r)), \mathcal{B}(x, r + \delta) \subset \mathcal{B}(0, K + c).$$

Let $x = (x_1, \dots, x_d) \in \mathcal{B}(0, K)$ and $y = (y_1, \dots, y_d) \in \pi(\mathcal{C}(\pi^{-1}(x), r))$. Then $\pi^{-1}(y) \in \mathcal{C}(\pi^{-1}(x), r)$ and hence

$$\theta(\pi^{-1}(x), \pi^{-1}(y)) \leq \frac{r}{R}, \tag{4.1}$$

where $\theta = \theta(\pi^{-1}(x), \pi^{-1}(y))$ satisfies

$$\cos \theta = \frac{\langle \pi^{-1}(x), \pi^{-1}(y) \rangle}{R^2} = \frac{\sum_{i=1}^{d+1} x_i y_i}{R^2}.$$

We write $\|\cdot\|$ for the Euclidean distance in \mathbb{R}^d . We then have

$$\|x - y\|^2 = \sum_{i=1}^d (x_i - y_i)^2 = 2R^2 - 2R^2 \cos \theta - (x_{d+1} - y_{d+1})^2.$$

But, since $\pi^{-1}(x), \pi^{-1}(y) \in \mathbb{S}_R$, we get that $x_{d+1} = \sqrt{R^2 - \sum_{i=1}^d x_i^2}$ and $y_{d+1} = \sqrt{R^2 - \sum_{i=1}^d y_i^2}$ and using the fact that $x \in \mathcal{B}(0, K)$ and $y \in \mathcal{B}(0, K + c)$ we obtain that

$$(x_{d+1} - y_{d+1}) \rightarrow 0 \text{ as } R \rightarrow \infty,$$

uniformly over all x and y in $\mathcal{B}(0, K)$ and $\mathcal{B}(0, K + c)$ respectively. From (4.1) and the monotonicity of the $\cos \varphi$ function for φ small, we obtain that

$$\cos \theta \geq \cos \frac{r}{R},$$

and hence for all R sufficiently large we have that

$$\|x - y\| \leq r + \delta.$$

To prove the other inclusion, it suffices to show that

$$\mathcal{B}(0, K + c) \cap \pi(\mathcal{C}(\pi^{-1}(x), r))^c \subset \mathcal{B}(0, K + c) \cap \mathcal{B}(x, r - \delta)^c, \quad (4.2)$$

since $\mathcal{B}(0, K + c)^c$ is disjoint from $\pi(\mathcal{C}(\pi^{-1}(x), r))$ and $\mathcal{B}(x, r - \delta)$. Finally (4.2) can be proved using similar arguments to the ones employed above.

To prove the last statement of the lemma, we write

$$\mu(A) - \text{vol}(\pi(A)) = \int_{\pi(A)} \left(\frac{1}{\cos(\alpha(x))} - 1 \right) dx,$$

where $\alpha(x)$ is the angle at the origin between $\pi^{-1}(x)$ and $(0, \dots, 0, -R)$.

Using that $\alpha(x) \rightarrow 0$ as $R \rightarrow \infty$ uniformly over all $x \in \mathcal{B}(0, K)$ we get the desired convergence. \square

Proof of Lemma 2.4. Since $\zeta(0)$ is uniformly distributed on $\mathcal{C}(L)$, we have that $\pi(\zeta(0))$ has density function given by

$$f_R^0(x) = \frac{\mathbf{1}(x \in \mathcal{B}(0, L + cn\varepsilon))}{\cos(\alpha(x))\mu(\mathcal{C}(L))},$$

where $\alpha(x)$ is the angle between $\pi^{-1}(x)$ and $(0, \dots, 0, -R)$. Also $z(0)$ has density function given by

$$f^0(x) = \frac{\mathbf{1}(x \in \mathcal{B}(0, L + cn\varepsilon))}{\text{vol}(\mathcal{B}(0, L + cn\varepsilon))}.$$

From Lemma 2.3 we get that $f_R(x) \rightarrow f(x)$ as $R \rightarrow \infty$ uniformly over all x , and hence the maximal coupling will succeed with probability tending to 1 as $R \rightarrow \infty$. So, for $\pi(\zeta(0))$ and $z(0)$ we have used the maximal coupling. Given $\pi(\zeta(0))$ and $z(0)$, then the density function of $\pi(\zeta(1))$ is given by

$$f_R^1(x) = \frac{\mathbf{1}(x \in \pi(\mathcal{C}(\zeta(0), \varepsilon)))}{\cos(\alpha(x))\mu(\mathcal{C}(\zeta(0), \varepsilon))}$$

and $\xi(1)$ has density function given by

$$f^1(x) = \frac{\mathbf{1}(x \in \mathcal{B}(\xi(0), \varepsilon))}{\text{vol}(\mathcal{B}(\xi(0), \varepsilon))}.$$

Thus if the coupling of the starting points has succeeded, then we can use the maximal coupling to couple the first steps and continuing this way we can couple the first n steps. Hence, we get that the probability that the coupling succeeds tends to 1 as $R \rightarrow \infty$. \square

Lemma 4.1. *Let $(A_i)_{i=1, \dots, n}$ be measurable sets in \mathbb{R}^d such that $\text{vol}(\partial A_i) = 0$, for all i . Then the function defined by*

$$x = (x_1, \dots, x_n) \mapsto \text{vol}(\cup_{i=1}^n (x_i + A_i))$$

is continuous.

Proof. Let A^δ stand for the δ -enlargement of the set A , i.e. $A^\delta = A + \mathcal{B}(0, \delta)$. If $\text{vol}(\partial A) = 0$, then it is easy to see that

$$\text{vol}(A^\delta) \rightarrow \text{vol}(A) \quad \text{as } \delta \rightarrow 0. \quad (4.3)$$

Let $x = (x_1, \dots, x_n)$ be such that $\|x\| < \delta$. We then have that

$$\text{vol}((\cup_{i=1}^n (x_i + A_i)) \triangle (\cup_{i=1}^n A_i)) \leq \text{vol}(\cup_{i=1}^n ((x_i + A_i) \triangle A_i)) \leq \sum_{i=1}^n \text{vol}((x_i + A_i) \triangle A_i).$$

But $\text{vol}((x_i + A_i) \triangle A_i) = 2(\text{vol}((x_i + A_i) \cup A_i) - \text{vol}(A_i))$ and $\text{vol}((x_i + A_i) \cup A_i) \leq \text{vol}(A_i^\delta)$, and hence by (4.3) we get the desired convergence. \square

5 Concluding Remarks and Questions

1. We recall here the detection problem as discussed in [8]. Let $\Pi = \{X_i\}$ be a Poisson point process in \mathbb{R}^d of intensity λ . We now let each point X_i of the Poisson process move according to an independent standard Brownian motion $(\xi_i(s))_{s \geq 0}$. Let u be another particle originally placed at the origin and which is moving according to a deterministic function f . We define the detection time of u analogously to (2.9) via

$$T_{\text{det}}^f = \inf\{s \geq 0 : \exists i \text{ s.t. } X_i + \xi_i(s) \in \mathcal{B}(f(s), r)\}.$$

Then from [8, Lemma 3.1] we have that

$$\mathbb{P}(T_{\text{det}}^f > t) = \exp(-\lambda \mathbb{E}[\text{vol}(\cup_{s \leq t} \mathcal{B}(\xi(s) + f(s), r))]).$$

In terms of the detection probabilities, Theorem 1.1 then gives that

$$\mathbb{P}(T_{\text{det}}^f > t) \leq \mathbb{P}(T_{\text{det}}^0 > t),$$

where T_{det}^0 stands for the detection time when u does not move at all. This means that the best strategy for u to stay undetected for long time is to stay put. This is an instance of the Pascal principle, which is discussed in [3] and [6] for a similar model in the discrete lattice.

2. Let (D_s) be a collection of open sets as in Theorem 1.1. We showed that

$$\mathbb{E}[\text{vol}(\cup_{s \leq t} (\xi(s) + D_s))] \geq \mathbb{E}[\text{vol}(\cup_{s \leq t} (\xi(s) + B_s))],$$

where the sets (B_s) are balls as defined in Theorem 1.1. Does the stochastic domination inequality

$$\mathbb{P}(\text{vol}(\cup_{s \leq t} (\xi(s) + D_s)) \geq \alpha) \geq \mathbb{P}(\text{vol}(\cup_{s \leq t} (\xi(s) + B_s)) \geq \alpha) \quad \forall \alpha$$

also hold?

3. Fix an open set D in \mathbb{R}^d . For any function $f : \mathbb{R}_+ \rightarrow \mathbb{R}^d$, consider the Wiener sausage with drift f determined by D , namely $W_f^D(t) = \cup_{s \leq t} (D + \xi(s) + f(s))$. For which sets D is it true that the expected volume of W_f^D is minimized when $f = 0$?

4. Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}^d$ be a measurable function. Consider the convex hull of $C_f(t) = (\xi + f)[0, t]$. Is the expected volume of $C_f(t)$ minimized when $f = 0$?

Acknowledgements

We are grateful to Philippe Charmoy, Jason Miller, Alexandre Stauffer and Rongfeng Sun for useful comments and discussions.

References

- [1] A. Burchard and M. Schmuckenschläger. Comparison theorems for exit times. *Geom. Funct. Anal.*, 11(4):651–692, 2001.
- [2] M. D. Donsker and S. R. S. Varadhan. Asymptotics for the Wiener sausage. *Comm. Pure Appl. Math.*, 28(4):525–565, 1975.
- [3] A. Drewitz, J. Gärtner, A.F. Ramírez, and R. Sun. Survival probability of a random walk among a poisson system of moving traps, 2010. arXiv:1010.3958v1.
- [4] Kiyosi Itô and Henry P. McKean, Jr. *Diffusion processes and their sample paths*. Springer-Verlag, Berlin, 1974. Second printing, corrected, Die Grundlehren der mathematischen Wissenschaften, Band 125.
- [5] Jean-François Le Gall. Fluctuation results for the Wiener sausage. *Ann. Probab.*, 16(3):991–1018, 1988.
- [6] M. Moreau, G. Oshanin, O. Bénichou, and M. Coppey. Lattice theory of trapping reactions with mobile species. *Phys. Rev.*, E 69, 2004.
- [7] P. Mörters and Y. Peres. *Brownian Motion*. Cambridge University Press, 2010.
- [8] Y. Peres, A. Sinclair, P. Sousi, and A. Stauffer. Mobile geometric graphs: Detection, Coverage and Percolation, 2010. arXiv:1008.0075v2.
- [9] Y. Peres and P. Sousi. Brownian motion with variable drift: 0-1 laws, hitting probabilities and Hausdorff dimension, 2010. arXiv:1010.2987v1.
- [10] G. Pólya and G. Szegő. Inequalities for the capacity of a condenser. *Amer. J. Math.*, 67:1–32, 1945.
- [11] Frank Spitzer. Electrostatic capacity, heat flow, and Brownian motion. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 3:110–121, 1964.
- [12] Walter William Whitman. *Some Strong Laws for Random Walks and Brownian Motion*. ProQuest LLC, Ann Arbor, MI, 1964. Thesis (Ph.D.)—Cornell University.